

# Tracking and Prediction of Arterial Blood Pressure From Childhood to Young Adulthood in 868 Patients With Type 1 Diabetes

A multicenter longitudinal survey in Germany and Austria

INA KNERR, MD, PHD<sup>1</sup>  
 AXEL DOST, MD, PHD<sup>2</sup>  
 RUDOLF LEPLER, MD<sup>3</sup>  
 KLEMENS RAILE, MD<sup>4</sup>  
 EDITH SCHOBER, MD, PHD<sup>5</sup>  
 WOLFGANG RASCHER, MD, PHD<sup>1</sup>

REINHARD W. HOLL, MD, PHD<sup>6</sup>  
 ON BEHALF OF THE DIABETES DATA  
 ACQUISITION SYSTEM FOR PROSPECTIVE  
 SURVEILLANCE (DPV) SCIENTIFIC  
 INITIATIVE GERMANY AND AUSTRIA

**OBJECTIVE** — Arterial blood pressure was followed in 868 patients with type 1 diabetes aged 6.0–19.9 years in 95 centers in Germany and Austria.

**RESEARCH DESIGN AND METHODS** — European blood pressure reference data for 28,043 children and adolescents were used with respect to age and sex. Data were stratified into three groups: prepubertal, pubertal, and postpubertal.

**RESULTS** — Up to 4% of the participants in the younger age-groups and 13.9% of the postpubertal patients exhibited blood pressure values >97th centile. Blood pressure levels correlated with A1C level and BMI Z score. Tracking of blood pressure revealed that children with elevated blood pressure had higher blood pressure in adolescence and young adulthood.

**CONCLUSIONS** — Patients with higher blood pressure in childhood showed elevated blood pressure later in life. We need to focus on the diagnosis of hypertension in children with type 1 diabetes and to study the efficacy of early intervention.

*Diabetes Care* 31:726–727, 2008

The objective of this survey was to follow arterial blood pressure and the prevalence of hypertension as a risk factor for cardiovascular disease (CVD) in a large cohort of young patients with type 1 diabetes. A total of 868 patients with type 1 diabetes from 6.0 to 19.9 years of age, who were treated in 95 Diabetes Centers and Pediatric Care Clinics in Germany and Austria, formed the study cohort for this report.

## RESEARCH DESIGN AND METHODS

At each of the first through fourth annual clinical visits, BMI

(calculated as weight in kilograms divided by the square of height in meters), A1C, and blood pressure at rest were recorded. A1C measurements were standardized to the Diabetes Control and Complications Trial reference of 4.05–6.05% (1); blood pressure levels were measured on a single occasion in a relaxed, sitting position at the upper arm with proper cuff size using sphygmomanometer or semi-automated Dinamap (Critikon, Tampa, FL). Data were collected from 1977 to 2006 with informed consent according to the Declaration of Helsinki, using the diabetes data

acquisition system for prospective surveillance (DPV), as previously described (2,3). Of 1,353 patients who had been screened, 962 were enrolled, and data sets from childhood to young adulthood were completed for 868 participants (96% Caucasian and 4% other ethnicity; age at diagnosis  $5.9 \pm 2.4$  years; 432 female and 436 male). Patients with other diseases or permanent medication, including anti-hypertensive drugs, were excluded. European blood pressure reference data for 28,043 children and adolescents were used with respect to age and sex (4). German reference data for BMI, obtained from 17,275 female subjects and 17,147 male subjects in a comparable time span, were applied as reported earlier (3,5). Blood pressure and BMI values were derived using the least median of squares (LMS) Box-Cox power transformation method, which adjusts the distribution of the parameters for skewness and allows individual data to be expressed as SDS or Z score (6,7). Data were stratified in three groups according to age: 6.0–9.9 years (prepubertal), 10.0–15.9 years (pubertal), and 16.0–19.9 years (postpubertal). Statistical analysis was performed using Pearson correlation, Kruskal-Wallis test, and Wilcoxon's signed-rank test. A P value of <0.05 was considered significant. In addition, mixed multivariate models with systolic or diastolic blood pressure during adulthood as the dependent variable and blood pressure during childhood (6–9.9 years), sex, migration background, age at onset, current age, observation period, BMI, smoking status, and treatment center (random effect) as potential confounders were evaluated (SAS proc glimmix).

**RESULTS** — Within these three age-groups, mean  $\pm$  SD values for A1C were  $7.4 \pm 1.4$ ,  $7.9 \pm 1.3$ , and  $8.4 \pm 1.7\%$ ; for BMI Z score  $0.24 \pm 0.73$ ,  $0.37 \pm 0.78$ , and  $0.60 \pm 0.89$ ; for systolic blood pressure (SBP)  $106 \pm 7$ ,  $116 \pm 8$ , and  $127 \pm$

From the <sup>1</sup>Children's and Adolescents' Hospital, University of Erlangen-Nuremberg, Erlangen, Germany; <sup>2</sup>University Children's Hospital, Jena, Germany; <sup>3</sup>Children's Hospital Wilhelmstift, Hamburg, Germany; <sup>4</sup>University Children's Hospital, Berlin, Germany; <sup>5</sup>University Children's Hospital, Vienna, Austria; and the <sup>6</sup>Division of Applied Information Technology, University of Ulm, Ulm, Germany.


Address correspondence and reprint requests to Ina Knerr, MD, PhD, Children's and Adolescents' Hospital, University of Erlangen-Nuremberg, D-91054 Erlangen, Germany. E-mail: ina.knerr@uk-erlangen.de. Received for publication 19 July 2007 and accepted in revised form 24 December 2007.

Published ahead of print at <http://care.diabetesjournals.org> on 9 January 2008. DOI: 10.2337/dc07-1392.

**Abbreviations:** CVD, cardiovascular disease; DBP, diastolic blood pressure; SBP, systolic blood pressure.

© 2008 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.



**Figure 1**—Tracking of systolic (A) and diastolic (B) blood pressure (BP) from childhood to young adulthood in patients with type 1 diabetes. Prepubertal (6.0–9.9 years) blood pressure Z score quartiles are given together with mean pubertal (10.0–15.9 years) and postpubertal (16.0–19.9 years) blood pressure Z scores in 868 patients with type 1 diabetes (of whom 432 were female and 436 male). Patients were stratified according to their prepubertal blood pressure Z scores in three categories: blood pressure Z score <25th centile ( $n = 217$ ), 25th–75th centile ( $n = 438$ ), and >75th centile ( $n = 213$ ).

11 mmHg; and for diastolic blood pressure (DPB)  $65 \pm 6$ ,  $68 \pm 7$ , and  $72 \pm 7$  mmHg. Mean blood pressure Z scores in the three age-groups increased with age (increases of 0.09, 0.12, and 0.52 for SBP,  $P < 0.000001$ , and 0.63 to 0.69 for DBP,  $P < 0.0001$ ). In the prepubertal and pubertal age-groups, up to 4% of participants exhibited blood pressure values >97th centile. However, 13.9% of patients in the postpubertal group had blood pressure levels >97th centile. Blood pressure values correlated to A1C level and BMI Z score ( $r = 0.2148$  and  $0.3663$ , respectively;  $P < 0.0001$ ).

A mixed model with adult blood pressure as the dependent variable and adjustment for sex, migration background, age at onset, current age, observation period,

BMI, smoking status, and treatment center (random effect) revealed significant effects of childhood blood pressure (age 6–9.9 years) as evidence for tracking ( $P < 0.000001$  for SBP and DBP) (Fig. 1). Elevation of childhood blood pressure by 1 SD increases adult blood pressure Z score by 0.43 (systolic) or 0.38 (diastolic).

**CONCLUSIONS**— In this survey, we followed blood pressure in patients with type 1 diabetes from childhood to young adulthood. Because patients with higher blood pressure in childhood, or even with hypertension, showed elevated blood pressure later in life, early intervention is feasible. The use of blood pressure determinations in patients with diabetes can, therefore, increase the prediction of a future CVD risk from childhood. It has been shown earlier, using ambulatory profiles of 24-h blood pressure, that daytime and nocturnal blood pressure is more pronounced in the course of type 1 diabetes compared with that in healthy control subjects (8). Although we can only imagine the beneficial effects of lowering blood pressure in the hypertensive individuals, preventing the development of atherosclerosis early in life is mandatory because morbidity and mortality of CVD are increased up to 10-fold in patients with type 1 diabetes (9). In our study, blood pressure values correlated significantly with A1C level and BMI Z score, which also exhibit adverse longitudinal changes in patients at risk for CVD (1,10). Similarly, progression of carotid intima-media thickness, a measure of atherosclerosis, is strongly associated with age, blood pressure, and A1C in subjects with type 1 diabetes (1).

Moreover, by showing the advantages of having lower blood pressure levels early in life on blood pressure levels in young adulthood, this study provides additional evidence for the importance of monitoring blood pressure as a risk factor in young patients with type 1 diabetes. In conclusion, we need to focus on the early detection of hypertension in children with type 1 diabetes and to study the efficacy of treatment in affected individuals early in life.

## References

- Nathan DM, Lachin J, Cleary P, Orchard T, Brillon DJ, Backlund JY, O'Leary DH, Genuth S; Diabetes Control and Complications Trial; Epidemiology of Diabetes Interventions and Complications Research Group: Intensive diabetes therapy and carotid intima-media thickness in type 1 diabetes mellitus. *N Engl J Med* 348:2294–2303, 2003
- Schwab KO, Doerfer J, Hecker W, Grulich-Henn J, Wiemann D, Kordonouri O, Beyer P, Holl RW, the DPV Initiative of the German Working Group for Pediatric Diabetology: Spectrum and prevalence of atherogenic risk factors in 27,358 children, adolescents, and young adults with type 1 diabetes: cross-sectional data from the German diabetes documentation and quality management system (DPV). *Diabetes Care* 29:218–225, 2006
- Knerr I, Wolf J, Reinehr T, Stachow R, Grabert M, Schober E, Rascher W, Holl RW: The 'accelerator hypothesis': relationship between weight, height, body mass index and age at diagnosis in a large cohort of 9,248 German and Austrian children with type 1 diabetes mellitus. *Diabetologia* 48:2501–2504, 2005
- de Man SA, Andre JL, Bachmann H, Grobbee DE, Ibsen KK, Laaser U, Lippert P, Hofman A: Blood pressure in childhood: pooled findings of six European studies. *J Hypertens* 9:109–114, 1991
- Krohmeier-Hauschild K, Wabitsch M, Kunze D, Geller F, Geiß HC, Hesse V, von Hippel A, Jaeger U, Johnson D, Korte W, Menner K, Müller G, Müller JM, Niemann-Pilatus A, Remer T, Schaefer F, Wittchen H-U, Zabransky S, Zellner K, Ziegler A, Hebebrand J: [Percentiles of body mass index in children and adolescents evaluated from different regional German studies]. *Monatsschr Kinderheilkd* 149:807–818, 2001 [in German]
- Cole TJ, Freeman JV, Preece MA: Body mass index reference curves for the UK, 1990. *Arch Dis Child* 73:25–29, 1995
- Jackson LV, Thalange NK, Cole TJ: Blood pressure centiles for Great Britain. *Arch Dis Child* 92:298–303, 2007
- Holl RW, Pavlovic M, Heinze E, Thon A: Circadian blood pressure during the early course of type 1 diabetes: analysis of 1,011 ambulatory blood pressure recordings in 354 adolescents and young adults. *Diabetes Care* 22:1151–1157, 1999
- Dahl-Jorgensen K, Larsen JR, Hanssen KF: Atherosclerosis in childhood and adolescent type 1 diabetes: early disease, early treatment? *Diabetologia* 48:1445–1453, 2005
- Srinivasan SR, Myers L, Berenson GS: Changes in metabolic syndrome variables since childhood in prehypertensive and hypertensive subjects: the Bogalusa Heart Study. *Hypertension* 48:33–39, 2006