

Corneal Confocal Microscopy Detects Early Nerve Regeneration After Pancreas Transplantation in Patients With Type 1 Diabetes

SANJAY MEHRA, MD¹
MITRA TAVAKOLI, MSC²
PANAGIOTIS A. KALLINIKOS, PHD²
NATHAN EFRON, PHD³

ANDREW J.M. BOULTON, MD²
TITUS AUGUSTINE, MD¹
RAYAZ A. MALIK, MD²

OBJECTIVE — Corneal confocal microscopy (CCM) is a rapid, noninvasive, clinical examination technique that quantifies small nerve fiber pathology. We have used it to assess the neurological benefits of pancreas transplantation in type 1 diabetic patients.

RESEARCH DESIGN AND METHODS — In 20 patients with type 1 diabetes undergoing simultaneous pancreas and kidney transplantation (SPK) and 15 control subjects, corneal sensitivity was evaluated using noncontact corneal esthesiometry, and small nerve fiber morphology was assessed using CCM.

RESULTS — Corneal sensitivity (1.54 ± 0.28 vs. 0.77 ± 0.02 , $P < 0.0001$), nerve fiber density (NFD) (13.8 ± 2.1 vs. 42 ± 3.2 , $P < 0.0001$), nerve branch density (NBD) (4.04 ± 1.5 vs. 26.7 ± 2.5 , $P < 0.0001$), and nerve fiber length (NFL) (2.23 ± 0.2 vs. 9.69 ± 0.7 , $P < 0.0001$) were significantly reduced, and nerve fiber tortuosity (NFT) (15.7 ± 1.02 vs. 19.56 ± 1.34 , $P = 0.04$) was increased in diabetic patients before pancreas transplantation. Six months after SPK, 15 patients underwent a second assessment and showed a significant improvement in NFD (18.04 ± 10.48 vs. 9.25 ± 1.87 , $P = 0.001$) and NFL (3.60 ± 0.33 vs. 1.84 ± 0.33 , $P = 0.002$) with no change in NBD (1.38 ± 0.74 vs. 1.38 ± 1.00 , $P = 1.0$), NFT (15.58 ± 1.20 vs. 16.30 ± 1.19 , $P = 0.67$), or corneal sensitivity (1.23 ± 0.39 vs. 1.54 ± 0.42 , $P = 0.59$).

CONCLUSIONS — Despite marked nerve fiber damage in type 1 diabetic patients undergoing pancreas transplantation, small fiber repair can be detected within 6 months of pancreas transplantation using CCM. CCM is a novel noninvasive clinical technique to assess the benefits of therapeutic intervention in human diabetic neuropathy.

Diabetes Care 30:2608–2612, 2007

Somatic polyneuropathy is one of the most common long-term complications of diabetes and is the main initiating factor for foot ulceration and lower extremity amputation (1,2). As 80% of amputations are preceded by foot ulceration, an effective means of detecting and treating peripheral neuropathy would have a major medical, social, and eco-

nomonic impact. With the exception of optimal glycemic control, there are currently no licensed treatments that prevent, slow, or arrest the development of neuropathy (1). The development of new treatments is of paramount importance, but they are hampered by a lack of clinically relevant surrogate end points favored by regulatory authorities (1). We have relied on

tests that quantify predominantly large nerve fiber dysfunction, which were principally developed to aid diagnosis and not to assess nerve repair and hence a therapeutic response (3). Thus, nerve conduction studies are useful but only detect an abnormality in large myelinated nerve fibers, whereas thermal and pain thresholds assess thinly myelinated (A δ) and unmyelinated (C) fiber function. Heart rate variability during respiratory stimuli indicates parasympathetic vagal efferent function, and blood pressure change during orthostatic manipulation evaluates sympathetic vasomotor efferents.

Although these tests correlate with axonal loss (4–7), there are major shortcomings when they are used to define therapeutic efficacy in clinical intervention trials (8). These tests do not target the specific fiber types that may benefit from the therapeutic intervention and demonstrate a limited ability to detect regeneration and repair. Only sural nerve biopsy with electron microscopy (9,10) and the assessment of intraepidermal nerve fiber density using skin-punch biopsies (11,12) directly assess nerve damage and repair; however, both are invasive procedures.

No available therapy (including glycemic control) has previously been shown to result in an improvement in diabetic neuropathy. Even in the most dramatic example of “curing” type 1 diabetes with pancreas transplantation, in 115 patients followed over 10 years, neurological function, nerve conduction studies, and autonomic function were only prevented from worsening and failed to show an improvement (13). This result is in keeping with the lack of improvement in heart rate variability 43 months after simultaneous pancreas and kidney transplantation (SPK) (14). Neuropathy is, of course, extremely severe at this stage, as evidenced recently by the demonstration of severe intraepidermal nerve fiber depletion in pancreas transplant recipients, suggesting that long-term follow-up is necessary to assess posttransplant nerve fiber regeneration (15).

Our previous work suggested that

From the ¹Transplant Unit, Manchester Royal Infirmary, Manchester, U.K.; the ²Division of Cardiovascular Medicine, University of Manchester and Manchester Royal Infirmary, Manchester, U.K.; and the ³Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Australia.

Address correspondence and reprint requests to Dr. R.A. Malik, Division of Cardiovascular Medicine, University of Manchester, Manchester, M13 9NT, U.K. E-mail: rayaz.a.malik@man.ac.uk.

Received for publication 4 May 2007 and accepted in revised form 2 July 2007.

Published ahead of print at <http://care.diabetesjournals.org> on 10 July 2007. DOI: 10.2337/dc07-0870.

Abbreviations: CCM, corneal confocal microscopy; NBD, nerve branch density; NFD, nerve fiber density; NFL, nerve fiber length; NFT, nerve fiber tortuosity; SPK, simultaneous pancreas and kidney transplantation.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

small nerve fibers are the earliest to undergo damage and retain the ability to repair longer than large fibers (9,10). More recently, we have used corneal confocal microscopy (CCM) to demonstrate that corneal nerve fiber damage is directly related to the severity of somatic neuropathy (16,17) and to intraepidermal nerve fiber pathology (18). These data led us to propose that CCM, a noninvasive and reiterative test might be an ideal surrogate end point for evaluating therapeutic efficacy in clinical trials of human diabetic neuropathy (19). To test this hypothesis, we have assessed corneal sensitivity and used CCM to evaluate corneal nerve fiber morphology at baseline and 6 months after SPK in patients with type 1 diabetes.

RESEARCH DESIGN AND METHODS

Twenty type 1 diabetic patients undergoing SPK and 15 nondiabetic healthy control subjects were studied at baseline, and 15 diabetic patients underwent repeat assessment 6 months after SPK transplantation.

Corneal sensitivity

The noncontact corneal esthesiometer uses a puff of air expressed through a bore of 0.5-mm diameter with an electronic pressure sensor, which displays the force exerted in millibars. The stimulus jet is mounted on a slit lamp positioned 1 cm from the eye and aligned to the center of the cornea. Corneal sensitivity was assessed using our established methodology (20).

Confocal microscopy

Patients underwent examination with a Tomey Confoscan corneal confocal microscope model P4 (Tomey, Erlangen, Germany). One eye of each subject was selected at random and anesthetized with 1 drop of 0.4% benoxinate hydrochloride (oxybuprocaine hydrochloride; Minims). The objective lens of the confocal microscope was disinfected (isopropyl alcohol 70% vol/vol, swabs), and 1 drop of Viscotears liquid gel (carbomer 940; Ciba Vision Ophthalmics) was applied onto the tip of the lens and advanced forward until the gel touched the cornea, allowing optical but not physical contact between the objective lens and corneal epithelium. The entire cornea was scanned in ~2 min and en face two-dimensional images (lateral resolution ~1–2 μm and final image size of 768 \times 576 pixels) were acquired. Three to five high-quality images of Bowman's layer were examined as it contains

the main nerve plexus. The investigator who examined the cornea and who performed morphometric measurements of the images was masked with respect to the identity of the patient. The following parameters were quantified to define corneal nerve fiber damage and repair: 1) nerve fiber density (NFD), the total number of major nerves per square millimeter of corneal tissue; 2) nerve fiber length (NFL), the total length of all nerve fibers and branches per square millimeter of corneal tissue; 3) nerve branch density (NBD), the number of branches emanating from each nerve trunk per square millimeter of corneal tissue (16); and 4) nerve fiber tortuosity (NFT), a parameter mathematically derived from the images (17). Measures 1 and 3 were determined using morphometric software incorporated within the Tomey instrument, measure 2 was determined using third-party image analysis software (Scion Image for Windows; Scion, Frederick, MD), and measure 4 was calculated using a MATLAB function (MATLAB version 6.5; MathWorks, Natick, MA) that was created for this purpose (17). Corneal nerve morphology was quantified fully in each patient in ~30 min. To estimate the error in measuring NFD, NFL, and NBD, we acquired images and determined each of these parameters for 15 subjects on two occasions separated by at least 48 h. The coefficients of variation of these parameters were 12% for NFD, 9% for NFL, and 24% for NBD.

Statistics

SPSS 11.05.0 for Windows was used to compute the results. The data are expressed as means \pm SEM, and the analysis includes descriptive and frequency statistics. ANOVA with Scheffé post hoc tests was used to study differences between means. $P < 0.05$ was considered statistically significant.

RESULTS

Baseline evaluation

Twenty type 1 diabetic patients aged 41 \pm 1 years, with diabetes duration of 27 \pm 2 years and A1C 8.9 \pm 1.4% who were undergoing SPK, and 15 age-matched (46 \pm 3 years) healthy control subjects were examined. Corneal sensitivity was significantly reduced in diabetic patients (1.54 \pm 0.28 mbar) compared with control subjects (0.77 \pm 0.02 mbar, $P < 0.0001$). Corneal NFD (13.8 \pm 2.1 vs. 42 \pm 3.2, $P < 0.0001$), NBD (4.04 \pm 1.5




Figure 1—Image of corneal nerves in Bowman's layer, showing six nerve fibers with a typical beaded appearance, mild tortuosity, and adequate branching in a control subject (A) compared with marked loss of nerve fibers with one nerve in a patient undergoing pancreas transplantation (B) and improvement in NFD with increased numbers of nerves, 6 months after transplantation (C).

vs. 26.7 \pm 2.5, $P < 0.0001$), and NFL (2.23 \pm 0.2 vs. 9.69 \pm 0.7, $P < 0.0001$) were significantly reduced, and NFT (15.7 \pm 1.02 vs. 19.56 \pm 1.34, $P = 0.04$) was increased in diabetic patients (Fig. 1B) undergoing pancreas transplantation compared with control subjects (Fig. 1A and Table 1).

After transplantation

Fifteen patients underwent repeat assessment 6 months after SPK (Table 2). A1C fell significantly into the normal range (5.5 \pm 0.1 vs. 8.6 \pm 0.4%, $P = 0.007$), confirming successful pancreas transplantation. There was a trend for improvement in corneal sensitivity (1.23 \pm

Table 1—Corneal confocal nerve fiber morphology in control subjects and in type 1 diabetic patients undergoing SPK

	Age (years)	NCCA (mbar)	NFD (n/mm ²)	NBD (n/mm ²)	NFL (mm/mm ²)	NFT (Tc)
Control (n = 15)	46 ± 3	0.77 ± 0.02	42.04 ± 3.2	26.73 ± 2.5	9.69 ± 0.7	19.56 ± 1.34
Before SPK (n = 20)	41 ± 1.8	1.54 ± 0.28	13.88 ± 2.1	4.04 ± 1.5	2.23 ± 0.28	15.76 ± 1.02
P	NS	<0.0001	<0.0001	<0.0001	<0.0001	0.04

Data are means ± SEM with significant difference. NCCA, noncontact corneal esthesiometer; Tc, tortuosity coefficient.

0.39 vs. 1.54 ± 0.42, $P = 0.59$). Corneal NFD (18.04 ± 10.48 vs. 9.25 ± 1.87, $P = 0.001$) (Fig. 2A) and NFL (3.60 ± 0.33 vs. 1.84 ± 0.33, $P = 0.002$) (Fig. 2B) improved significantly. No change was observed for either NBD (1.38 ± 0.74 vs. 1.38 ± 1.00, $P = 1.0$) or NFT (15.58 ± 1.20 vs. 16.30 ± 1.19, $P = 0.67$).

CONCLUSIONS— The natural history of nerve damage in patients with type 1 diabetes is not entirely clear. Longitudinal data from the Rochester cohort support the contention that the duration and severity of exposure to hyperglycemia are related to the progression and hence severity of neuropathy rather than its onset (21). A recent study of patients with type 1 diabetes showed a high prevalence of diabetic neuropathy, which progressed in a significant proportion of patients and was related not only to glycemic control but also to conventional cardiovascular risk factors such as hypertension and lipids (22).

The replacement of functioning islet β -cells by pancreas transplantation has been considered to be the most logical treatment for patients with type 1 diabetes to normalize blood glucose and ameliorate long-term complications. Although pancreas transplantation takes ~5 years to prevent progression and 10 years to reverse the lesions of diabetic nephropathy (23), a recent study has demonstrated an improvement and/or stabilization of diabetic retinopathy after a median follow-up of only 17 months (24). For diabetic neuropathy, the largest and longest follow-up series to date has shown that pancreas transplantation improved both motor and sensory nerve conduction as

well as sudomotor function in the hand and foot within 1 year, which was maintained throughout follow-up for 10 years (13,25). However, autonomic function did not improve (13), as confirmed by another recent study (14). At this stage, most patients receiving transplantation will have severe nerve fiber damage as evidenced by marked depletion of intraepidermal nerve fibers (15). In addition, concomitant uremia, which often coexists in these patients, may contribute to nerve damage and limit repair, although renal transplantation in patients with diabetic nephropathy has not been shown to halt progression of neuropathy (26), and there was no difference in outcomes when pancreas alone transplantation was compared with combined pancreas and renal transplantation (13).

To determine a therapeutic response to an intervention in diabetic neuropathy, a standardized set of end points, which include clinical and neurophysiological evaluation combined with quantitative sensory and autonomic function testing, have been recommended (1,27,28). Pancreas transplantation has previously been shown to improve large nerve fiber conduction and, in particular, upper limb motor and sensory action potentials as well as sudomotor function within 1 year (13), but no impact on autonomic function was seen (13,14). Our studies show no relationship between quantitative sensory tests evaluating small fiber function and unmyelinated fiber pathology (9). Thus, although nerve conduction studies and quantitative sensory testing are useful and well-validated measures to help diagnose and assess progression of diabetic

neuropathy, their utility in evaluating a therapeutic response may be limited (29).

Although more detailed and reproducible measures that accurately quantify small fiber neuropathy such as skin or nerve biopsy will reduce this variability, they are invasive (9–12). Our recent work using the corneal confocal microscope suggests that this noninvasive and reiterative test might be an ideal surrogate end point for use in clinical trials of human diabetic neuropathy and therefore may be appropriate for assessing the benefits of pancreas transplantation (16–18). The cornea is richly innervated by the ophthalmic division of the trigeminal nerve via the anterior ciliary nerves. Corneal nerves have been studied by detailed light and electron microscopy (30,31) and immunohistology (32,33). CCM provides a novel approach to study corneal nerve morphology (34). The main advantage of this technique is that it enables a noninvasive in vivo evaluation of the human cornea at 700 \times magnification, with excellent resolution and contrast. Initial studies were descriptive, and hence their usefulness for the purposes of interpreting small fiber degeneration and regeneration is limited. However, a study from our group in control subjects refined and significantly improved the quantification of C and A- δ corneal nerve fibers (35), showing high concordance with previous histological studies (30).

One may argue that corneal innervation has little relevance to diabetic somatic neuropathy, characterized by a distal loss of nerve fibers innervating the lower limbs, and therefore has limited application as a measure of peripheral neu-

Table 2—A1C and corneal confocal nerve fiber morphology at baseline and 6 months after SPK in 15 patients with type 1 diabetes

	A1C	NCCA (mbar)	NFD (n/mm ²)	NBD (n/mm ²)	NFL (mm/mm ²)	NFT (Tc)
Before SPK	8.6 ± 0.4	1.54 ± 0.42	9.25 ± 1.87	1.38 ± 0.74	1.84 ± 0.33	16.30 ± 1.19
6 months after SPK	5.5 ± 0.1	1.23 ± 0.39	18.04 ± 1.48	1.38 ± 1.0	3.60 ± 0.33	15.58 ± 1.20
P	0.007	0.59	0.001	1.0	0.002	0.67

Data are means ± SEM with significant difference. NCCA, noncontact corneal esthesiometer; Tc, tortuosity coefficient.




Figure 2—NFD in diabetic patients before SPK at baseline compared with that of age-matched nondiabetic control subjects ($P < 0.0001$) with a significant improvement after SPK ($P = 0.001$) (A), and NFL in diabetic patients before SPK at baseline compared with that of age-matched nondiabetic control subjects ($P < 0.0001$) with a significant improvement after SPK ($P = 0.002$) (B). Results are means \pm SEM.

ropathy in diabetic patients. However, diabetic patients have a reduction in corneal sensitivity and a reduction in corneal nerve fiber bundles, which correlate with the Michigan Neuropathy Screening Instrument, a quantitative measure of somatic neuropathy (36). We have also

shown a progressive reduction in corneal sensitivity (20) and significant corneal nerve pathology, which relates to the severity of neuropathy assessed using neurophysiology, quantitative sensory testing (16,17), and, in particular, intraepidermal nerve fiber density (18).

In the present study, we have demonstrated a highly significant loss of corneal nerve fibers in type 1 diabetic patients undergoing pancreas transplantation, which confirms previous studies demonstrating severe neuropathy in patients undergoing pancreas transplantation (13–15). However, despite this considerable baseline damage, we have shown a significant improvement in corneal NFD and NFL within 6 months of transplantation, indicating an early repair process with the restoration of euglycemia. These findings are in contrast to previous studies in diabetic nephropathy (23), retinopathy (24), and particularly neuropathy (13,14,25), in which at best a prevention of progression in nerve damage was shown only after several years of euglycemia. However, these latter studies focused heavily on electrophysiology and quantitative sensory assessment, which predominantly measure large fiber function and, to a lesser extent, small fiber function. When small fiber function was assessed in the form of sudomotor function, it is of relevance that significant improvement was demonstrated within 1 year of SPK (13).

Our study using CCM focused on detailed pathology as opposed to function of the small fibers and demonstrated repair despite significant baseline damage. These observations support the view that in clinical intervention trials for diabetic neuropathy, perhaps the focus should be on assessment of small fiber damage and repair (1,8). Until recently, this could only be provided by costly, time-consuming, and, most importantly, invasive procedures such as nerve (10) and skin biopsy (11,12). We now show that CCM, a noninvasive and hence reiterative test, might be an ideal surrogate end point for assessing the benefits of pancreas transplantation and indeed for assessing therapeutic efficacy of other therapies in clinical trials of human diabetic neuropathy.

Acknowledgments—This work was supported by grant R01 NS46259-01 from the National Institutes of Health.

References

1. Boulton AJM, Malik RA, Arezzo JC, Soslenko JM: Diabetic somatic neuropathies. *Diabetes Care* 27:1458–86, 2004
2. Abbott CA, Vileikyte L, Williamson S, Carrington AL, Boulton AJM: Multicenter study of the incidence of and predictive risk factors for diabetic neuropathic foot ulceration. *Diabetes Care* 21:1071–1075,

- 1998
3. Dyck PJ, Thomas PK, Asbury AK, Winegrad AI, Porte D Jr: *Diabetic Neuropathy*. Philadelphia, Saunders, 1987
 4. Behse F, Buchthal F, Carlsen F: Nerve biopsy and conduction studies in diabetic neuropathy. *J Neurol Neurosurg Psychiatry* 40:1072–1082, 1977
 5. Dyck PJ, Karnes JL, Daube J, O'Brien PC, Service FJ: Clinical and neuropathological criteria for the diagnosis and staging of diabetic polyneuropathy. *Brain* 108:861–880, 1985
 6. Veves A, Malik RA, Lye RH, Masson EA, Sharma AK, Schady W, Boulton AJ: The relationship between sural nerve morphometric findings and measures of peripheral nerve function in mild diabetic neuropathy. *Diabet Med* 8:917–921, 1991
 7. Levy DM, Terenghi G, Gu XH, Abraham RR, Springall DR, Polak JM: Immunohistochemical measurements of nerves and neuropeptides in diabetic skin: relationship of tests of neurological function. *Diabetologia* 35:889–897, 1992
 8. Mojaddidi M, Quattrini C, Tavakoli M, Malik RA: Recent developments in the assessment of efficacy in clinical trials of diabetic neuropathy. *Curr Diab Rep* 5:417–422, 2005
 9. Malik RA, Veves A, Walker D, Siddique I, Lye RH, Schady W, Boulton AJ: Sural nerve fibre pathology in diabetic patients with mild neuropathy: relationship to pain, quantitative sensory testing and peripheral nerve electrophysiology. *Acta Neuropathol (Berl)* 101:367–374, 2001
 10. Malik RA, Tesfaye S, Newrick PG, Walker D, Rajbhandari SM, Siddique I, Sharma AK, Boulton AJ, King RH, Thomas PK, Ward JD: Sural nerve pathology in diabetic patients with minimal but progressive neuropathy. *Diabetologia* 48:578–585, 2005
 11. Singleton JR, Smith AG, Bromberg MB: Increased prevalence of impaired glucose tolerance in patients with painful sensory neuropathy. *Diabetes Care* 24:1448–1453, 2001
 12. Sumner C, Seth S, Griffin J, Cornblath D, Polydefkis M: The spectrum of neuropathy in diabetes and impaired glucose tolerance. *Neurology* 60:108–111, 2003
 13. Navarro X, Sutherland DE, Kennedy WR: Long-term effects of pancreatic transplantation on diabetic neuropathy. *Ann Neurol* 42:727–736, 1997
 14. Boucek P, Saudek F, Adamec M, Janousek L, Koznarova R, Havrdova T, Skibova J: Spectral analysis of heart rate variation following simultaneous pancreas and kidney transplantation. *Transplant Proc* 35:1494–1498, 2003
 15. Boucek P, Havrdova T, Voska L, Lodererova A, Saudek F, Lipar K, Janousek L, Adamec M, Sommer C: Severe depletion of intraepidermal nerve fibers in skin biopsies of pancreas transplant recipients. *Transplant Proc* 37:3574–3575, 2005
 16. Malik RA, Kallinikos P, Abbott CA, van Schie CH, Morgan P, Efron N, Boulton AJM: Corneal confocal microscopy: a non-invasive surrogate of nerve fibre damage and repair in diabetic patients. *Diabetologia* 46:683–688, 2003
 17. Kallinikos P, Berhanu M, O'Donnell C, Boulton AJM, Efron N, Malik RA: Corneal nerve tortuosity in diabetic patients with neuropathy. *Invest Ophthalmol Vis Sci* 45:418–422, 2004
 18. Quattrini C, Tavakoli M, Jeziorska M, Kallinikos P, Tesfaye S, Finnigan J, Marshall A, Boulton AJ, Efron N, Malik RA: Surrogate markers of small fiber damage in human diabetic neuropathy. *Diabetes* 56:2148–2154, 2007
 19. Hossain P, Sachdev A, Malik RA: Early detection of diabetic peripheral neuropathy using corneal confocal microscopy. *Lancet* 366:1340–1343, 2005
 20. Tavakoli M, Kallinikos PA, Efron N, Boulton AJ, Malik RA: Corneal sensitivity is reduced and relates to the severity of neuropathy in patients with diabetes. *Diabetes Care* 30:1895–1897, 2007
 21. Dyck PJ, Davies JL, Wilson DM, Service FJ, Melton LJ 3rd, O'Brien PC: Risk factors for severity of diabetic polyneuropathy: intensive longitudinal assessment of the Rochester Diabetic Neuropathy Study cohort. *Diabetes Care* 22:1479–1486, 1999
 22. Tesfaye S, Chaturvedi N, Eaton SE, Ward JD, Manes C, Ionescu-Tirgoviste C, Witte DR, Fuller JH, EURODIAB Prospective Complications Study Group: Vascular risk factors and diabetic neuropathy. *N Engl J Med* 352:341–350, 2005
 23. Fioretto P, Steffes MW, Sutherland DE, Goetz FC, Mauer M: Reversal of lesions of diabetic nephropathy after pancreas transplantation. *N Engl J Med* 339:69–75, 1998
 24. Giannarelli R, Coppelli A, Sartini M, Aragona M, Boggi U, Vistoli F, Rizzo G, Del Prato S, Mosca F, Marchetti P: Effects of pancreas-kidney transplantation on diabetic retinopathy. *Transpl Int* 18:619–622, 2005
 25. Kennedy WR, Navarro X, Goetz FC, Sutherland DE, Najarian JS: Effects of pancreatic transplantation on diabetic neuropathy. *N Engl J Med* 322:1031–1037, 1990
 26. Van der Vliet JA, Navarro X, Kennedy WR, Goetz FC, Barbosa JJ, Sutherland DE, Najarian JS: A long-term follow-up of polyneuropathy in diabetic kidney transplant recipients. *Diabetes* 37:1247–1252, 1988
 27. Report and recommendations of the San Antonio conference on diabetic neuropathy. *Diabetes* 37:1000–1004, 1988
 28. Proceedings of a consensus development conference on standardized measures in diabetic neuropathy. *Diabetes Care* 15:1080–1107, 1992
 29. Dyck PJ, Bushek W, Spring EM, Karnes JL, Litchy WJ, O'Brien PC, Service FJ: Vibratory and cooling detection thresholds compared with other tests in diagnosing and staging diabetic neuropathy. *Diabetes Care* 10:432–440, 1987
 30. Müller LJ, Pels L, Vrensen GFJM: Ultrastructural organisation of human corneal nerves. *Invest Ophthalmol Vis Sci* 37:476–488, 1996
 31. Müller LJ, Vrensen GFJM, Pels L, Nunes Cardozo B, Willekens B: Architecture of human corneal nerves. *Invest Ophthalmol Vis Sci* 38:985–994, 1997
 32. Tervo K, Tervo T, Eränkö L, Vannas A, Cuello AC, Eränkö O: Substance P-immunoreactive nerves in the human cornea and iris. *Invest Ophthalmol Vis Sci* 23:671–674, 1982
 33. Tovainen M, Tervo T, Partanen M, Vannas A, Hervonen A: Histochemical demonstration of adrenergic nerves in the stroma of human cornea. *Invest Ophthalmol Vis Sci* 28:398–400, 1987
 34. McGhee CNJ, Keller PR: In vivo confocal microscopy of living tissue: the cornea at cellular level. *Eyeneews* 5:14–20, 1985
 35. Oliveira-Soto L, Efron N: Morphology of corneal nerves using confocal microscopy. *Cornea* 20:374–384, 2001
 36. Rosenberg ME, Tervo TM, Immonen IJ, Muller LJ, Gronhagen-Riska C, Vesaluoma MH: Corneal structure and sensitivity in type 1 diabetes mellitus. *Invest Ophthalmol Vis Sci* 41:2915–2921, 2000