



Effect of Sitagliptin on Kidney Function and Respective Cardiovascular Outcomes in Type 2 Diabetes: Outcomes From TECOS

Diabetes Care 2016;39:2304–2310 | DOI: 10.2337/dc16-1415

Jan H. Cornel,¹ George L. Bakris,² Susanna R. Stevens,³ Michael Alvarsson,⁴ Willem A. Bax,¹ Lee-Ming Chuang,⁵ Samuel S. Engel,⁶ Renato D. Lopes,³ Darren K. McGuire,⁷ Axel Riefflin,⁸ Helena Wachslicht Rodbard,⁹ Isaac Sinay,¹⁰ Tsvetalina Tankova,¹¹ Julio Wainstein,¹² Eric D. Peterson,³ and Rury R. Holman,¹³ on behalf of the TECOS Study Group

OBJECTIVE

To evaluate chronic kidney disease (CKD) and cardiovascular outcomes in TECOS (Clinical trial reg. no. NCT00790205, clinicaltrials.gov) participants with type 2 diabetes and cardiovascular disease treated with sitagliptin, a dipeptidyl peptidase 4 inhibitor, according to baseline estimated glomerular filtration rate (eGFR).

RESEARCH DESIGN AND METHODS

We used data from 14,671 TECOS participants assigned in a double-blind design to receive sitagliptin or placebo added to existing therapy, while aiming for glycemic equipoise between groups. Cardiovascular and CKD outcomes were evaluated over a median period of 3 years, with participants categorized at baseline into eGFR stages 1, 2, 3a, and 3b (≥ 90 , 60–89, 45–59, or 30–44 mL/min/1.73 m², respectively).

RESULTS

Participants with eGFR stage 3b were older, were more often female, and had a longer duration of diabetes. Four-point major adverse cardiovascular event rates increased with lower baseline eGFR (3.52, 3.55, 5.74, and 7.34 events/100 patient-years for stages 1–3b, respectively). Corresponding adjusted hazard ratios for stages 2, 3a, and 3b versus stage 1 were 0.93 (95% CI 0.82–1.06), 1.28 (1.10–1.49), and 1.39 (1.13–1.72), respectively. Sitagliptin therapy was not associated with cardiovascular outcomes for any eGFR stage (interaction *P* values were all >0.44). Kidney function declined at the same rate in both treatment groups, with a marginally lower but constant eGFR difference (-1.3 mL/min/1.73 m²) in those participants who were assigned to sitagliptin. Treatment differences in these eGFR values remained after adjustment for region, baseline eGFR, baseline HbA_{1c}, time of assessment, and within-study HbA_{1c} levels.

CONCLUSIONS

Impaired kidney function is associated with worse cardiovascular outcomes. Sitagliptin has no clinically significant impact on cardiovascular or CKD outcomes, irrespective of baseline eGFR.

¹Noordwest Ziekenhuisgroep, Alkmaar, the Netherlands

²University of Chicago Medicine, Chicago, IL

³Duke Clinical Research Institute, Duke University School of Medicine, Durham, NC

⁴Department of Endocrinology, Metabolism and Diabetes, Karolinska University Hospital, Solna, Stockholm, Sweden

⁵Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan

⁶Merck and Co., Inc., Kenilworth, NJ

⁷Department of Medicine, University of Texas Southwestern Medical Center, Dallas, TX

⁸GMP, Husby, Germany

⁹Endocrine and Metabolic Consultants, Rockville, MD

¹⁰Unit of Diabetes, Instituto Cardiovascular de Buenos Aires, Buenos Aires, Argentina

¹¹Clinical Center of Endocrinology, Medical University, Sofia, Bulgaria

¹²E. Wolfson Medical Center, Holon, Israel

¹³Diabetes Trials Unit, Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Oxford, U.K.

Corresponding author: Rury R. Holman, rury.holman@dtu.ox.ac.uk.

Received 1 July 2016 and accepted 16 September 2016.

Clinical trial reg. no. NCT00790205, clinicaltrials.gov.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc16-1415/-/DC1>.

This article is featured in a podcast available at <http://www.diabetesjournals.org/content/diabetes-core-update-podcasts>.

© 2016 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

Patients with type 2 diabetes mellitus are at high risk for macrovascular and microvascular complications (1), with type 2 diabetes being a key risk factor for the development of chronic kidney disease (CKD). CKD further increases the risk of adverse cardiovascular (CV) outcomes, especially in patients with known CV disease (2–5), and both microalbuminuria and macroalbuminuria are associated independently with an increased risk of CV events (6). Accordingly, the potential impact of type 2 diabetes therapies on CV and CKD outcomes is a major consideration in the long-term management strategy of the disease. Intensified glucose control and multiple CV risk factor therapies can reduce CV risk in general, and diabetic nephropathy in particular (7,8), but there is a paucity of data on the effectiveness of specific type 2 diabetes treatment regimens with regard to these two outcomes.

The Trial Evaluating Cardiovascular Outcomes With Sitagliptin (TECOS) (9) showed that adding sitagliptin, compared with placebo, to usual care in patients with type 2 diabetes and established CV disease did not have an impact on the risk of major CV outcomes, hospitalization for heart failure, or adverse events in general. The aim of the present post hoc analysis was to evaluate CV and CKD outcomes in TECOS participants with type 2 diabetes and CV disease when treated with sitagliptin, a dipeptidyl peptidase 4 inhibitor (DPP-4i), according to their baseline estimated glomerular filtration rate (eGFR) stage.

RESEARCH DESIGN AND METHODS

The rationale and design of TECOS (10), as well as its primary outcomes and safety measures (9), have been reported. Briefly, 14,735 participants from 38 countries were enrolled in the study between December 2008 and July 2012. Eligible participants were ≥ 50 years old with type 2 diabetes, established atherosclerotic CV disease, and HbA_{1c} values in the range 6.5–8.0% (48–64 mmol/mol) and were receiving stable-dose monotherapy or dual-combination therapy with metformin, pioglitazone, or sulfonylurea, or insulin with or without metformin. Patients with eGFR values < 30 mL/min/1.73 m² were excluded from the study.

Participants were randomized in a double-blind manner to receive sitagliptin

100 mg/day or placebo, with a lower dose of 50 mg/day for those with eGFR values 30–50 mL/min/1.73 m². During the study, sitagliptin doses were adjusted, based on at least annual eGFR values, to 50 mg/day if eGFR values were 30–50 mL/min/1.73 m² and to 25 mg/day if eGFR values were < 30 mL/min/1.73 m². If a sustained eGFR recovery occurred, sitagliptin doses could also be up-titrated.

Treatment for type 2 diabetes and its comorbidities was provided by usual care providers, based on local guidelines. Any other glucose-lowering agent could be added, except for a glucagon-like peptide 1 receptor agonist or an open-label DPP-4i, with rosiglitazone use discouraged.

The study was managed and all data were adjudicated and analyzed by academic partners (Duke Clinical Research Institute and the University of Oxford Diabetes Trials Unit). The database was held at and independently verified by the Duke Clinical Research Institute.

Ascertainment of CV Outcomes

An independent clinical events committee, blinded to treatment allocation, adjudicated all events of death, myocardial infarction (MI), stroke, hospitalization for unstable angina, and hospitalization for heart failure (10). The clinical events committee, which was independent of both the sponsor and the TECOS Executive Committee, remained blinded to study treatment assignment. The primary CV composite outcome was a 4-point major adverse CV event (MACE), defined as time to CV death, nonfatal MI, nonfatal stroke, or hospitalization for unstable angina.

Ascertainment of Kidney Function

Kidney function during the trial was assessed by annual usual care measurements of eGFR, calculated using the Modification of Diet in Renal Disease Study Equation (11). For a subset of participants, usual care urinary albumin-to-creatinine ratio (UACR) measurements were also available.

Statistical Analysis

Participants were categorized at baseline into eGFR stages 1, 2, 3a, and 3b (≥ 90 , 60–89, 45–59, and 30–44 mL/min/1.73 m², respectively) (12) and into three UACR groups according to their baseline values (normoalbuminuria < 30 mg/g, microalbuminuria 30–300 mg/g, and

macroalbuminuria > 300 mg/g). Baseline characteristics for the intention-to-treat (ITT) population were summarized as mean (± 1 SD) or median (25th, 75th percentile) for quantitative data and as percentages for categorical data.

Separate Kaplan-Meier plots for the primary 4-point MACE outcome were created for each eGFR stage, split by assigned study treatment or by HbA_{1c} level above or below the median. Possible associations between CV outcomes and the CKD stage or the UACR category were evaluated using Cox proportional hazards regression models, with region included as a stratification variable and adjustment covariates taken from models developed previously for the Nateglinide and Valsartan in Impaired Glucose Tolerance Outcomes Research (NAVIGATOR) trial (13–15). Less than 4% of the adjustment variables had missing values apart from LDL cholesterol (24%), hemoglobin (34%), and UACR (65%). For modeling purposes, missing baseline data were imputed using SAS PROC MI (SAS Institute, Cary, NC). CV outcome rates are presented as the total number of events and as events/100 patient-years of follow-up. Adjusted hazard ratios and 95% CIs are presented for each of eGFR stages 2–3b, with stage 1 as the reference group. Models were repeated with the addition of the treatment-by-eGFR interaction.

Kidney-related outcomes included changes from baseline in eGFR and UACR, as specified in the TECOS protocol and statistical analysis plan (9). Repeated-measures ANOVA was used to test for between-treatment group eGFR and UACR differences over 4 years, with the overall difference summarized as the least squares mean difference and 95% CI. Overall least squares mean differences are presented for all patients and separately for each eGFR stage, with *P* values for the treatment group-by-eGFR stage interaction. Models were adjusted for region, baseline eGFR or UACR, and for study visit. The model for the association between treatment and eGFR changes over 4 years was also performed with adjustment for baseline HbA_{1c} level and change in HbA_{1c} level from baseline to each visit.

Data were analyzed with SAS version 9.4. *P* values < 0.05 were considered to be statistically significant, with no adjustments made for multiple testing.

0.93 (95% CI 0.82–1.06), 1.28 (1.10–1.49), and 1.39 (1.13–1.72), respectively. Rates increased similarly for the secondary 3-point MACE outcome (CV death, non-fatal MI, or nonfatal stroke) and all other secondary outcomes, except for hospitalization for unstable angina. There were no significant interactions (all $P > 0.44$) between continuous eGFR measurements and randomized treatment allocation (Supplementary Table 2).

Sitagliptin treatment did not have an impact on the primary 4-point MACE outcome irrespective of baseline eGFR stage, as shown by Kaplan-Meier curves (Supplementary Fig. 1). Lower event rates were seen in those participants who had within-study HbA_{1c} values below the median, compared with those with HbA_{1c} values greater than the median (Supplementary Fig. 2).

Table 2 shows the CV outcomes by baseline UACR category. CV outcomes worsened with increasing albuminuria, except for 4-point MACE, MI, stroke, and hospitalization for unstable angina. The modeled impact of continuous baseline eGFR and UACR values on the 4-point MACE primary outcome are shown in Supplementary Fig. 3. Rates increase substantially with eGFR values <60 mL/min/1.73 m² and with UACR values >30 mg/g.

Kidney Outcomes

The mean eGFR reduction over 4 years from baseline was greater in the sitagliptin group than in the placebo group (-4.0 ± 18.4 vs. -2.8 ± 18.3 mL/min/1.73 m²). The mean eGFR value was marginally lower in the sitagliptin group at the first postrandomization visit and remained

consistently lower thereafter (Fig. 1A), with an overall estimated least squares mean difference of -1.34 mL/min/1.73 m² (95% CI -1.76 to -0.91 , $P < 0.001$) (Table 3). The 4-year between-treatment group differences were similar for each eGFR stage, with no significant interactions of treatment effect by eGFR stage (Table 3 and Supplementary Fig. 4). The slight eGFR difference between treatment groups remained after adjusting for time from randomization when eGFR was measured, baseline eGFR, baseline HbA_{1c} level, change in HbA_{1c} level over time, and region (Supplementary Table 3), with an estimated overall mean difference of -1.43 mL/min/1.73 m² (95% CI -1.88 to -0.98 , $P < 0.0001$).

In the subset of participants with UACR data, the median value was marginally and consistently lower in the sitagliptin group compared with the placebo group (Fig. 1B), with an estimated overall mean difference of -0.18 mg/g (95% CI -0.35 to -0.02 , $P = 0.031$) (Table 3). The 4-year UACR between-treatment group differences were similar for each eGFR stage, with no significant interactions of treatment effect by eGFR stage (Table 3).

CONCLUSIONS

TECOS was a global clinical trial demonstrating that the addition of sitagliptin to usual care in patients with type 2 diabetes and established CV disease did not affect rates of major atherosclerotic CV events in a setting of glycemic equipoise. This study shows that, although CV events are more frequent in patients with lower levels of kidney

function, there is no interaction with the addition of sitagliptin. Kidney function declined at the same rate in both the sitagliptin and placebo groups, but with a slightly lower and constant eGFR difference in those assigned to receive sitagliptin.

CV Outcomes

The primary 4-point MACE outcome rate was progressively higher in participants with lower eGFR levels, as was the 3-point MACE outcome rate in those with an increased UACR. These data confirm earlier observational studies in which both eGFR and albuminuria have been shown to be independently associated with increased mortality and morbidity (3). Interestingly, the rise in the rate of the primary 4-point MACE outcome starts with a UACR as low as 30 mg/g, emphasizing albuminuria as a strong predictor of risk, as shown previously by Matsushita et al. (16). As would be expected, TECOS participants with reduced baseline kidney function were less likely to be taking metformin and were more often receiving insulin therapy, but with no difference in the rates of sulfonylurea use. Despite the varying glucose-lowering strategies, sitagliptin therapy had no effect on the primary 4-point MACE outcome at any eGFR or UACR level and thus appears to be safe with respect to CV outcomes in patients with decreased kidney function.

CKD Outcomes

Although the mean eGFR during the trial was marginally lower in the sitagliptin group compared with the placebo group, even when adjusted for glycemic

Table 2—Association of CV end points with baseline UACR categories

End point	Total number of events (events/100 patient-years)			Adjusted hazard ratio (95% CI) vs. normoalbuminuria (UACR <30 mg/g)		
	Normoalbuminuria UACR <30 mg/g	Microalbuminuria UACR 30–300 mg/g	Macroalbuminuria UACR >300 mg/g	UACR 30–300 mg/g	UACR >300 mg/g	P value
CV death, MI, stroke, or hospitalization for UA	381 (3.54)	165 (5.03)	46 (7.13)	1.19 (0.99–1.43)	1.33 (0.96–1.83)	0.0797
CV death, MI, or stroke	331 (3.05)	155 (4.71)	46 (7.13)	1.28 (1.05–1.56)	1.52 (1.10–2.11)	0.0066
CV death	119 (1.03)	79 (2.26)	24 (3.41)	1.86 (1.39–2.49)	2.27 (1.43–3.60)	<0.0001
Hospitalization for UA	65 (0.58)	12 (0.35)	0 (0)	0.56 (0.30–1.06)		0.2018
MI	174 (1.58)	63 (1.88)	22 (3.36)	1.04 (0.77–1.40)	1.52 (0.95–2.42)	0.2172
Stroke	79 (0.71)	35 (1.03)	12 (1.78)	1.16 (0.77–1.75)	1.75 (0.92–3.32)	0.2179
All-cause death	203 (1.76)	105 (3)	34 (4.83)	1.45 (1.14–1.84)	1.82 (1.25–2.66)	0.0006
Hospitalization for heart failure	94 (0.84)	53 (1.57)	20 (3.07)	1.63 (1.15–2.29)	2.78 (1.68–4.59)	<0.0001

UA, unstable angina.




Figure 1—A: eGFR over 4 years ($N = 13,604$). B: UACR over 4 years ($N = 3,832$). Data are plotted at each visit as the mean (± 1 SD) for eGFR and the median (25th, 75th percentile) for UACR among patients with the measurement at the visit. Patients without baseline and at least one postbaseline measure are not shown at any visit.

control, the eGFR decline was the same. Mean UACR values were also marginally lower with sitagliptin than with placebo in the 26% of TECOS participants who had these data available. It is uncertain whether these small offsets in eGFR and UACR would have any long-term clinical implications. Similar observations of a decrease in UACR have been made in post hoc pooled analyses of phase 3 DPP-4i studies using linagliptin (17). Animal studies (18) in streptozotocin-induced diabetic rats found that DPP-4i treatment improved albuminuria, with

similar improvements in creatinine clearance. Although microalbuminuria rates can be reduced by improving glucose control (19), the minimal effects of sitagliptin on eGFR and UACR appear to be unrelated to its glucose-lowering effects since they are not explained by baseline HbA_{1c} levels or HbA_{1c} level changes during the trial. Although the small eGFR offset occurs early and is stable over time for all GFR categories, there is no evidence of progression, and the offset appears to be similar for other DPP-4i agents as well (20).

TECOS did not show a clinically relevant improvement of kidney outcomes in patients treated with sitagliptin. Microvascular complication rates are related to HbA_{1c} levels (21), and can be reduced with improved glycemic control (22,23), but in TECOS there was only a small difference in HbA_{1c} levels between treatment groups because the study aimed to achieve glycemic equipoise in order to minimize possible glycemic confounding effects on the outcomes of interest. The decline of kidney function in individuals with diabetes is often, but not always, preceded by glomerular hyperfiltration, possibly as early as the stage of impaired fasting glucose (24). Glomerular hyperfiltration is associated with high glucose levels and changes in tubuloglomerular feedback related to alterations in vasoactive mediators, such as nitric oxide and cyclooxygenase-2-derived prostanoids, resulting in glomerular hypertension. These changes contribute to the inflammatory nature of diabetes, which affects the vasculature and is directly associated with the genesis of microalbuminuria. Ultimately, in a subgroup of people, there is a decline in kidney function, with some individuals progressing to end-stage renal disease (25,26). Early studies (26,27) suggested that microalbuminuria was a predictor of faster declines in kidney function, but over the past decade the data clearly indicate it is a marker of increased CV disease risk in various pathophysiologic conditions, such as diabetes.

Strengths and Limitations

The strength of the current study was the large number of patients studied in a double-blind prospective manner. Limitations include the fact that follow-up may be relatively short for evaluating the risk of the development of diabetic nephropathy, especially in view of the biphasic change in GFR with initial hyperfiltration followed by a decrease in GFR. Furthermore, no differences were taken into account for eGFR stage 1 and CKD stage 1, in which microalbuminuria must be present and which may be a somewhat different class of patients.

Conclusion

Reduced eGFR and increased UACR were associated with a significantly increased risk of CV events, but we observed no clinically significant effect of sitagliptin treatment on CV outcomes or CKD

Table 3—Estimated mean 4-year eGFR and UACR between-treatment group differences (sitagliptin minus placebo), overall and by baseline eGFR stages

	Baseline value	Mean between-group treatment difference (95% CI) [†]	P value [‡]
eGFR (N = 13,604), mL/min/1.73 m²			
Overall	75.1 ± 21.0	−1.34 (−1.76 to −0.91)	<0.0001
Stage 1 (eGFR ≥90 mL/min/1.73 m ²)	104 ± 14	−0.22 (−1.19 to 0.75)	Interaction P = 0.14
Stage 2 (eGFR 60–89 mL/min/1.73 m ²)	73 ± 9	−1.42 (−2.05 to −0.79)	
Stage 3a (eGFR 45–59 mL/min/1.73 m ²)	53 ± 4	−1.33 (−2.45 to −0.21)	
Stage 3b (eGFR 30–44 mL/min/1.73 m ²)	39 ± 4	−2.25 (−4.27 to −0.23)	
UACR (N = 3,832), mg/g			
Overall	11.1 (3.9, 35.0)	−0.18 (−0.35 to −0.02)	0.031
Stage 1 (eGFR ≥90 mL/min/1.73 m ²)	11.0 (4.7, 30.2)	−0.18 (−0.53 to 0.16)	Interaction P = 0.68
Stage 2 (eGFR 60–89 mL/min/1.73 m ²)	9.7 (3.5, 29.2)	−0.20 (−0.42 to 0.02)	
Stage 3a (eGFR 45–59 mL/min/1.73 m ²)	14.3 (4.1, 55.4)	−0.30 (−0.70 to 0.09)	
Stage 3b (eGFR 30–44 mL/min/1.73 m ²)	27.7 (9.7, 126.6)	0.23 (−0.54 to 1.00)	

eGFR data are presented as the mean ± 1 SD, and UACR data are presented as median (25th, 75th percentiles). [†]The estimated mean difference and P value are derived from repeated measures over the 4-year time frame. Estimated mean differences for UACR are modeled and presented with a Box-Cox transformation. Models include region, baseline eGFR stage, time of measure, treatment, and the eGFR stage-by-treatment interaction. The UACR analysis also adjusts for continuous baseline UACR.

progression in patients with different CKD categories at baseline.

Acknowledgments. The authors thank the investigators, staff, and participants in the TECOS trial without whose efforts and collaboration this work would not have been possible. The authors also thank the following academic partners and contract research organizations for their assistance: Parexel International, Jubilant Clinsys, Clinogent, Canadian VIGOUR Centre, Green Lane Coordinating Centre, and South Australian Health and Medical Research Institute.

Funding. Funded by Merck Sharp & Dohme, a subsidiary of Merck and Co., Inc.

Duality of Interest. J.H.C. has received personal fees from Merck and Co., Inc., Eli Lilly, and AstraZeneca. G.L.B. reports serving as consultant for Merck and Co., Inc., Bayer, Boehringer Ingelheim, AbbVie, Janssen, NxStage, AstraZeneca, and Sanofi. S.S.E. is an employee of Merck Sharp & Dohme, a subsidiary of Merck & Co., Inc., the manufacturer of sitagliptin. R.D.L. reports receiving research grants from Bristol-Myers Squibb and GlaxoSmithKline and serving as a consultant for Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Merck and Co., Inc., Pfizer, and Portola. D.K.M. has received personal fees from Boehringer Ingelheim, Janssen Research and Development LLC, Sanofi Group, Merck Sharp & Dohme, Daiichi Sankyo, Inc., Lilly USA, Novo Nordisk, GlaxoSmithKline, Takeda Pharmaceuticals North America, AstraZeneca, Lexicon, Regeneron, University of Oxford, Duke Clinical Research Institute, Partners Healthcare, and the Cleveland Clinic Foundation. H.W.R. conducts clinical trials, lectures, and serves on advisory boards for AstraZeneca, Boehringer Ingelheim, Janssen, Lilly, Merck and Co., Inc., Novo Nordisk, Sanofi, and Regeneron. I.S. has received grants, personal fees, and support from Merck and Co., Inc., AstraZeneca, Janssen, Boehringer Ingelheim, Novartis, Novo Nordisk,

GlaxoSmithKline, and Servier. T.T. has received lecture fees and serves on advisory boards for Merck and Co., Inc., Boehringer Ingelheim, Novo Nordisk, Sanofi, Eli Lilly, Novartis, Servier, and AstraZeneca. J.W. reports serving as a consultant for AstraZeneca, Novo Nordisk, Eli Lilly, Novartis, Boehringer Ingelheim, and Sanofi. E.D.P. has received grants and personal fees from Janssen, grants from Eli Lilly, and personal fees from AstraZeneca, Bayer, and Sanofi. R.R.H. has received grants and personal fees from Merck and Co., Inc.; grants from Bayer and AstraZeneca; personal fees from Amgen, Bayer, Intarcia, Novartis, and Novo Nordisk; and other support from GlaxoSmithKline, Janssen, and Takeda. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. J.H.C. and W.A.B. drafted and edited the manuscript. G.L.B., M.A., L.-M.C., R.D.L., A.R., H.W.R., I.S., T.T., and J.W. edited the manuscript. S.R.S. performed the statistical analyses and edited the manuscript. S.S.E., D.K.M., and E.D.P. edited the manuscript and contributed to the study design and the data analysis and interpretation. R.R.H. drafted and edited the manuscript and contributed to the study design and data analysis and interpretation. R.R.H. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Parts of this study were presented in abstract form at the 75th Scientific Sessions of the American Diabetes Association, Boston, MA, 5–9 June 2015; the 51st EASD Annual Meeting, Stockholm, Sweden, 14–18 September 2015; and the Scientific Programme of the World Diabetes Congress 2015, Vancouver, British Columbia, Canada, 30 November–4 December 2015.

References

1. Stamler J, Vaccaro O, Neaton JD, Wentworth D. Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the Multiple

Risk Factor Intervention Trial. *Diabetes Care* 1993; 16:434–444

2. Anavekar NS, McMurray JJ, Velazquez EJ, et al. Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. *N Engl J Med* 2004;351:1285–1295

3. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004;351:1296–1305

4. Tonelli M, Muntner P, Lloyd A, et al.; Alberta Kidney Disease Network. Risk of coronary events in people with chronic kidney disease compared with those with diabetes: a population-level cohort study. *Lancet* 2012;380:807–814

5. Fox CS, Matsushita K, Woodward M, et al.; Chronic Kidney Disease Prognosis Consortium. Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without diabetes: a meta-analysis. *Lancet* 2012;380:1662–1673

6. Mahmoodi BK, Matsushita K, Woodward M, et al.; Chronic Kidney Disease Prognosis Consortium. Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without hypertension: a meta-analysis. *Lancet* 2012;380:1649–1661

7. UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet* 1998;352:854–865

8. Gaede P, Lund-Andersen H, Parving H-H, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med* 2008;358:580–591

9. Green JB, Bethel MA, Armstrong PW, et al.; TECOS Study Group. Effect of sitagliptin on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2015;373:232–242

10. Green JB, Bethel MA, Paul SK, et al. Rationale, design, and organization of a randomized, controlled Trial Evaluating Cardiovascular Outcomes with Sitagliptin (TECOS) in patients with type 2 diabetes and established cardiovascular disease. *Am Heart J* 2013;166:983–989.e7

11. Rossing P, Rossing K, Gaede P, Pedersen O, Parving HH. Monitoring kidney function in type 2 diabetic patients with incipient and overt diabetic nephropathy. *Diabetes Care* 2006;29:1024–1030
12. Levey AS, Tangri N, Stevens LA. Classification of chronic kidney disease: a step forward. *Ann Intern Med* 2011;154:65–67
13. Preiss D, Giles TD, Thomas LE, et al. Predictors of stroke in patients with impaired glucose tolerance: results from the Nateglinide and Valsartan in Impaired Glucose Tolerance Outcomes Research trial. *Stroke* 2013;44:2590–2593
14. Preiss D, Thomas LE, Sun JL, et al. Predictors of cardiovascular events in a contemporary population with impaired glucose tolerance: an observational analysis of the Nateglinide and Valsartan in Impaired Glucose Tolerance Outcomes Research (NAVIGATOR) trial. *BMJ Open* 2012;2:e001925
15. Wong YW, Thomas L, Sun JL, et al. Predictors of incident heart failure hospitalizations among patients with impaired glucose tolerance: insight from the Nateglinide And Valsartan in Impaired Glucose Tolerance Outcomes Research study. *Circ Heart Fail* 2013;6:203–210
16. Matsushita K, Coresh J, Sang Y, et al.; CKD Prognosis Consortium. Estimated glomerular filtration rate and albuminuria for prediction of cardiovascular outcomes: a collaborative meta-analysis of individual participant data. *Lancet Diabetes Endocrinol* 2015;3:514–525
17. Groop PH, Cooper ME, Perkovic V, Emser A, Woerle HJ, von Eynatten M. Linagliptin lowers albuminuria on top of recommended standard treatment in patients with type 2 diabetes and renal dysfunction. *Diabetes Care* 2013;36:3460–3468
18. Liu WJ, Xie SH, Liu YN, et al. Dipeptidyl peptidase IV inhibitor attenuates kidney injury in streptozotocin-induced diabetic rats. *J Pharmacol Exp Ther* 2012;340:248–255
19. Cull CA, Manley S, Frighi V, Holman RR, Turner RC. UK Prospective Diabetes Study (UKPDS). X. Urinary albumin excretion over 3 years in diet-treated type 2, (non-insulin-dependent) diabetic patients, and association with hypertension, hyperglycaemia and hypertriglyceridaemia. *Diabetologia* 1993;36:1021–1029
20. Cooper ME, Perkovic V, McGill JB, et al. Kidney disease end points in a pooled analysis of individual patient-level data from a large clinical trials program of the dipeptidyl peptidase 4 inhibitor linagliptin in type 2 diabetes. *Am J Kidney Dis* 2015;66:441–449
21. Stratton IM, Adler AI, Neil HA, et al. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ* 2000;321:405–412
22. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–853
23. Holman RR, Paul SK, Bethel MA, Matthews DR, Neil HA. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008;359:1577–1589
24. Melsom T, Mathisen UD, Ingebreetsen OC, et al. Impaired fasting glucose is associated with renal hyperfiltration in the general population. *Diabetes Care* 2011;34:1546–1551
25. Vora JP, Dolben J, Dean JD, et al. Renal hemodynamics in newly presenting non-insulin dependent diabetes mellitus. *Kidney Int* 1992;41:829–835
26. Bakris GL, Molitch M. Microalbuminuria as a risk predictor in diabetes: the continuing saga. *Diabetes Care* 2014;37:867–875
27. Lambers Heerspink HJ, Gansevoort RT. Albuminuria is an appropriate therapeutic target in patients with CKD: the pro view. *Clin J Am Soc Nephrol* 2015;10:1079–1088