

Diagnostic Accuracy of Point-of-Care Testing for Diabetic Ketoacidosis at Emergency-Department Triage

β -hydroxybutyrate versus the urine dipstick

SANJAY ARORA, MD¹
SEAN O. HENDERSON, MD¹

THEODORE LONG, MD²
MICHAEL MENCHINE, MD, MPH¹

RESEARCH DESIGN AND METHODS

OBJECTIVE—In the emergency department, hyperglycemic patients are screened for diabetic ketoacidosis (DKA) via a urine dipstick. In this prospective study, we compared the test characteristics of point-of-care β -hydroxybutyrate (β -OHB) analysis with the urine dipstick.

RESEARCH DESIGN AND METHODS—Emergency-department patients with blood glucose ≥ 250 mg/dL had urine dipstick, chemistry panel, venous blood gas, and capillary β -OHB measurements. DKA was diagnosed according to American Diabetes Association criteria.

RESULTS—Of 516 hyperglycemic subjects, 54 had DKA. The urine dipstick had a sensitivity of 98.1% (95% CI 90.1–100), a specificity of 35.1% (30.7–39.6), a positive predictive value of 15% (11.5–19.2), and a negative predictive value of 99.4% (96.6–100) for DKA. Using the manufacturer-suggested cutoff of >1.5 mmol/L, β -OHB had a sensitivity of 98.1% (90.1–100), a specificity of 78.6% (74.5–82.2), a positive predictive value of 34.9% (27.3–43), and a negative predictive value of 99.7% (98.5–100) for DKA.

CONCLUSIONS—Point-of-care β -OHB and the urine dipstick are equally sensitive for detecting DKA (98.1%). However, β -OHB is more specific (78.6 vs. 35.1%), offering the potential to significantly reduce unnecessary DKA work-ups among hyperglycemic patients in the emergency department.

Rapid and accurate identification of patients with diabetic ketoacidosis (DKA) is critical but is complicated by the fact that DKA only affects a small percentage of the total number of patients with hyperglycemia (1,2). Current clinical practice in the emergency department is to screen hyperglycemic patients with a urine dipstick for ketones and conduct a laboratory evaluation for DKA on patients whose results screen positive (3). Although the urine dipstick is easily accessible, inexpensive, rapid, and has excellent sensitivity for DKA, its poor specificity (estimated at $<50\%$) results

in a large number of false-positive tests and unnecessary work-ups (4). This has led the American Diabetes Association (ADA) to discourage using the urine dipstick and encourage using serum ketones for DKA screening (5). The development of point-of-care β -hydroxybutyrate (β -OHB) devices has made following these ADA recommendations feasible in the emergency department (6). The goal of this prospective study was to compare the test characteristics of a point-of-care β -OHB meter with the urine dipstick for screening for DKA at emergency-department triage.

Setting

This prospective, observational study was conducted at a large, urban emergency department. Institutional review board approval was obtained, and study participants provided written informed consent. The cohort studied represents a convenience sample of emergency-department patients with triage capillary blood glucose ≥ 250 mg/dL enrolled on 320 individual days over a 2-year period based on research-assistant availability. Subjects were excluded if they were 1) critically ill, 2) in police custody, 3) suffering from acute psychosis, or 4) unable to give informed consent.

Study procedures and data collection

Included patients had a urine dipstick (Siemens Multistix 10SG) for ketones (recorded as positive or negative), a serum chemistry panel, a venous blood gas to determine serum pH, and a point-of-care capillary β -OHB measurement (Precision Xtra meter; Medisense/Abbott Laboratories). DKA was defined according to the following ADA criteria: serum glucose ≥ 250 mg/dL; anion gap >10 mmol/L; carbon dioxide ≤ 18 mmol/L; and pH ≤ 7.30 (7). Subjects' race, ethnicity, age, sex, previous history of diabetes, and insulin use were recorded.

Sample size considerations

On the basis of previous literature, we assumed that the true sensitivity of β -OHB for detecting DKA was between 98 and 100% and determined that a sample including 54 cases of DKA would confirm this point estimate with a confidence limit no lower than 90% (8–12). The study was stopped after the 54th DKA patient was enrolled.

Statistical analysis

Data were analyzed using Stata version 10.0 (StataCorp, College Station, TX). Diagnostic test characteristics with 95%

From the ¹Department of Clinical Emergency Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California; and the ²Keck School of Medicine, University of Southern California, Los Angeles, California.

Corresponding author: Sanjay Arora, sanjay.arora@usc.edu.

Received 4 October 2010 and accepted 7 January 2011.

DOI: 10.2337/dc10-1844

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc10-1844/-/DC1>.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

CI) were calculated. According to the manufacturer's guidelines, point-of-care β -OHB ≤ 1.5 mmol/L was considered negative and >1.5 mmol/L was considered positive. The difference in specificity for β -OHB and urine dipstick ketones was assessed using the McNemar test with a Yates correction (13). We explored the optimal capillary β -OHB cutoff by creating a receiver-operator curve and analyzing diagnostic test characteristics at various cutoffs.

RESULTS

Participants

A total of 859 patients were screened, 616 provided informed consent, and 516 (83.7%) had all data elements necessary for data analysis (excluded patients: 49 with no urine dipstick, 30 with no serum pH via venous blood gas, and 21 with no serum chemistry panel). In our final sample, 54 of 516 subjects (10.5%) met the ADA criteria for DKA. See Table 1 for sample characteristics.

Diagnostic accuracy

At the manufacturer-suggested cutoff of >1.5 mmol/L, the sensitivity of capillary β -OHB and the urine dipstick for DKA were identical at 98.1% (95% CI 90.1–100). However, the specificity of β -OHB was significantly higher than that of the urine dipstick (78.6 [74.5–82.2] vs. 35.1% [30.7–39.6]; $P < 0.01$). Positive and negative predictive values of capillary β -OHB were 34.9 (27.3–43) and 99.7% (98.5–100), respectively. Positive and negative predictive values of the urine dipstick were 15 (11.5–19.2) and 99.4% (96.6–100), respectively. Test characteristics of β -OHB were stable across a wide range of potential cutoffs (Table 1). The receiver-operator curve suggested that the optimal β -OHB cutoff is >2 mmol/L, where sensitivity remains at 98.1% but specificity improves to 82.3% (Supplementary Data).

CONCLUSIONS—Approximately one-fourth of emergency-department patients have diabetes, and almost half have poor glycemic control (2). Common clinical emergency-department practice is to consider the diagnosis of DKA in all patients presenting with blood glucose ≥ 250 mg/dL regardless of the reason for the visit. Because emergency-department overcrowding has stretched resources to the breaking point, there is increasing emphasis on developing, testing, and using

Table 1—Demographic and laboratory characteristics of the study sample

	DKA	No DKA
n (%)	54 (10.5)	462 (89.5)
Age (years)	41 (30–48)	48 (40–57)
Female (%)	27.8	35.3
Race/ethnicity (%)		
Non-Hispanic white	0	3.1
African American	16.7	13.6
Hispanic	79.2	80.2
Asian	4.2	2.6
Other	0.5	0
Admitted (%)	95.2	35.9
Newly diagnosed diabetes (%)	25.9	14.6
Insulin treated (%)	56.6	44.5
Venous pH	7.19 (7.10–7.25)	7.39 (7.36–7.41)
β -OHB (mmol/L)	0.3 (0.2–1.2)	4.9 (3.7–5.6)
HCO ₃ (mmol/L)	8 (6–11)	23 (21–26)
Glucose (mg/dL)	678 (448–955)	409 (328–409)
Anion gap (mmol/L)	29 (26–34)	13 (11–17)
Positive urine dipstick ketones (%)	98.1	64.9

Data are median (interquartile range), unless otherwise indicated.

the most rapid, accurate, and easy-to-use DKA screening tool available. The results of this study show that the specificity of capillary point-of-care β -OHB is superior to the urine dipstick.

Two retrospective emergency-department studies have shown that capillary point-of-care β -OHB is 100% sensitive for DKA (8,9). Charles et al. (10) likewise found that β -OHB was 100% sensitive in a prospective study of seven DKA patients. Naunheim et al. (11) reported a β -OHB sensitivity for DKA of 98% in a prospective sample of 160 patients (57 with DKA). Our study is the first to prospectively compare the test characteristics of point-of-care β -OHB against the urine dipstick in a cohort of emergency-department patients. Using β -OHB in lieu of the urine dipstick to screen our sample could have reduced full laboratory work-ups for DKA by 56.9%.

Our study has several limitations. First, we did not provide specific training to the nurses who interpreted the urine dipsticks. We felt this was unnecessary because they are 1) Clinical Laboratory Improvement Amendment certified, 2) familiar and comfortable with this process, and 3) do so regularly in routine clinical practice. Second, the urine dipstick was considered only negative or positive, and consideration of trace/small ketones versus moderate/large ketones may impact the findings. Third, we did not send blood samples to the laboratory for an enzymatic analysis to correlate the

values measured by the point-of-care β -OHB meter because, despite early concerns, current-generation β -OHB meters have been shown to be highly accurate compared with laboratory enzymatic analysis (6,14). Fourth, although there are many ways to define DKA, we chose to use the ADA definition because this is commonly used in DKA literature, allowing for direct comparison between our results and previous reports.

In conclusion, point-of-care capillary β -OHB testing at a cutoff of >1.5 mmol/L was equally sensitive (98.1%) but markedly more specific (78.6 vs. 35.1%) than the urine dipstick for the detection of DKA in our prospective sample of emergency-department patients with hyperglycemia. Utilizing point-of-care β -OHB testing has the potential to substantially reduce comprehensive laboratory evaluations for DKA among hyperglycemic emergency-department patients.

Acknowledgments—The authors thank Abbott Laboratories for their donation of the β -OHB test strips needed to complete this study.

No potential conflicts of interest relevant to this article were reported.

The authors alone had sole control of study design, data collection and analysis, and dissemination of findings. S.A. conceived of the project, helped design the study, and contributed to the writing and revising of the manuscript. S.O.H. helped design the study and assisted with collection of data. T.L. collected

and organized data and participated in the writing of the manuscript. M.M. helped design the study, performed all statistical analyses, and helped write and revise the manuscript.

Parts of this article were presented in abstract form at the President's Session of the 70th Scientific Sessions of the American Diabetes Association, Orlando, Florida, 25–29 June 2010.

References

1. Ginde AA, Pelletier AJ, Camargo CA Jr. National study of U.S. emergency department visits with diabetic ketoacidosis, 1993-2003. *Diabetes Care* 2006;29:2117–2119
2. Menchine MAS, Camargo CA, Ginde AA. Prevalence of undiagnosed and sub-optimally controlled diabetes by point-of-care hba1c in unselected emergency department patients. *Acad Emerg Med*. In press
3. American Diabetes Association. Tests of glycemia in diabetes. *Diabetes Care* 1998;21:S69–S71
4. Schwab TM, Hendey GW, Soliz TC. Screening for ketonemia in patients with diabetes. *Ann Emerg Med* 1999;34:342–346
5. American Diabetes Association. Clinical practice recommendations. *Diabetes Care* 2004;27:S94–S102
6. Byrne HA, Tieszen KL, Hollis S, Dornan TL, New JP. Evaluation of an electrochemical sensor for measuring blood ketones. *Diabetes Care* 2000;23:500–503
7. Kitabchi AE, Umpierrez GE, Miles JM, Fisher JN. Hyperglycemic crises in adult patients with diabetes. *Diabetes Care* 2009;32:1335–1343
8. Harris S, Ng R, Syed H, Hillson R. Near patient blood ketone measurements and their utility in predicting diabetic ketoacidosis. *Diabet Med* 2005;22:221–224
9. Taboulet P, Haas L, Porcher R, et al. Urinary acetoacetate or capillary beta-hydroxybutyrate for the diagnosis of ketoacidosis in the Emergency Department setting. *Eur J Emerg Med* 2004;11:251–258
10. Charles RA, Bee YM, Eng PH, Goh SY. Point-of-care blood ketone testing: screening for diabetic ketoacidosis at the emergency department. *Singapore Med J* 2007;48:986–989
11. Naunheim R, Jang TJ, Banet G, Richmond A, McGill J. Point-of-care test identifies diabetic ketoacidosis at triage. *Acad Emerg Med* 2006;13:683–685
12. Flahault A, Cadilhac M, Thomas G. Sample size calculation should be performed for design accuracy in diagnostic test studies. *J Clin Epidemiol* 2005;58:859–862
13. Hawass NE. Comparing the sensitivities and specificities of two diagnostic procedures performed on the same group of patients. *Br J Radiol* 1997;70:360–366
14. Wallace TM, Meston NM, Gardner SG, Matthews DR. The hospital and home use of a 30-second hand-held blood ketone meter: guidelines for clinical practice. *Diabet Med* 2001;18:640–645