



Ketone Body Infusion Increases Circulating Erythropoietin and Bone Marrow Glucose Uptake

<https://doi.org/10.2337/dc18-1421>

Katrine M. Lauritsen,^{1,2}
Esben Søndergaard,^{1,2,3} Mads Svart,^{1,2}
Niels Møller,^{1,3} and Lars C. Gormsen⁴

Sodium–glucose cotransporter 2 (SGLT2) inhibition was originally developed as a facile way of lowering blood glucose through increased renal elimination. However, clinical trials have demonstrated that SGLT2 inhibition not only improves glycemic control but also increases circulating ketone bodies and hematocrit (1,2). Recently, SGLT2 inhibitor–driven modest hyperketonemia has attracted much focus as a possible explanation of the improved cardiac outcome in patients treated with SGLT2 inhibitors (3), whereas the increased hematocrit has mostly been ascribed to hemoconcentration due to increased osmotic diuresis. In this observation, we present data indicating that hyperketonemia directly stimulates circulating erythropoietin concentrations and is associated with increased bone marrow activity assessed by ¹⁸F-fluorodeoxyglucose (FDG) positron emission tomography/computed tomography (PET/CT).

Seventeen healthy subjects undergoing either cardiac (study A) or brain (study B) PET/CT as part of protocols involving experimental hyperketonemia were investigated. Study A (4) consisted of 8 subjects (3 women) aged 60 (50–68) years with BMI 26 (22–35) kg/m², and study B (5) of 9 subjects (4 women) aged 62 (56–69) years with BMI 24 ± 2 kg/m². Both

studies were randomized, single-blinded crossover trials in which participants were infused with either 7.5% Na-3-β-hydroxybutyrate (3-OHB) (infusion rate 0.18 g/kg/h [study A] and 0.22 g/kg/h [study B]) or saline (0.9%). In study A, bone marrow palmitate and glucose uptake was measured with dynamic ¹¹C-palmitate and ¹⁸F-FDG PET (*n* = 8). Bone marrow time-radioactivity curves were measured in two vertebrae in the spine, and the input function was derived from either arterial samples (¹¹C-palmitate) or from a volume of interest drawn in the aorta (¹⁸F-FDG). Uptake rates were obtained using linear graphical analysis (Logan plot for ¹⁸F-FDG and Patlak analysis for ¹¹C-palmitate).

A serum sample from the end of each study day (study A: *t* = 390 min, study B: *t* = 240 min) was analyzed for erythropoietin by a chemiluminescent immunoassay on Immulite 2000 (Siemens Healthineers, Erlangen, Germany).

By design, 3-OHB concentrations increased rapidly during ketone body infusions (Fig. 1A), reaching 3.8 ± 0.2 mmol/L (study A) and 5.5 ± 0.4 mmol/L (study B). Ketone body levels were undetectable on saline study days.

As seen in Fig. 1C, erythropoietin concentrations were significantly greater

during hyperketonemia than after saline infusions (9.9 ± 1.1 vs. 7.6 ± 1.0 IU/L, *P* = 0.01). Hyperketonemia was also associated with 25% increased bone marrow ¹⁸F-FDG uptake compared with the saline study day (0.64 ± 0.08 vs. 0.51 ± 0.12 mL plasma/mL tissue, *P* = 0.04) (Fig. 1D) but did not affect bone marrow ¹¹C-palmitate uptake (1.7 ± 0.4 vs. 1.9 ± 0.6 μmol/g/min, *P* = 0.62). No correlation was observed between the increase in erythropoietin concentration and increase in bone marrow glucose uptake (*r* = −0.09, *P* = 0.84).

Some data supporting a connection between ketone bodies and erythropoietin exist. Thus, it has been shown that the hematocrit of infants born to mothers with diabetes is correlated with maternal 3-OHB concentrations and also that 4 weeks of SGLT2 inhibition by empagliflozin results in both increased levels of ketone bodies and a most-likely-transient increase in erythropoietin (2). In addition, dapagliflozin treatment causes not only hemoconcentration but also increased total red cell mass and transient increases in reticulocytes and erythropoietin.

Although the levels of hyperketonemia obtained in our experimental protocols were markedly greater than what has

¹Department of Endocrinology and Internal Medicine, Aarhus University Hospital, Aarhus, Denmark

²Danish Diabetes Academy, Odense University Hospital, Odense, Denmark

³Steno Diabetes Center Aarhus, Aarhus, Denmark

⁴Department of Nuclear Medicine and PET Centre, Aarhus University Hospital, Aarhus, Denmark.

Corresponding author: Lars C. Gormsen, lars.christian.gormsen@clin.au.dk.

Received 3 July 2018 and accepted 4 September 2018.

Clinical trial reg. nos. NCT02814474 (study A) and NCT02357550 (study B), clinicaltrials.gov.

© 2018 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.




Figure 1—A: 3-OHB concentrations during hyperketonemia (open boxes) and saline (closed circles). Error bars are SEM. EPO, erythropoietin. B: Representative CT and PET images (coronal and axial projections) of subject no. 8 on the ketone body infusion study day. PET images were summed from 5–50 min and represents average standardized uptake values (SUV). As seen, the bone marrow was clearly visible. C: Erythropoietin levels. D: Bone marrow glucose uptake (volume of distribution) assessed by ¹⁸F-FDG PET/CT.

been observed during SGLT2 inhibitor treatment (4–5 vs. ~0.6 mmol/L) (3), it is possible that a more modest but sustained increase in ketone bodies has a similar effect on erythropoietin levels. Erythropoietin is primarily produced in specialized renal cells, and increased secretion of the hormone in response to, e.g., altitude-induced hypoxia is not necessarily closely related to the stimulus. In other words, we may have only detected the initial increase in erythropoietin levels that the experimental hyperketonemia would eventually have produced.

Erythropoietin stimulates red blood cell maturation and release from the bone marrow, an energy-consuming process visible as avid ¹⁸F-FDG uptake on

PET/CT. The time to maximum effect of a single dose of human recombinant erythropoietin on reticulocyte cell count is 200–300 h, and it is therefore not surprising that we were unable to demonstrate a correlation between erythropoietin levels and bone marrow activity. Alternatively, the increase in bone marrow glucose uptake could be the result of increased leukopoiesis or thrombopoiesis or increased blood flow. We find this unlikely since 1) ketone bodies have been reported to be anti-inflammatory, 2) it takes several weeks of ketone body ingestion to increase thrombocytes discretely, and 3) bone marrow ¹¹C-palmitate uptake was unaffected.

In conclusion, experimental hyperketonemia results in ~30% concomitant

increase in erythropoietin levels and bone marrow glucose uptake. Although the level of hyperketonemia in our study was markedly higher than what is typically observed during SGLT2 inhibition, our data provide a possible explanation for the unknown link between SGLT2 inhibitor treatment and increased erythropoiesis.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Funding. The study was supported by grants from the Danish Council for Strategic Research, the Danish Council for Independent Research | Medical Sciences, Aase Danielsen Fund, Ferd. Og Ellen Hindsgaulls Fund, the Danish Medical Association, the Novo Nordisk Foundation, the Danish Diabetes Association, the Danish Diabetes

Academy supported by the Novo Nordisk Foundation, the Health Research Fund of Central Denmark Region, and the A.P. Møller Foundation.

Author Contributions. N.M. and L.C.G. conceptualized the study and wrote the protocol. K.M.L., E.S., M.S., and L.C.G. researched the data. E.S. contributed to discussion. K.M.L. wrote the manuscript, and E.S., M.S., N.M., and L.C.G. reviewed and edited the manuscript. N.M. and L.C.G. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Zinman B, Wanner C, Lachin JM, et al.; EMPA-REG OUTCOME Investigators. Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med* 2015;373:2117–2128
2. Ferrannini E, Baldi S, Frascerra S, et al. Renal handling of ketones in response to sodium–glucose cotransporter 2 inhibition in patients with type 2 diabetes. *Diabetes Care* 2017;40:771–776
3. Ferrannini E, Mark M, Mayoux E. CV protection in the EMPA-REG OUTCOME trial: a “thrifty substrate” hypothesis. *Diabetes Care* 2016;39:1108–1114
4. Gormsen LC, Svart M, Thomsen HH, et al. Ketone body infusion with 3-hydroxybutyrate reduces myocardial glucose uptake and increases blood flow in humans: a positron emission tomography study. *J Am Heart Assoc* 2017;6:e005066
5. Svart M, Gormsen LC, Hansen J, et al. Regional cerebral effects of ketone body infusion with 3-hydroxybutyrate in humans: reduced glucose uptake, unchanged oxygen consumption and increased blood flow by positron emission tomography. A randomized, controlled trial. *PLoS One* 2018;13:e0190556