

This article has been temporarily retracted, as it will be presented in abstract form at the 69th Scientific Sessions, 5–9 June 2009, New Orleans, Louisiana

For information about Scientific Sessions, the annual meeting of the American Diabetes Association, please visit <http://scientificsessions.diabetes.org>.

Serum uric acid levels improve prediction of incident Type 2 Diabetes in individuals with impaired fasting glucose: The Rancho Bernardo Study

Caroline Kaercher Kramer, MD^{1,2} Denise von Mühlen, MD, PhD¹
Simerjot Kaur Jassal, MD¹ Elizabeth Barrett-Connor, MD¹

¹Division of Epidemiology, Department of Family and Preventive Medicine, School of Medicine, University of California, San Diego, La Jolla, California

²Division of Endocrinology, Hospital de Clinicas de Porto Alegre, RS, Brazil

Correspondence and Reprint requests:

Elizabeth Barrett-Connor, MD

Email: ebarrettconnor@ucsd.edu

Submitted 12 February 2009 and accepted 25 March 2009.

This is an uncopyedited electronic version of an article accepted for publication in *Diabetes Care*. The American Diabetes Association, publisher of *Diabetes Care*, is not responsible for any errors or omissions in this version of the manuscript or any version derived from it by third parties. The definitive publisher-authenticated version will be available in a future issue of *Diabetes Care* in print and online at <http://care.diabetesjournals.org>.