

Supplemental Table

Linear association analysis among metabolic/vascular variables and pathogen seropositivities in the study subjects.

<i>Variable</i>	HSV-1 Titer	HSV-2 Titer	Chlamydia titer	Enterovirus titer	Semi- quantitative score	Quantitative index
Age	r=0.18 p=0.04	r=0.09 p=NS	r=-0.02 p=NS	r=-0.22 p=0.01	r=0.11 p=NS	r=0.12 p=NS
Body mass index	r=0.11 p=NS	r=0.18 p=0.04	r=0.03 p=NS	<i>r=0.15</i> <i>p=0.07</i>	r=0.009 p=NS	r=0.13 p=NS
Waist-to-hip ratio	r=0.11 p=NS	r=0.09 p=NS	r=0.06 p=NS	r=0.05 p=NS	r=0.03 p=NS	r=0.12 p=NS
Systolic blood pressure	r=0.14 p=0.1	r=0.03 p=NS	r=0.04 p=NS	r=0.24 p=0.009	<i>r=0.16</i> <i>p=0.06</i>	r=0.24 p=0.01
Diastolic blood pressure	r=0.20 p=0.03	<i>r=0.15</i> <i>p=0.07</i>	r=0.01 p=NS	r=0.18 p=0.05	r=0.13 p=NS	r=0.22 p=0.02
Fasting glucose	r=0.02 p=NS	r=0.20 p=0.03	r=0.05 p=NS	r=0.06 p=NS	r=0.04 p=NS	r=0.09 p=NS
HbA _{1c}	r=0.08 p=NS	r=0.09 p=NS	r=0.01 p=NS	r=0.007 p=NS	r=0.06 p=NS	r=0.06 p=NS
Total cholesterol	<i>r=0.16</i> <i>p=0.06</i>	r=0.12 p=NS	r=0.009 p=NS	r=0.14 p=NS	r=0.006 p=NS	r=0.08 p=NS
LDL-cholesterol	r=0.08 p=NS	r=0.04 p=NS	r=0.01 p=NS	r=0.07 p=NS	r=0.008 p=NS	r=0.12 p=NS
HDL-cholesterol	r=-0.07 p=NS	r=-0.01 p=NS	r=0.02 p=NS	r=-0.04 p=NS	r=0.09 p=NS	r=0.11 p=NS
Triglycerides	r=0.08 p=NS	r=0.05 p=NS	R=0.02 p=NS	r=0.12 p=NS	r=0.02 p=NS	r=0.07 p=NS
C-reactive protein	r=0.04 p=NS	r=0.04 p=NS	r=0.04 p=NS	r=0.11 p=NS	r=0.06 p=NS	r=0.01 p=NS
sTNFR1	r=0.01 p=NS	r=0.19 p=0.03	r=0.14 p=NS	r=0.19 p=0.03	r=0.09 p=NS	r=0.07 p=NS
sTNFR2	r=0.01 p=NS	r=0.05 p=NS	r=0.09 p=NS	r=0.19 p=0.03	r=0.12 p=NS	r=0.03 p=NS
Insulin sensitivity	r=-0.05 p=NS	r=-0.29 p=0.002	r=0.005 p=NS	r=0.09 p=NS	r=0.03 p=NS	r=-0.13 p=NS
Insulin secretion	r=-0.05 p=NS	r=-0.03 p=NS	r=-0.07 p=NS	r=0.04 p=NS	r=0.05 p=0.NS	r=-0.06 p=NS