



Update on Insulinomas or The Case of the Missing (Pro)Insulinoma

The diagnosis of hypoglycemia with insulin hypersecretion is one of the most intriguing challenges facing physicians. The entities to exclude constitute a host of rare organic maladies and less rare psychologic maladies, among them "lay-press-induced hypoglycemia." The possibility of an insulinoma formed a major plot element in a recent novel in which murder was suspected and the detective was "a top Washington endocrinologist" who had done diabetes research and "thanked (his) lucky stars (he) didn't have to write another grant application (1)." In some instances, the diagnosis can be established on clinical grounds without resort to the laboratory. A case in point is a recent patient with abrupt onset of profound and recurrent hypoglycemia. One of his pharmacy bottles—labeled acetazolamide—was found, after initiating the workup, to contain acetohexamide and the pharmacy was able to confirm its error. Frequently, however, physicians are quite dependent on the laboratory and provocative testing for establishing the diagnosis in these patients.

The gold standard for the biochemical diagnosis of insulinoma remains the 72-h fast (2,3). The presence of detectable circulating insulin in the face of symptomatic hypoglycemia has been regarded as the end point, yet in research-quality insulin assays that permit the detection of ≤ 1 $\mu\text{U}/\text{ml}$, that criterion is not appropriate. Ser-

vice et al. (3) have recommended a threshold of 6 $\mu\text{U}/\text{ml}$ (36 pM) for inappropriate hyperinsulinism in the face of hypoglycemia. If the patient becomes hypoglycemic within the first few hours, the fast is relatively straightforward to perform and interpret. However there is easily a role for an accurate test that is simple and inexpensive compared to: 1) the discomfort for the patient and logistical problems of close supervision and frequent sampling by health-care personnel in a prolonged fast not resolved within the first few hours; or 2) a variety of provocative tests (e.g., tolbutamide injection, calcium infusion, C-peptide suppression either by insulin infusion to hypoglycemia or by hyperinsulinemic-euglycemic clamp) that are labor intensive, sometimes risky, and usually require a research environment. Current economic pressures on the use of days of hospitalization and the diagnostic uncertainties alluded to above further support the potential benefits of such a test. A screening test may prove capable of making a biochemical diagnosis of inappropriate β -cell hypersecretion, possibly distinguishing degrees of tissue differentiation (carcinoma, adenoma, "hyperplasia") or at least establishing a level of certainty for excluding the diagnosis and reducing the number of patients who will require extensive evaluation.

Pre- or intraoperative localization, including intraoperative ultrasound, may permit enucleation of a discrete tumor. If the biochemical diagnosis is made but no discrete tumor is anatomically identifiable, up to 90% pancreatic resection is the recommended procedure in one major surgical text (4) but is considered controversial in another (5). The appropriateness of extensive resection when tumor cannot be identified, even in the operating room, deserves reevaluation in 1988, given the possibility of resulting diabetes, the possible need for postoperative medical therapy in any case, and new

From the Division of Endocrinology/Metabolism, University of Cincinnati Medical Center, Cincinnati, Ohio.

Address correspondence and reprint requests to Robert M. Cohen, MD, Division of Endocrinology/Metabolism, University of Cincinnati Medical Center, 231 Bethesda Avenue, Cincinnati, OH 45267-0547.

therapeutic options. The total operative mortality of up to 8–10% for islet tumor surgery (5) and the report that up to 24% of patients will not have hypoglycemia relieved by a single operation (6) should give the referring physician pause. In cases when a diagnosis has not been made even in the operating room, it may remain unclear even after the tissue is on the pathologist's table.

Using histomorphometric techniques, Goudswaard et al. (7) compared 1) pancreases from normal fetuses, neonates, children up to age 3 yr, adults age 44–71 yr and children with hyperinsulinism; and 2) the nontumor tissue from eight adults with benign insulin-producing adenomas. They concluded "it seems that any morphological pattern, including a normal pattern, may be identified as nesidioblastosis [or islet cell hyperplasia], as long as the clinical condition fulfills the criteria for organic hyperinsulinism" (7). The meaning and clinical utility of the pathologic diagnosis of islet cell hyperplasia, in the absence of standards by which to compare the response of the normal human β -cell mass not only to age but to obesity and perhaps other factors, are uncertain. There have not been adequate studies of the response of islet cell hyperplasia to single and combined medical therapies including not only diazoxide but also new therapeutic options such as a long-acting somatostatin analog (8,9) and the calcium channel blockers verapamil and diltiazem, which have been reported anecdotally to control hypoglycemia in some patients (10–12). If it becomes possible to treat these patients medically, it will become important to determine that inappropriate insulin hypersecretion is due to an entity other than a discrete tumor.

Since the discovery of proinsulin, studies from a number of laboratories using various earlier assay techniques have demonstrated abnormalities in serum proinsulin in patients with insulinomas (reviewed in refs. 13 and 14). Major technical improvements in proinsulin measurement have occurred in the last few years as a result of the availability of biosynthetic human proinsulin (15–23). The newer assays probably will provide an improvement over the previous assays in the diagnosis of β -cell hypersecretion, partly by virtue of better-defined specificity and partly by making proinsulin measurements easier and more available. Still, there are several subtleties to the interpretation of any proinsulin assay for this use. C-peptide is present in the circulation at ~ 100 times and insulin at ~ 20 times the concentration of proinsulin. For proinsulin measurements to be helpful in difficult cases, the contribution of either C-peptide or insulin to the total proinsulin immunoreactivity should be $< 1\%$. Therefore C-peptide cross-reactivity in the proinsulin assay should be $< 0.01\%$ (on a molar basis) and insulin cross-reactivity $< 0.05\%$. Also, the extent and pattern of cross-reactivity of the intermediates in the process of conversion of proinsulin to insulin need to be taken into account. They may contribute to variations in the normal range between laboratories and in the correlation between measured hormone concentration and the degree of hypoglycemia observed.

We had the opportunity to measure samples from 20 patients with proven tumors in a specific proinsulin radioimmunoassay. The fasting serum proinsulin concentration from patients with β -cell tumors were higher than those from normal and obese control subjects by $> 3SD$ (17), a degree of discrimination not achieved by measuring fasting insulin alone. Heding and Kruse (24) reported in abstract form in 1984 that 204 of 206 patients with islet tumors had abnormal plasma proinsulin concentrations, a remarkable sensitivity of 99%. In a small series, mean circulating percent proinsulin (by gel filtration) differed between histologically defined subsets which in turn predicted the response to hormonal/pharmacologic intervention (25). But because of the rarity of these tumors, none of these laboratories has had the opportunity to establish 1) both sensitivity and specificity of proinsulin measurements for the diagnosis of hypoglycemia with insulin hypersecretion in the context of adequate clinical and pathologic data, 2) the relationship of proinsulin results to the islet pathology, or 3) the usefulness of postsurgical proinsulin measurements in predicting recurrent hypoglycemia in a large series of patients. These questions remain to be answered in the coming years and we look forward to seeing how, in the long run, the use of proinsulin measurements will contribute to unravelling the mysteries of hypoglycemia.

ROBERT M. COHEN, MD
FLORENCE CAMUS, MD

REFERENCES

1. Ravin N: *Seven North*. New York, Jove, 1985
2. Sherwin RS, Felig P: Hypoglycemia. In *Endocrinology and Metabolism*. 2nd ed. Felig P, Baxter JD, Broadus AE, Frohman LA, Eds. New York, McGraw-Hill, 1987, p. 1179–202
3. Service FJ, van Heerden JA, Sheedy PF: Insulinoma. In *Hypoglycemic Disorders*. Service FJ, Ed. Boston, GK Hall, 1983, p. 111–29
4. Bradley EL III, Zeppa R: The Pancreas. In *Textbook of Surgery*. 13th ed. Sabiston DC, Ed. Philadelphia, PA, Saunders, 1986, p. 1196–97
5. Silen W, Steer ML: The Pancreas. In *Principles of Surgery*. 4th ed. Schwartz SI, Shires GT, Spencer FC, Storer EH, Eds. New York, McGraw-Hill, 1984, p. 1365–67
6. Stefanini P, Carboni M, Patrassi N, Basoli A: Beta-islet cell tumors of the pancreas: results of a study on 1067 cases. *Surgery* 75:597–609, 1974
7. Goudswaard WB, Houthoff HJ, Koudstaal J, Zwierstra RP: Nesidioblastosis and endocrine hyperplasia of the pancreas: a secondary phenomenon. *Hum Pathol* 17:46–54, 1986
8. Osei K, O'Dorisio TM: Malignant insulinoma: effects of a somatostatin analog (compound 201-995) on serum glucose, growth, and gastro-entero-pancreatic hormones. *Ann Intern Med* 103:223–25, 1985
9. Kvols LK, Buck M, Moertel CG, Schutt AJ, Rubin J, O'Connell MJ, Hahn RG: Treatment of metastatic islet cell carcinoma with a somatostatin analog (SMS 201-995). *Ann Intern Med* 107:162–68, 1987
10. Taniguchi H, Murakami, Morita S, Kazumi T, Yoshino G,

- Maeda M, Baba S: Calcium antagonist (diltiazem) for reversal of hypoglycemic symptoms in insulinoma (Letter). *Lancet* 2:501, 1977
11. De Marinis L, Barbarino A: Calcium antagonists and hormone release. I. Effects of verapamil on insulin release in normal subjects and patients with islet-cell tumor. *Metabolism* 29:599–604, 1980
 12. Ulbrecht JS, Schmeltz R, Aarons JH, Greene DA: Insulinoma in a 94-year-old woman: long-term therapy with verapamil. *Diabetes Care* 9:186–88, 1986
 13. Robbins DC, Tager HS, Rubenstein AH: Biological and clinical importance of proinsulin. *N Engl J Med* 310:1165–75, 1984
 14. Heding LG, Faber O, Kasperska-Czyzłowa T, Sestoft L, Turner R: Radioimmunoassay of proinsulin and hyperproinsulinemic states. In *Proinsulin, Insulin and C-Peptide*. Baba S, Kaneko T, Yanaihara N, Ed. Amsterdam, Excerpta Med., 1979, p. 254–61
 15. Frank BH, Pettee JM, Zimmerman RE, Burck PJ: The production of human proinsulin and its transformation to human insulin and C-peptide. In *Peptides. Synthesis-Structure-Function, Proc 7th American Peptide Symp.* Rich DE, Gross E, Eds. Madison, WI, Pierce Chemical, 1982, p. 729–38
 16. Cohen RM, Nakabayashi T, Blix PM, Rue PA, Shoelson SE, Root MA, Frank BH, Revers RR, Rubenstein AH: A radioimmunoassay for circulating human proinsulin. *Diabetes* 34:84–91, 1985
 17. Cohen RM, Given BD, Licinio-Paixao J, Provow SA, Rue PA, Frank BH, Root MA, Polonsky KS, Tager HS, Rubenstein AH: Proinsulin radioimmunoassay in the evaluation of insulinomas in familial hyperproinsulinemia. *Metabolism* 35:1137–46, 1986
 18. Deacon CF, Conlon JM: Measurement of circulating human proinsulin concentrations using a proinsulin-specific antiserum. *Diabetes* 34:491–97, 1985
 19. Hartling SG, Dinesen B, Kappelgard A-M, Faber OK, Binder C: ELISA for human proinsulin. *Clin Chem Acta* 156:289–97, 1986
 20. Ward WK, Paquette TL, Frank BH, Porte D: A sensitive radioimmunoassay for human proinsulin, with sequential use of antisera to C-peptide and insulin. *Clin Chem* 32:728–33, 1986
 21. Gray IP, Siddle K, Frank BH, Hales CN: Characterization and use in immunoradiometric assay of monoclonal antibodies directed against human proinsulin. *Diabetes* 36:684–88, 1987
 22. Naylor BA, Matthews DR, Turner RC: A soluble-phase proinsulin radioimmunoassay and its use in diagnosis of hypoglycemia. *Ann Clin Biochem* 24:352–63, 1987
 23. Bowsher RR, Apathy JM, Ferguson AL, Frank BH: An improved radioimmunoassay (RIA) for proinsulin in human serum and plasma (Abstract). *Diabetes* 36:94A, 1987
 24. Heding LG, Kruse V: Usefulness of fasting proinsulin in the diagnosis of insulinoma (Abstract). *Diabetes* 33:148A, 1984
 25. Berger M, Bordi C, Cüppers H-J, Berchtold P, Gries FA, Münterfering H, Sailer R, Zimmermann H, Orci L: Functional and morphologic characterization of human insulinomas. *Diabetes* 32:921–31, 1983