

Errata

Kelley DE, Bray GA, Pi-Sunyer FX, Klein S, Hill J, Miles J, Hollander P: Clinical efficacy of orlistat therapy in overweight and obese patients with insulin-treated type 2 diabetes. *Diabetes Care* 25:1033–1041, 2002

Figures 3 and 4 of the above-listed article were mistakenly switched. The correct figures and legends are shown below. Please disregard the original erratum published in *Diabetes Care* 25:1495, 2002.




Figure 3—Fasting serum glucose levels over time with placebo (○) or 120 mg orlistat (●). $P = 0.02$, least-squares mean difference from placebo in the change from baseline over 52 weeks.




Figure 4—HbA_{1c} over 1 year of double-blind treatment with placebo (○) or 120 mg orlistat (●). $P = 0.002$, least-squares mean difference from placebo in the change from baseline over 52 weeks.

Arday DR, Fleming BB, Keller DK, Pendergrass PW, Vaughn RJ, Turpin JM, Nicewander DA: Variation in diabetes care among states: do patient characteristics matter? *Diabetes Care* 25:2230–2237, 2002

In the above-listed article, the URL for the online appendix is incorrect. The online appendix can be accessed via <http://care.diabetesjournals.org>.

American Diabetes Association: Standards of medical care for patients with diabetes mellitus (Position Statement). *Diabetes Care* 26 (Suppl. 1):S33–S50, 2003

Please note that the online version of this ADA position statement has been revised to reflect a correction in the reference list. The corrected version can be found at http://care.diabetesjournals.org/cgi/content/full/26/suppl_1/s33.