

Diabetes Reporting as a Cause of Death

Results from the Translating Research Into Action for Diabetes (TRIAD) study

LAURA N. McEWEN, PHD¹
 CATHERINE KIM, MD, MPH²
 MARY HAAN, MPH, DRPH³
 DEBASHIS GHOSH, PHD⁴
 PAULA M. LANTZ, PHD, MS⁵
 CAROL M. MANGIONE, MD, MSPH⁶

MONIKA M. SAFFORD, MD⁷
 DAVID MARRERO, PHD⁸
 THEODORE J. THOMPSON, MS⁹
 WILLIAM H. HERMAN, MD, MPH¹
 THE TRIAD STUDY GROUP*

OBJECTIVE — To determine the frequency of reporting of diabetes on death certificates of decedents with known diabetes, define factors associated with reporting of diabetes, and describe trends in reporting over time.

RESEARCH DESIGN AND METHODS — Data were obtained from 11,927 participants with diabetes who were enrolled in the Translating Research Into Action for Diabetes study, a multicenter prospective observational study of diabetes care in managed care. Data on decedents ($n = 540$) were obtained from the National Death Index. The primary dependent variable was the presence of ICD-10 codes for diabetes on the death certificate. Covariates included age at death, sex, race/ethnicity, education, income, duration of diabetes, type of diabetes, diabetes treatment, smoking status, and number of comorbidities.

RESULTS — Diabetes was recorded on 39% of death certificates and as the underlying cause of death for 10% of decedents with diabetes. Diabetes was significantly less likely to be reported on the death certificates of decedents with diabetes dying of cancer. Predictors of recording diabetes anywhere on the death certificate included longer duration of diabetes and insulin treatment. Longer duration of diabetes, insulin treatment, and fewer comorbidities were associated with recording of diabetes as the underlying cause of death.

CONCLUSIONS — Diabetes is much more likely to be reported on the death certificates of diabetic individuals who die of cardiovascular causes. Reporting of diabetes on death certificates has been stable over time. Death certificates underestimate the prevalence of diabetes among decedents and present a biased picture of the causes of death among people with diabetes.

Diabetes Care 29:247–253, 2006

Death certificates are commonly used to rank leading causes of death, estimate all-cause and cause-specific mortality, and describe trends in mortality over time (1–3). Each year the National Center for Health Statistics publishes a report on the leading causes of death in the U.S. and years of life lost (4–

6). These data are used by agencies such as the National Institutes of Health and the Centers for Disease Control and Prevention to define the burden of disease in the U.S. and to determine funding levels (7).

There are three ways that diabetes can be recorded on a death certificate (8). In part I, diabetes may be recorded as either the underlying cause of death or as an antecedent condition, and in part II, diabetes may be recorded as a significant condition contributing to death but not resulting in the underlying cause given in part I (10). National mortality estimates are based on a simple analysis of the underlying cause of death listed on death certificates.

There are many problems related to the reliability and validity of cause-of-death information on death certificates. These problems arise from inaccuracy of diagnosis, variation in interpreting causal sequences and conditions contributing to death, changing perceptions of the causal role of diseases, variations in nosological coding, lack of training in death certificate completion, and improper completion of death certificates (8,9). Coding of cause of death is especially problematic when a decedent has multiple chronic conditions because a single disease may not adequately describe the cause of death.

Clearly, not everyone with diabetes dies of diabetes. Studies dating back to the 1970s have shown that diabetes is often not recorded as a cause of death for people with diabetes (8,9,11–21). For decedents with a history of diabetes, diabetes was listed anywhere on the death certificate only 38% of the time in 1986 and 36% of the time in 1993 (16,17).

Individuals who have diabetes recorded as a cause of death may not be representative of all decedents with diabetes. For example, in the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study, diabetes was recorded as an underlying cause of death among women with diabetes (21%) more frequently than among men with diabetes (7%) (22). Hispanic or Latino heritage may also be associated with more frequent recording of diabetes on death

From the ¹Departments of Internal Medicine and Epidemiology, University of Michigan, Ann Arbor, Michigan; the ²Departments of Internal Medicine and Obstetrics and Gynecology, University of Michigan, Ann Arbor, Michigan; the ³Department of Epidemiology, University of Michigan, Ann Arbor, Michigan; the ⁴Department of Biostatistics, University of Michigan, Ann Arbor, Michigan; the ⁵Department of Health Management and Policy, University of Michigan, Ann Arbor, Michigan; the ⁶Department of Medicine, University of California, Los Angeles, California; the ⁷Deep South Center on Effectiveness at the Birmingham Veterans Affairs Medical Center and the University of Alabama at Birmingham, Birmingham, Alabama; the ⁸Indiana University School of Medicine, Indianapolis, Indiana; and the ⁹Division of Diabetes Translation, Centers for Disease Control and Prevention, Atlanta, Georgia.

Address correspondence and reprint requests to William H. Herman, MD, MPH, Internal Medicine/Metabolism, Endocrinology, and Diabetes, 1500 East Medical Center Dr., 3920 Taubman Center, Ann Arbor, MI 48109-0354. E-mail: wherman@umich.edu.

Received for publication 1 June 2005 and accepted in revised form 9 November 2005.

*A complete list of TRIAD Study Group members can be found in the APPENDIX.

Abbreviations: NDI, National Death Index; TRIAD, Translating Research Into Action for Diabetes.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2006 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

certificates, especially when the death is due to cardiovascular disease (8,23).

Will et al. (17) have hypothesized that diabetes reporting on death certificates may improve because of the increasing prevalence of diabetes, highly publicized results of major clinical trials, and increased media coverage of diabetes. To determine the frequency of reporting of diabetes on death certificates of decedents with known diabetes, the factors associated with more complete reporting of diabetes, and whether recording practices have changed over time, we examined the reporting of diabetes on death certificates in the Translating Research Into Action for Diabetes (TRIAD) study, a multicenter prospective observational study of diabetes care in managed care.

RESEARCH DESIGN AND METHODS

— TRIAD has been described in detail elsewhere (24). In brief, six Translational Research Centers collaborated with 10 managed care health plans and 68 provider groups that serve ~180,000 people with diabetes. The health plans are geographically and ethnically diverse (Hawaii, California, Texas, Indiana, Michigan, Pennsylvania, and New Jersey). The TRIAD study population consists of a stratified, random sample of adults with diabetes from the patient populations of these health plans. Study participants were ≥ 18 years of age, were community-dwelling, were not pregnant, were English- or Spanish-speaking, and had been continuously enrolled in the health plan for at least 18 months and had made at least one claim for health services during the previous 18 months. Participants were sampled from provider groups that had at least 50 participants with diabetes enrolled in the health plan. Recruitment began in 1999 and was completed in September 2001. The study protocol was reviewed and approved by the institutional review boards at all six Translational Research Centers. All participants provided informed consent.

We used a survey to collect information from TRIAD participants. The survey was administered by computer-assisted telephone interview or in writing by mail. We also reviewed medical records. We obtained information on TRIAD decedents from National Death Index (NDI) Plus searches (25). All deaths were verified by matching name, social security number (available for ~52% of participants), date of birth, and sex of the dece-

dent with data supplied by the NDI. The sensitivity of NDI has been shown to range from 87 to 98% (26). Different combinations of identifiers excluding social security number correctly identify 83–92% of deceased individuals and 92–99% of living individuals, making NDI an accurate source of ascertaining vital status even without social security number (27).

Vital status was determined for all TRIAD participants ($n = 11,927$) as of 1 January 2003. All information regarding ICD-10 codes for the underlying and contributing causes of death were derived from the NDI file. For our analyses, we included all TRIAD participants ($n = 11,927$), 540 (4.5%) of whom were identified through NDI, as having died before 1 January 2003.

Main outcome measures and covariates

We used the American Heart Association definitions to group causes of death according to underlying cause (28). We investigated two dichotomous dependent variables: the presence of diabetes as defined by ICD-10 codes E10–E14 on the death certificates as either the underlying cause of death or as a cause appearing anywhere on the death certificate. Covariates included age at death, sex, race/ethnicity, education, income, age at diagnosis of diabetes, duration of diabetes, type of diabetes, diabetes treatment, smoking status, and number of comorbidities. Type 1 diabetes was defined by insulin use (without the use of oral antidiabetic agents) and age at diagnosis of diabetes ≤ 30 years; all others were defined as having type 2 diabetes (29). Number of comorbidities was defined using the Charlson index, a weighted measure of comorbid conditions associated with mortality (30). The Charlson Index has been extensively studied and provides an accurate and valid measure of comorbidity (31).

Statistical analyses

Bivariate analyses were performed for all-cause mortality and separately for cardiovascular (ICD-10 codes I00–I99) and noncardiovascular mortality (all other ICD-10 codes). Unadjusted odds ratios were constructed using each variable singularly in a logistic model predicting either diabetes as the underlying cause of death or diabetes listed anywhere on the death certificate.

To simultaneously adjust for covariates that predict the recording of diabetes

on death certificates, we constructed multivariable models. We modeled diabetes as the underlying cause of death, and diabetes listed anywhere on the death certificate separately. We also modeled diabetes listed anywhere on the death certificate separately for cardiovascular and noncardiovascular causes. Single imputations were generated for missing values for covariates from the TRIAD patient survey using the transcan function in S-PLUS edition 6.1 (Insightful, Seattle, WA). Variables obtained through chart review were not imputed. For multivariable analyses, we excluded all TRIAD decedents who did not consent to medical record review ($n = 173$, 32%, excluded). Those who were excluded from the multivariable models were more likely to be nonwhite, to have less education, and to not use insulin for diabetes treatment. Before constructing the models, we assessed a correlation matrix with exposure variables. Age at diagnosis of diabetes, duration of diabetes, type of diabetes, and age at death were correlated at Spearman's $r > 0.5$. To decrease colinearity between predictors in the multivariable models, we only included duration of diabetes because it had the strongest bivariate association with each outcome. Although education and income were correlated at Spearman's $r > 0.5$, we included both in the multivariable models because they may have different predictive properties when modeling health-related outcomes (32).

In constructing the models, we used multivariable logistic regression, first including the variables that were found to be significant in bivariate tests of association between predictors and outcomes. For models of diabetes listed anywhere on the death certificate, we had a large sample size and were able to include variables that were significantly associated with mortality in previous studies (sex and race/ethnicity) (22,23,33) and variables that might explain the differences previously seen in mortality by race/ethnicity (education and income) or variables that might account for the differences seen in bivariate tests of association (Charlson index and smoking). For models of diabetes as the underlying cause of death and for models of diabetes as any listed cause separated by cardiovascular and noncardiovascular causes, sample size was smaller and we included only duration of diabetes, treatment for diabetes, and Charlson Index. All models included variables for TRIAD site to account for possible geo-

Table 1—Distribution of underlying causes of death for TRIAD decedents (n = 540) and odds ratio (95% CI) of recording diabetes anywhere on the death certificate using cardiac causes as the reference group, 2000–2002

Underlying cause of death	ICD-10 codes	Percent of deaths by underlying cause of death	Odds ratio of having diabetes recorded anywhere	Lower 95% CI	Upper 95% CI
Cardiac	I00–09, I11, I13, I20–25, I27, I30–52	36%	1.00	—	—
Cancer	C00–99	21%	0.43*	0.25	0.73
Other	all other	12%	0.56	0.30	1.03
Diabetes	E10–14	10%	—	—	—
Cerebrovascular disease	I60–69	6%	0.71	0.32	1.59
Respiratory	J00–09, J19–99	5%	0.89	0.54	2.33
All other cardiovascular disease	I10, I12, I14–19, I28–29, I53–59, I70–99	3%	1.16	0.42	3.25
Accidents, suicide, assault	V00–99, W00–99, X00–99, Y00–99	2%	0.27	0.06	1.26
Infections	A00–99, B00–99	2%	0.27	0.06	1.26
Renal failure	N17–19	2%	0.33	0.07	1.58
Influenza and pneumonia	J10–18	1%	0.75	0.13	4.18

* $P \leq 0.05$.

graphic or organizational differences in reporting diabetes on death certificates. In separate analyses, we used hierarchical logistic regression models (SAS GLIMMIX Macro with penalized quasi-likelihood estimation method [34]) with random intercepts for health plans and provider groups to account for the clustered study design (health plan, provider group, and participant levels) and the correlation among participant characteristics within health plans and provider groups. We did not find significant clustering at any level, so the results are not shown.

All analyses were performed using SAS version 8.02 (Research Triangle Institute, Research Triangle Park, NC).

RESULTS— Ten percent of decedents had diabetes recorded as the underlying cause of death, and 39% had diabetes recorded anywhere on their death certificates. Table 1 shows the distribution of underlying causes of death and the odds of recording diabetes anywhere on the death certificate by underlying cause of death. Compared with dying of cardiac disease, diabetes was recorded more frequently for those who died of all other cardiovascular disease, although this difference was not statistically significant. The more frequent recording of death due to all other cardiovascular disease was driven by those who died of hypertension or hypertensive renal disease, of whom 80% had diabetes recorded on the death certificate. For all other causes of death, diabetes was reported less frequently than

it was for cardiac disease. This result was statistically significant only when the cause of death was cancer (Table 1).

The unadjusted characteristics associated with recording diabetes as the underlying cause of death were treatment with insulin and having fewer comorbidities (Table 2). The only factors that were significantly associated with recording diabetes anywhere on the death certificate were treatment with insulin and diabetes duration 5–14 years and ≥ 15 years (Table 2).

When we looked only at those who died of cardiovascular causes, no unadjusted characteristics were significantly associated with recording of diabetes anywhere on the death certificate. Among people who died of noncardiovascular causes, only treatment with insulin was significantly associated with recording of diabetes.

In multivariable analyses of underlying cause of death, longer duration of diabetes and having fewer comorbidities were associated with the recording of diabetes (Table 2). In multivariable analyses of all-cause mortality, we found that recording of diabetes anywhere on the death certificate was only associated with longer duration of diabetes (Table 2).

In models of cardiovascular mortality, no variables predicted recording of diabetes anywhere on the death certificate. In models of noncardiovascular mortality, longer duration of diabetes was associated with the recording of diabetes

anywhere on the death certificate (Table 2).

CONCLUSIONS— Attribution of death to diabetes is often a subjective judgment on the part of certifying physicians (10). Physicians may not record diabetes on the death certificates of decedents because they may not have known that the decedent was diagnosed with diabetes before death, may not have believed that diabetes contributed to the decedent's death, or may not have listed diabetes because of space constraints. The standard death certificate includes only three to four lines for part I and one to two lines for part II. People with diabetes may have multiple chronic conditions such as hypertension, dyslipidemia, cardiovascular disease, and renal disease, which may compete with diabetes for space on the death certificate.

The results of our study that used knowledge of physician-diagnosed diabetes to determine diabetes status are consistent with those of previous studies that used data from the 1986 and 1993 National Mortality Follow-Back Survey and information from next of kin or personal informants. We found that 10% of diabetic decedents had diabetes recorded as the underlying cause of death, and 39% had diabetes recorded anywhere on their death certificates. Earlier studies found that 10 and 11% of diabetic decedents had diabetes recorded as the underlying cause of death, and 38 and 36% had diabetes recorded anywhere (16,17). Al-

Table 2—Factors associated with the recording of diabetes on death certificates and ORs and their associated 95% CIs in the TRIAD decedent population, 2000–2002

Characteristic	n	Diabetes listed as the underlying cause of death on death certificate (%)		Adjusted OR (95% CI)	Diabetes listed anywhere on death certificate (%)	Unadjusted OR (95% CI)	Fully adjusted OR (95% CI)	Adjusted OR (95% CI) for those who died of CVD causes predicting diabetes listed anywhere on death certificate	Adjusted OR (95% CI) for those who died of non-CVD causes predicting diabetes listed anywhere on death certificate
		Entire cohort of deceased subjects	Diabetes listed as the underlying cause of death on death certificate (%)						
Age at death (mean)	540	—	—	367	—	1.01 (0.99, 1.02)	367	164	203
Sex (Referent: Male)	70.3	—	0.99 (0.96–1.01)	†	—	*	*	†	†
Female	47	8	0.71 (0.40–1.27)	†	37	0.88 (0.62–1.24)	0.77 (0.49–1.23)	†	†
Race (referent: white)									
Hispanic	14	11	1.16 (0.50–2.69)	†	38	0.98 (0.57–1.67)	0.74 (0.33–1.65)	†	†
Black	23	13	1.39 (0.71–2.71)		40	1.09 (0.71–1.69)	1.32 (0.70–2.48)		
Other	16	6	0.58 (0.22–1.58)		37	0.94 (0.57–1.55)	1.62 (0.69–3.82)		
Education (referent: some high school or less)				†				†	†
High school graduate	26	10	1.23 (0.60–2.53)		39	1.08 (0.69–1.68)	0.98 (0.54–1.77)		
College	35	11	1.24 (0.63–2.41)		41	1.18 (0.78–1.76)	1.08 (0.59–2.00)	†	†
Income (referent: <\$15,000)				†				†	†
\$15,000–\$40,000	34	7	0.65 (0.33–1.28)		37	0.91 (0.61–1.34)	0.84 (0.49–1.44)		
≥\$40,000	18	13	1.21 (0.59–2.50)		40	1.04 (0.64–1.67)	1.54 (0.75–3.15)	†	†
Age at diabetes diagnosis (referent: <30 years)				†			*	†	†
≥30 years	93	10	0.92 (0.31–2.70)		38	0.85 (0.44–1.66)			
Duration of diabetes (referent: <5 years)									
5–14 years	42	11	1.87 (0.74–4.73)		41	1.90 (1.14–3.18)	2.31 (1.16–4.59)	0.74 (0.27–2.03)	6.50 (2.27–18.65)
15+ years	39	11	1.93 (0.76–4.89)		42	1.94 (1.16–3.37)	2.12 (1.02–4.40)	0.77 (0.28–2.14)	5.53 (1.80–17.03)
Type of diabetes (referent: type 2 diabetes)				†			*	†	†
Type 1 diabetes	4	14	1.56 (0.45–5.50)		62	2.70 (1.10–6.63)			

Diabetes treatment (referent: non-insulin)	43	13	1.84(1.04-3.26)	1.22	(0.59-2.53)	45	1.65(1.16-2.34)	1.46	(0.91-2.36)	1.44	(0.71-2.92)	1.44	(0.76-2.76)
Smoker (referent: nonsmoker)	19	9	0.87(0.41-1.84)	0.78	(0.65-0.95)	33	0.76(0.48-1.20)	0.82	(0.46-1.46)	0.94	(0.80-1.09)	0.94	(0.82-1.08)
Charlson index (mean)	4.02	—	0.80(0.67-0.97)	0.0614		—	0.95(0.87-1.04)	0.94	(0.85-1.04)	0.0318		0.1055	
R ²				0.0856						0.1258		0.2946	
Hosmer-Lemeshow goodness-of-fit P value													

*Not included in the fully adjusted model due to high collinearity with other variables in the model. †Not included in the adjusted model because of small cell sizes. CVD, cardiovascular disease; OR, odds ratio.

though those studies hypothesized that the frequency of recording of diabetes on death certificates would increase, results from our study suggest that the frequency of recording has remained constant despite an increase in the prevalence of diabetes since 1986 (17).

In our study, we were able to ascertain more factors from the decedents and from their medical records before death to investigate predictors of recording. These factors included race/ethnicity, education, income, diabetes treatment, smoking status, and number of comorbidities. Using these in multivariable models, we found that longer duration of diabetes consistently predicted recording diabetes as the underlying cause of death and as a cause of death anywhere on the death certificate for both all-cause mortality and noncardiovascular causes of death. Insulin treatment and having fewer comorbidities were also associated with the reporting of diabetes. This finding suggests that more severe diabetes and having fewer comorbidities are likely to trigger the physician to list diabetes as a cause or contributor to death.

Recording of diabetes on death certificates was most frequent when the cause of death was cardiac, highlighting the well-recognized association between diabetes and cardiovascular disease. Nevertheless, only 39% of diabetic patients dying of cardiovascular disease had diabetes recorded anywhere on their death certificates. Diabetes was significantly less likely to be recorded on death certificates when the cause of death was cancer. National statistics that have used the recording of diabetes on death certificates as an indicator of comorbid diabetes in patients with cancer suggest that individuals with diabetes do not die of cancer as frequently as individuals without diabetes. In a recent publication, Coughlin et al. (35) demonstrated that diabetes may contribute to the development of cancer and that diabetes is an independent predictor of mortality from cancer of the colon, pancreas, female breast, and, among men, of the liver and bladder. Our results, and the realization that death certificates of decedents with cancer are less likely to report diabetes, highlight the need for careful studies of cancer in people with diabetes.

Certainly diabetes is not a cause of or a contributor to death in all decedents with diabetes. Death certificates are intended to record and count events, but people often use them as a tool to measure the burden of disease (3,5,7). Unfortu-

nately, diabetes mortality rates based on death certificates fail to account for all deaths among people with diabetes. Using death certificates to define whether a decedent had diabetes thus underestimates mortality in all people with diabetes and presents a biased perspective on causes of death in people with diabetes. These problems are less of an issue in cohort studies that define diabetes status before death and so cohort studies may be the preferred design for studying diabetes mortality.

At least two limitations were present in our study. First, we limited our multivariable analyses to decedents who consented to medical record review, which excluded 32% of decedents. Those who were excluded from the multivariable models were more likely to be nonwhite, have less education, and not use insulin for diabetes treatment. When multivariate models were run for all decedents but excluded variables obtained from the medical record review, the results were not substantially changed. Second, we studied only managed care enrollees, and our findings may not extend to other populations. Indeed, our finding that <50% of diabetic subjects died of cardiac, cerebrovascular, or other cardiac disease (compared with 65% in other studies [8]) may reflect better access to care and better risk factor control in managed care.

Adding a series of check boxes for common conditions of interest (similar to those on a clinical encounter form) would likely improve recording of diabetes as a coexistent condition on death certificates (9). Without such an intervention, death certificates alone are not a good source of data to estimate the strength of associations between the presence of diabetes and specific causes of death. Similarly, death certificates probably do not provide a valid basis for determining the effect of diabetes on life expectancy, since the average age at death may differ by causes for which the completeness of recording of diabetes may differ. In conclusion, death certificates underestimate the prevalence of diabetes among decedents and present a biased picture of the causes of death among people with diabetes.

APPENDIX

The TRIAD Study Group

Pacific Health Research Institute. Principal Investigator: J. David Curb, MD, MPH; Coinvestigators: Beth Waitzfelder, PhD; Adams Dudley, MD, MBA; Chien-

Wen Tseng, MD, MPH; Project Coordinator: Suzanne Firrell; Analyst: Qimei He, PhD.

Indiana University Translational Research Center. Principal Investigator: David G. Marrero, PhD; Coinvestigators: Ronald T. Ackermann, MD, MPH; Matthew J. Bair, MD, MS; Ed Brizendine, MS; Aaron E. Carroll, MD; Ann Chou, PhD; Gilbert C. Liu, MD, MS; Paris Roach, MD; Changyr Shen, PhD; Morris Weinberger, PhD; Madelyn L. Wheeler, MS, RD, CD, FADA, CDE; Project Coordinator: Susanna R. Williams, MSPH.

Kaiser Foundation Research Institute. Principal Investigator and study chairman: Joseph V. Selby, MD, MPH; Co-Principal Investigator: Andrew J. Karter, PhD; Coinvestigator: Assiamira Ferrara, MD, PhD; Project Director: Bix E. Swain, MS.

University of California Los Angeles School of Medicine. Principal Investigator: Carol M. Mangione, MD; Co-Principal Investigator: Arleen Brown, MD, PhD; Coinvestigators: Kenrik Duru, MD; Susan Ettner, PhD; Shaista Malik, MD, MPH; Martin F. Shapiro, MD; Project Director: Rebecca Brusuelas; Analysts: Neil Steers, PhD; Peter Guierrez; Administrative Assistant: Carole Nagy.

University of Medicine and Dentistry of New Jersey. Principal Investigator: Norman Lasser, MD, PhD; Coinvestigators: Dorothy A. Caputo, MA, APRN, BC-ADM, CDE; Jesse C. Crosson, PhD; Stephen Crystal, PhD; David S. Kountz, MD; Shou-En Lu, PhD; Patricia Prata, MPH, CHES; Monika Safford, MD; Stephen H. Schneider, MD; Weichung (Joe) Shih, PhD; Leslie-Faith Morrill Taub, DNSc, AND-C, GNP-BC; Program Coordinator: Sonja Ross, MS, MHS; Research Assistant: Ahmus H. Mahmoud, MB, ChB; TRIAD-wide Program Specialist: Gabrielle J. Davis, BS, CHES; TRIAD-wide Program Assistant: Lucyna Lis, MA.

The University of Michigan Health System. Principal Investigator: William H. Herman, MD, MPH; Coinvestigators: Michele Heisler, MD; Susan Johnson, MD; Catherine Kim, MD, MPH; Project Director: Jennifer Goewey, MSA; Research Associates: Ray Burke, MA; Laura McEwen, PhD; Bahman Tabaei, MPH; Honghong Zhou, MS; Administrative Assistant: Barbara Pearlman.

National Institute of Diabetes & Digestive & Kidney Diseases. Coinvestigator: Sanford A. Garfield, PhD.

Division of Diabetes Translation, Centers for Disease Control and Preven-

tion. Lead Consultant: K.M. Venkat Narayan, MD, MSc, MBA, MRCP; Collaborating Investigators: Gloria Beckles, MD, MSc; Patrick Boyle, PhD; Michael Engle-gau, MD; Tiffany Gary, PhD; Linda Geiss, MS; Bob Gerzoff, MS; Edward W. Gregg, PhD; Henry Kahn, MD, FACP; Jinan Saaddine, MD; Mark Stevens, MSPH, MA; Theodore Thompson, MS; Ed Tierney, MPH; Roldofo Valdez, PhD; David F. Williamson, PhD; Ping Zhang, PhD; Program Administrator: Bernice Moore, MBA; Management and Program Analyst: Shay Clayton.

TRIAD-wide Administrative Data Co-ordinator. Barbara K. Smith, MHSA.

TRIAD, Department of Veterans Affairs. Principal Investigator: Eve Kerr, MD, MPH; Co-Principal Investigator: Rodney Hayward, MD; Coinvestigators: Sarah Krein, PhD; John Peitte, PhD; Project Managers: Mary Hogan, PhD, RN; Emily Lipp, MPH; Fatima Makki, MPH, MSW; Data Manager: Jennifer Davis, MPH.

Acknowledgments— This study was jointly funded by Program Announcement number 04005 from the Centers for Disease Control and Prevention (Division of Diabetes Translation) and the National Institute of Diabetes and Digestive and Kidney Diseases. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of The Centers for Disease Control and Prevention and the National Institute of Diabetes and Digestive and Kidney Diseases.

Significant contributions to this study were made by members of the TRIAD study group (see APPENDIX). The authors acknowledge the participation of our health plan partners.

References

1. Greenberg JA: Biases in the mortality risk versus body mass index relationship in the NHANES-1 Epidemiologic Follow-Up Study. *Int J Obes Relat Metab Disord* 25:1071–1078, 2001
2. Taffet GE, Teasdale TA, Bleyer AJ, Kutka NJ, Luchi RJ: Survival of elderly men with congestive heart failure. *Age Ageing* 21:49–55, 1992
3. Anderson RN: Deaths: leading causes for 1999. *Natl Vital Stat Rep* 49:1–87, 2001
4. U.S. Department of Health and Human Services: *Health, United States, 2004: With Chartbook on Trends in the Health of Americans*. Hyattsville, MD, National Center for Health Statistics, 2004
5. Arias E, Anderson RN, Kung HC, Murphy SL, Kochanek KD: Deaths: final data for 2001. *Natl Vital Stat Rep* 52:1–115, 2003
6. Kochanek KD, Smith BL: Deaths: prelim-

inary data for 2002. *Natl Vital Stat Rep* 52:1–47, 2004

7. Gross CP, Anderson GF, Powe NR: The relation between funding by the National Institutes of Health and the burden of disease. *N Engl J Med* 340:1881–1887, 1999
8. Geiss LS, Herman WH, Smith PJ: Mortality in non-insulin-dependent diabetes. In *Diabetes in America*, 2nd ed. Harris MI, Cowie CC, Stern MP, Boyko EJ, Reiber GE, Bennett PH, Eds. Bethesda, MD, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, 1995, p. 233–258 (NIH publ. no. 95–1468)
9. Tierney EF, Geiss LS, Engulgau MM, Thompson TJ, Schaubert D, Shireley LA, Vukelic PJ, McDonough SL: Population-based estimates of mortality associated with diabetes: use of a death certificate check box in North Dakota. *Am J Public Health* 91:84–92, 2001
10. National Center for Health Statistics, Division of Vital Statistics: U.S. Standard Certificate of Death [article online], 1989. Available from <http://www.cdc.gov/nchs/data/dvs/std-dcrt.pdf>. Accessed 22 October 2004
11. Centers for Disease Control and Prevention: From the Centers for Disease Control and Prevention: sensitivity of death certificate data for monitoring diabetes mortality: Diabetic Eye Disease Follow-Up Study, 1985–1990. *JAMA* 266:2812, 1991
12. Fuller JH, Elford J, Goldblatt P, Adelstein AM: Diabetes mortality: new light on an underestimated public health problem. *Diabetologia* 24:336–341, 1983
13. West K: *Epidemiology of Diabetes and Its Vascular Lesions*. New York, Elsevier, 1978
14. Fuller JH: Mortality trends and causes of death in diabetic patients. *Diabetes Metab* 19:96–99, 1993
15. Fuller JH: Causes of death in diabetes mellitus. *Horm Metab Res Suppl* 15:3–9, 1985
16. Bild DE, Stevenson JM: Frequency of recording of diabetes on U.S. death certificates: analysis of the 1986 National Mortality Followback Survey. *J Clin Epidemiol* 45:275–281, 1992
17. Will JC, Vinicor F, Stevenson J: Recording of diabetes on death certificates: has it improved? *J Clin Epidemiol* 54:239–244, 2001
18. Penman A: Excess mortality due to diabetes in Mississippi and the estimated extent of underreporting on death certificates. *J Miss State Med Assoc* 44:319–325, 2003
19. Ochi JW, Melton LJ 3rd, Palumbo PJ, Chu CP: A population-based study of diabetes mortality. *Diabetes Care* 8:224–229, 1985
20. Palumbo PJ, Elveback LR, Chu CP, Connolly DC, Kurland LT: Diabetes mellitus: incidence, prevalence, survivorship, and causes of death in Rochester, Minnesota,

- 1945–1970. *Diabetes* 25:566–573, 1976
21. Anderson EM, Lee JA, Pecoraro RE, Kopsell TD, Hallstrom AP, Siscovick DC: Underreporting of diabetes on death certificates, King County, Washington. *Am J Public Health* 83:1021–1024, 1993
 22. Kleinman JC, Donahue RP, Harris MI, Finucane FF, Madans JH, Brock DB: Mortality among diabetics in a national sample. *Am J Epidemiol* 128:389–401, 1988
 23. Pandey DK, Labarthe DR, Goff DC, Chan W, Nichaman MZ: Community-wide coronary heart disease mortality in Mexican Americans equals or exceeds that in non-Hispanic whites: the Corpus Christi Heart Project. *Am J Med* 110:81–87, 2001
 24. TRIAD: The Translating Research Into Action for Diabetes (TRIAD) study: a multicenter study of diabetes in managed care. *Diabetes Care* 25:386–389, 2002
 25. Centers for Disease Control and Prevention, National Center for Health Statistics: National Death Index (NDI), NDI Plus Searches. [article online], 2003. Available from http://www.cdc.gov/nchs/r&d/ndi/what_is_ndi.htm. Accessed 15 April 2003
 26. Cowper DC, Kubal JD, Maynard C, Hynes DM: A primer and comparative review of major United States mortality databases. *Ann Epidemiol* 12:462–468, 2002
 27. Williams BC, Demitrack LB, Fries BE: The accuracy of the National Death Index when personal identifiers other than Social Security number are used. *Am J Public Health* 82:1145–1147, 1992
 28. Luepker RV, Apple FS, Christenson RH, Crow RS, Fortmann SP, Goff D, Goldberg RJ, Hand MM, Jaffe AS, Julian DG, Levy D, Manolio T, Mendis S, Mensah G, Pajak A, Prineas RJ, Reddy KS, Roger VL, Rosamond WD, Shahar E, Sharrett AR, Sorlie P, Tunstall-Pedoe H: Case definitions for acute coronary heart disease in epidemiology and clinical research studies: a statement from the AHA Council on Epidemiology and Prevention, AHA Statistics Committee, World Heart Federation Council on Epidemiology and Prevention, the European Society of Cardiology Working Group on Epidemiology and Prevention, Centers for Disease Control and Prevention, and the National Heart, Lung, and Blood Institute. *Circulation* 108:2543–2549, 2003
 29. Klein R, Klein BE: Relation of glycemic control to diabetic complications and health outcomes. *Diabetes Care* 21 (Suppl. 3):C39–C43, 1998
 30. Charlson ME, Pompei P, Ales KL, MacKenzie CR: A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 40:373–383, 1987
 31. de Groot V, Beckerman H, Lankhorst GJ, Bouter LM: How to measure comorbidity: a critical review of available methods. *J Clin Epidemiol* 56:221–229, 2003
 32. Robbins JM, Vaccarino V, Zhang H, Kasl SV: Socioeconomic status and type 2 diabetes in African American and non-Hispanic white women and men: evidence from the Third National Health and Nutrition Examination Survey. *Am J Public Health* 91:76–83, 2001
 33. Will JC, Casper M: The contribution of diabetes to early deaths from ischemic heart disease: US sex and racial comparisons. *Am J Public Health* 86:576–579, 1996
 34. Littell RC, Milliken GA, Stroup WW, Wolfinger RD: *SAS System for Mixed Models*. Cary, NC, SAS Inst., 1996
 35. Coughlin SS, Calle EE, Teras LR, Petrelli J, Thun MJ: Diabetes mellitus as a predictor of cancer mortality in a large cohort of US adults. *Am J Epidemiol* 159:1160–1167, 2004