

Genetics of Gestational Diabetes Mellitus and Type 2 Diabetes

RICHARD M. WATANABE, PHD^{1,2}
 MARY HELEN BLACK, MS¹
 ANNY H. XIANG, PHD¹

HOUMAN ALLAYEE, PHD^{1,3}
 JEAN M. LAWRENCE, SCD⁴
 THOMAS A. BUCHANAN, MD⁵

The successful application of positional cloning, the process of disease susceptibility gene identification using gene mapping techniques, to identify the gene underlying cystic fibrosis (1) opened the era of gene hunting. The subsequent identification of susceptibility genes underlying numerous monogenic disorders led to the possibility that susceptibility genes for complex diseases could also be identified using the positional cloning approach. This, coupled with advances in both genotyping technology and the identification of large numbers of microsatellite markers across the genome, led to numerous genome-wide scans to identify susceptibility genes for various forms of diabetes. However, despite the genetic screening of genes known to be involved in the biology of diabetes (candidate genes) and positional cloning approaches, to date, few susceptibility genes for diabetes have been identified. Even less is known about the genetic basis for gestational diabetes mellitus (GDM). The inability to readily identify susceptibility genes for diabetes can be attributed to a variety of issues, including insufficient statistical power, etiologic heterogeneity, and the confounding effect of interactions with environmental factors. These same problems will likely apply to GDM, which appears to represent early stages of many forms of diabetes

outside of pregnancy. Here, we briefly review the current state of knowledge regarding the genetics of diabetes and discuss specific issues regarding the genetics of GDM.

GENETICS OF DIABETES

Type 1 diabetes and rare/monogenic forms of diabetes

The HLA region on chromosome 6 was identified very early on as a major susceptibility gene for type 1 diabetes (2–4), with haplotypes within the HLA region accounting for as much as 50% of cases of type 1 diabetes in Caucasians (4). While the contribution of HLA to genetic susceptibility to type 1 diabetes was readily identified, numerous genome scans have also identified at least 16 additional loci across the genome that may harbor susceptibility genes for type 1 diabetes (5). However, identification of the specific genes underlying these linkage regions has proven to be difficult, partly because of the presence of the large genetic contribution of HLA alleles.

Greater success has been found in identifying susceptibility genes for monogenic (6–10), rare (11,12), or syndromic (13) forms of diabetes. Rare variation in the insulin gene results in an autosomal dominant form of diabetes (11), and the 3,243 A-G mutation in the mitochondrial

tRNA (Leu-UUR) gene results in maternally inherited diabetes and deafness syndrome (12,14). Susceptibility genes identified in syndromic forms of diabetes include the Wolfram Syndrome gene (*WSF1*) and mutations in translation initiation factor 2- α kinase-3 gene (*EIF2AK3*), resulting in Wolcott-Rallison syndrome. Genes underlying susceptibility for the six different forms of maturity-onset diabetes of the young (MODY) have been identified (6–10,15). MODY is a form of diabetes characterized by an early age of onset, autosomal dominant inheritance, and β -cell dysfunction in the absence of insulin resistance (15). All six MODY genes are associated with the pancreatic β -cell, either altering transcriptional regulation or possibly altering β -cell mass or turnover (15).

The physiological consequence of genetic variation in MODY genes was characterized in a series of elegant studies by Polonsky and colleagues (16–18). Insulin secretory response to intravenous glucose infusion was assessed in nondiabetic patients and patients with known MODY variants in glucokinase (*GCK*, MODY2), hepatocyte nuclear factor-1 α (*HNF1A*, MODY3), and hepatocyte nuclear factor-4 α (*HNF4A*, MODY1). These studies demonstrated that the insulin secretory dose-response in patients with MODY susceptibility variants in *GCK* was less responsive and generally right-shifted compared with nondiabetic patients. In contrast, patients with MODY susceptibility variants in *HNF1A* or *HNF4A* appeared to have a normal secretory dose-response at lower glucose concentrations, but had maximal secretory responses that were substantially blunted compared with nondiabetic patients. Interestingly, the dose-response curves for *HNF1A* and *HNF4A* appeared to have similar characteristics, suggesting that these variants may have similar effects on the β -cell and overall insulin secretory response. However, subsequent studies in which patients were exposed to prolonged hyperglycemia, brought about by a 42-h glucose infusion, revealed that although blunted, increased insulin secretion observed after prolonged hyperglycemia, the so-called “priming effect” (19), could be observed in patients with *HNF1A* variants, whereas the priming

From the ¹Department of Preventive Medicine, Division of Biostatistics, Keck School of Medicine of USC, Los Angeles, California; the ²Department of Physiology and Biophysics, Keck School of Medicine of USC, Los Angeles, California; the ³Institute for Genetic Medicine, Keck School of Medicine of USC, Los Angeles, California; ⁴Research and Evaluation, Kaiser Permanente Southern California, Pasadena, California; and the ⁵Department of Medicine, Division of Diabetes and Endocrinology, Keck School of Medicine of USC, Los Angeles, California.

Address correspondence and reprint requests to Richard M. Watanabe, PhD, Department of Preventive Medicine, Keck School of Medicine of USC, 1540 Alcazar St., CHP-220, Los Angeles, CA 90089-9011. E-mail: rwatanab@usc.edu.

Received for publication 28 March 2006 and accepted in revised form 16 May 2006.

This article is based on a presentation at a symposium. The symposium and the publication of this article were made possible by an unrestricted educational grant from LifeScan, Inc., a Johnson & Johnson company.

Abbreviations: GDM, gestational diabetes mellitus; FUSION, Finland-U.S. Investigation of Non-Insulin-Dependent Diabetes Mellitus; MODY, maturity-onset diabetes of the young; SNP, single-nucleotide polymorphism.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

DOI: 10.2337/dc07-s205

© 2007 by the American Diabetes Association.

effect was absent in patient with *HNF4A* variants.

Type 2 diabetes

One of the first type 2 diabetes susceptibility genes was identified not by linkage analysis, but by association (20). Deeb et al. (20) observed association between the Pro12Ala polymorphism in peroxisome proliferator-activated receptor- γ and BMI and insulin sensitivity in a sample of Finnish subjects. Individuals with at least one copy of Ala had a lower BMI and a higher insulin sensitivity compared with individuals homozygous for Pro. In addition, they observed a significantly increased risk of type 2 diabetes (odds ratio = 4.25) for individuals homozygous for Pro in a sample of Japanese Americans. The lower frequency Ala allele appeared to play a "protective" role against type 2 diabetes, since the Ala allele was also associated with a lower transactivation of response elements and therefore higher insulin sensitivity (20).

Peroxisome proliferator-activated receptor- γ is a nuclear receptor involved in adipocyte differentiation (21) and a target for the thiazolidinedione class of insulin-sensitizing drugs (22), making it an attractive candidate gene for type 2 diabetes. The initial report by Deeb et al. (20) led many groups to assess the Pro12Ala polymorphism, resulting in a variety of positive and negative associations. This led to some question as to whether Pro12Ala was a true diabetes susceptibility variant. However, this was resolved by a meta-analysis performed by Altshuler et al. (23). In their analysis of over 3,000 subjects, Altshuler et al. observed a modest but significant genotype relative risk (RR = 1.25) for type 2 diabetes associated with the Pro allele (23). When these results were combined with previously published reports in a meta-analysis, a significant protective effect for the Ala allele with a risk ratio of 0.79 was observed (23).

Over 25 genome-wide linkage scans for type 2 diabetes have been completed to date (24). Despite the large number of studies, very few regions of the genome showed common evidence for linkage across studies. Regions showing the greatest replication among studies include chromosomes 1q, 12q, and 20q. Initially, seven studies showed overlapping evidence for linkage to chromosome 1q (24). This led to the formation of the International Type 2 Diabetes 1q Consortium, in which investigators from multiple studies

are pooling samples, resources, and an effort to fine-map susceptibility genes underlying this region.

One of the first genome-wide scans to be published came from Hanis et al. (25), who reported linkage results from Mexican Americans from Starr County, TX. The strongest linkage signal was observed on chromosome 2q, with the locus being dubbed *NIDDM1*. Subsequent analyses and fine-mapping efforts of this region led to the first type 2 diabetes susceptibility gene to be identified via positional cloning, calpain-10 (*CAPN10*), a ubiquitously expressed member of the calpain-like cysteine protease family (26). Three single-nucleotide polymorphisms (SNPs), SNP-43, -19, and -63, formed a high-risk haplotype within the Starr County Mexican-American sample. As with peroxisome proliferator-activated receptor- γ , subsequent studies have reported a mixture of both positive and negative associations in other populations leading to extensive discussion of whether *CAPN10* could be classified as a type 2 diabetes susceptibility gene. Evans et al. (27) were the first to demonstrate that another SNP with *CAPN10*, SNP-44, located 11 bp from SNP-43, was associated with type 2 diabetes in Caucasian subjects from the U.K. A subsequent meta-analysis by Weedon et al. (28) examined whether SNP-44 was associated with type 2 diabetes. Their analysis included Asian, Mexican-American, and Caucasian samples and concluded that *CAPN10* was associated with an overall modest increase in risk for type 2 diabetes (odds ratio = 1.17). However, examination of the combined Mexican-American samples alone suggests *CAPN10* may be associated with a higher risk for type 2 diabetes in this specific ethnic group (odds ratio = 2.13), compared with the Asian (odds ratio = 1.09) and Caucasian (odds ratio = 1.17) samples.

Most recently, the DeCode group attempting to positionally clone genes for a variety of common diseases reported association between type 2 diabetes and variation in the gene for transcription factor 7-like 2 (*TCF7L2*) (29). The initial genome-wide linkage analysis performed in large families from the population of Iceland revealed evidence for linkage on chromosome 10 (30). Subsequent fine-mapping of the chromosome 10 interval of interest revealed a microsatellite marker, DG10S478, to be associated with type 2 diabetes. This association was replicated in both Danish and U.S. Caucasian

samples. Individuals heterozygous for the risk allele had a relative risk for type 2 diabetes of 1.45, whereas individuals homozygous for the risk allele had a relative risk of 2.41. The mechanism by which *TCF7L2* confers risk for type 2 diabetes is unknown, but it has been hypothesized that it may regulate proglucagon gene expression in the enteroendocrine cells (29).

Chromosome 20

Linkage to chromosome 20q was also observed for multiple studies and was the impetus for the formation of the International Type 2 Diabetes Linkage Consortium. The 20q region was independently fine-mapped by several groups, resulting in the identification of a novel glucose transporter (*SLC2A10*) (31) that does not appear to confer susceptibility to type 2 diabetes (32,33), along with two type 2 diabetes susceptibility genes: protein-tyrosine phosphatase, nonreceptor-type 1 (*PTPN1*), and hepatocyte nuclear factor-4 α (*HNF4A*). *PTPN1* was first shown to be associated with type 2 diabetes in patients with diabetes and end-stage renal disease and participants of the Diabetes Heart Study (34). Subsequently, variation in this gene was also shown to be associated with insulin resistance and fasting glucose in Hispanic Americans from the Insulin Resistance Atherosclerosis Study Family Study (35). However, in one of the largest association studies of type 2 diabetes with over 3,000 cases and 3,000 control subjects from mainly Northern European Caucasian samples, no evidence for association with type 2 diabetes was found for single variants within *PTPN1* or with observed haplotypes (36).

The fall and rise of *HNF4A*. The second susceptibility gene to be identified in the 20q region is *HNF4A*, already described above as a susceptibility gene for MODY. *HNF4A* was initially rejected by one study as a type 2 diabetes susceptibility gene. The Finland-U.S. Investigation of Non-Insulin-Dependent Diabetes Mellitus (FUSION) study (37), an effort to positionally clone type 2 diabetes susceptibility genes in the Finnish population, was one of several groups (38–41) showing evidence for linkage in the 20q region (42). Because *HNF4A* fell within the linkage region and given that coding variants were known to confer susceptibility to MODY1, Ghosh et al. (42) sequenced this gene and tested variants for association for type 2 diabetes. None of the variants they identified were associated with type

2 diabetes, leading them to conclude that variation in *HNF4A* was “. . . unlikely to account for the linkage results on chromosome 20q.”

However, ~5 years later, the FUSION group along with Love-Gregory et al., jointly reported association between variation in the P2 promoter region of *HNF4A* and type 2 diabetes (43,44). Love-Gregory et al. (44), who were studying the Ashkenazi Jewish population, and the FUSION study, observed the associations independently while fine-mapping the 20q region. Both groups subsequently collaborated to ensure that common variants were genotyped in both samples, resulting in common SNPs showing association with type 2 diabetes in the P2 promoter region. In both studies, the associated SNPs accounted for a significant fraction of the evidence for type 2 diabetes in the 20q region (43,44). Type 2 diabetes-related quantitative trait data were not available in the Ashkenazi Jewish sample, but FUSION observed association between variation in the P2 promoter region and acute insulin response and disposition index in offspring of their affected patients (43), consistent with the presumed effect of *HNF4A* within the β -cell. The association between variation in the P2 promoter region of *HNF4A* and type 2 diabetes has been replicated in other populations (45–47).

GENETICS OF GDM

Familial clustering

So what about GDM? Is there a genetic basis for GDM? Surprisingly, there has been relatively little research in the area of GDM genetics per se. An essential first step in genetics research has been the determination of evidence for a genetic basis for the disease. This can come in the form of twin concordance studies or estimates of familial risk or heritability. However, performing such studies in a prospective fashion is fraught with numerous difficulties, primarily the need to identify women who will become pregnant. Studies are also difficult to perform in a retrospective fashion. The clinical definition for GDM has evolved over the years and differs slightly among countries (48–51). Furthermore, there has not been consistent screening for GDM, leading to possible bias in ascertainment, e.g., missed cases. Finally, there are difficulties in ascertaining families with multiple cases of GDM, which is partly related to the relatively low prevalence of GDM. There has been,

to our knowledge, only one unpublished attempt to estimate familiarity of GDM. In 1999, Williams and colleagues used the statewide medical record system in the state of Washington to identify and link sisters diagnosed with GDM using *International Classification of Diseases, Ninth Revision* (ICD-9), coding. Based on their initial screening, they estimated that the sibling risk ratio for GDM was 1.75 (M. Williams, personal communication), suggesting some evidence for a genetic basis for GDM. This risk, likely an underestimate, is significantly lower than the estimated sibling risk ratio for type 2 diabetes, which ranges from 2 to 4 (52).

The question of a genetic basis for GDM, however, is also closely tied to the debate of whether GDM is a unique disease state or whether pregnancy with its associated metabolic derangements simply provides us with a crystal ball with which to identify women who are susceptible to hyperglycemia and subsequent development of diabetes. If genetic variants associated with type 1 or type 2 diabetes are also associated with GDM, from a purely genetic perspective, it would be difficult to argue a unique genetic predisposition for GDM. This does not take into account the possibility of unique environmental exposures related to pregnancy that may interact with genetic variants to alter disease risk. Also, this does not negate the importance of using genetic information to improve treatment for GDM and minimize the deleterious effect of hyperglycemia on fetal outcomes.

There are studies that have examined the familial clustering of GDM and type 1 and type 2 diabetes. Examples include the studies of Dorner et al. (53), who showed increased familial aggregation of diabetes on the maternal side of offspring with type 1 diabetes whose mothers had GDM. Similarly, there is evidence for clustering of type 2 diabetes and impaired glucose tolerance in families with a GDM (54) and evidence for higher prevalence of type 2 diabetes in mothers of women with GDM (55). Thus, there is evidence of some link between both autoimmune and nonautoimmune forms of diabetes and GDM.

Candidate genes

Candidate genes related to both autoimmune and nonautoimmune forms of GDM have been assessed in a variety of cohorts. Freinkel et al. (56) examined whether HLA antigens were associated with GDM. They observed that HLA DR3 and DR4 antigens were uncommon over-

all, but nonetheless in higher frequency in women with GDM than in women with normal pregnancies. Similarly, Ober et al. (57) reported association between variation in the insulin receptor (*INSR*) in Caucasian and African-American women with GDM. They also noted that variation in *INSR* appeared to interact with both BMI and history of diabetes in mothers with GDM. Among Caucasian women, *INSR* variants also appeared to interact with variation in insulin-like growth factor-2 (*IGF2*). No associations between *INSR* and *IGF2* were observed in Hispanic women with GDM (57).

There have also been several reports of association between variation in *GCK* and GDM (58–63). Stoffel et al. (58) estimated that the frequency of *GCK* variants in GDM was ~5%, by extrapolating observations from 40 women with GDM. The relatively low frequency of *GCK* variants among GDM subjects was also confirmed by others (60,61). Despite the low frequency, the important contribution of *GCK* variation to risk for GDM can also be observed in families of *MODY2* patients. Saker et al. (60) noted that a large proportion of female members of *MODY2* families present with GDM. They further speculated that because variation in *GCK* typically results in subclinical hyperglycemia, the frequency of *GCK* variants may be higher and only detectable upon pregnancy (60). This possibility was confirmed by Ellard et al. (62) who used highly selective clinical criteria to select patients with GDM and tested whether they carried *GCK* variants (62). Their data suggest that the prevalence of *GCK* variants may be as high as 80% in a small subset of women with GDM selected by highly specific clinical criteria.

***HNF4A* in GDM.** Studies by a variety of investigators have identified a β -cell defect as being one of the primary characteristics of GDM (64–67). Given that β -cell function is a highly heritable trait (68–71), we became interested in trying to identify genes underlying β -cell dysfunction observed in GDM. This resulted in the BetaGene study, in which we are recruiting Mexican-American families of a proband with previous GDM (71,72). The reported associations between variation in the P2 promoter region of *HNF4A* and type 2 diabetes (43,44,46,73) led us to examine whether these variants might be associated with diabetes-related quantitative traits. Muller et al. (73) examined these variants for association with type 2 diabetes-related quantitative traits in

Pima Indians and only observed modest association with insulin resistance, as assessed by the euglycemic glucose clamp. In the previous reports by Silander et al. (43) and Love-Gregory et al. (44), the frequency of the minor allele for the associated SNP (rs2144908) was between 16 and 27%, depending on the sample. In contrast, the same allele had a frequency of 49% in the Mexican-American sample from the BetaGene study (72). When we examined this SNP for association with type 2 diabetes-related phenotypes, we observed a significant association with disposition index ($P = 0.035$) under an additive genetic model. Disposition index is a measure of β -cell compensation (67,74), and this association is consistent with the known biologic function of *HNF4A* in the pancreatic β -cells (75). Thus, variation in *HNF4A* may contribute to the β -cell dysfunction observed in GDM.

THE FUTURE — The field of genetics has come a long way since the days of Gregor Mendel and his pea experiments. During a period when positional cloning of complex disease genes by linkage analysis using microsatellite markers was reaching its zenith, a landmark article by Risch and Merikangas (76) appeared in the literature. This article provided a theoretical argument that large-scale association analysis was statistically more powerful than linkage analysis and therefore the preferable approach to identify genes underlying complex diseases. In that article, the authors are quoted as saying, “. . . imagine the time when all human genes (say 100,000 in total) have been found and that simple, diallelic polymorphisms in these genes have been identified.” Although far from reality at the time of publication, rapid advances in genotyping technology (77), drastic reductions in genotyping costs, the sequencing of the human genome (78,79), and the recent completion of the HapMap project (80) have now made whole genome association analysis a reality. These advances have now made it possible to select 250,000 to 500,000 SNPs across the human genome and rapidly genotype them in samples of 2,000–3,000 subjects in an affordable manner.

What does this new era of whole genome association mean for the genetics of GDM? It may now be possible to identify large case-control samples and perform whole genome association to identify regions of the genome harboring suscepti-

bility genes for GDM. The unanswered question of whether GDM per se has a genetic basis may also be indirectly addressed using whole genome association by incorporating carefully selected samples of type 2 diabetes cases as a secondary contrast group into the study design. SNPs showing association with GDM, but not type 2 diabetes, may represent susceptibility genes unique to GDM. Finally, the identification of genetic variants underlying disease alone will not have a major impact on clinical care for patients with GDM. Like the MODY studies of Polonsky and his colleagues (16–18), additional molecular and biochemical studies, and most importantly, clinical studies must be performed to understand the physiological and clinical consequences of genetic variation. Finally, we will need to assess how to maximize genetic information and how to best incorporate it into the clinical care setting.

Acknowledgments — R.M.W. was supported in part by a grant from the American Diabetes Association (05-RA-140). T.A.B. was supported in part by a Distinguished Clinical Scientist Award from the American Diabetes Association. The FUSION study was supported by intramural funds from the National Human Genome Research Institute (project no. OH95-C-N030) and by National Institutes of Health (NIH) Grants DK62370 and HG00376. The BetaGene Study was supported by NIH Grant DK61628 (to T.A.B.), with support from the University of Southern California General Clinical Research Center (M01-RR-00043).

References

- Riordan JR, Rommens JM, Kerem B-S, Alon N, Rozmahel R, Grzelczak Z, Zielenski J, Lok S, Plavsic N, Chou JL, Drummler ML, Iannuzzi MC, Collins FS, Tsui L-C: Identification of the cystic fibrosis gene: cloning and characterization of the complementary DNA. *Science* 245:1066–1073, 1989
- Barbosa J, Chern MM, Noreen H, Anderson VE: Analysis of linkage between the major histocompatibility system and juvenile, insulin-dependent diabetes in multiplex families: reanalysis of data. *J Clin Invest* 62:492–495, 1978
- Trucco M, Dorman JS: Immunogenetics of insulin-dependent diabetes mellitus in humans. *Crit Rev Immunol* 9:201–245, 1989
- Noble JA, Valdes AM, Cook M, Klitz W, Thomson G, Erlich HA: The role of HLA class II genes in insulin-dependent diabetes mellitus: molecular analysis of 180 Caucasian, multiplex families. *Am J Hum*

- Genet* 59:1134–1148, 1996
- Pugliese A: Genetics of type 1 diabetes. *Endocrinol Metab Clin N Am* 33:1–16, 2004
- Froguel P, Zouali H, Vionnet N, Velho G, Vaxillaire M, Sun F, Lesage S, Stoffel M, Takeda J, Passa P, Permutt MA, Beckmann JS, Bell GI, Cohen D: Familial hyperglycemia due to mutations in glucokinase. *N Engl J Med* 328:697–702, 1993
- Yamagata K, Furuta H, Oda N, Kaisaki PJ, Menzel S, Cox NJ, Fajans SS, Signorini S, Steffel M, Bell GI: Mutations in the hepatocyte nuclear factor-4 α gene in maturity-onset diabetes of the young (MODY1). *Nature* 384:458–460, 1996
- Yamagata K, Oda N, Kaisaki PJ, Menzel S, Furuta H, Vaxillaire M, Southam L, Cox RD, Lathrop GM, Boriraj W, Chen X, Cox NJ, Oda Y, Yano H, LeBeau MM, Yamada S, Nishigori H, Takeda J, Fajans SS, Hattersley AT, Iwasaki N, Hansen T, Pedersen O, Polonsky KS, Turner RC, Velho G, Chevre JC, Froguel P, Bell GI: Mutations in the hepatocyte nuclear factor-1 α gene in maturity-onset diabetes of the young (MODY3). *Nature* 384:455–458, 1996
- Stoffers DA, Ferrer J, Clarke WL, Habener JF: Early-onset type-II diabetes mellitus (MODY4) linked to IPF1. *Nat Genet* 17:138–139, 1997
- Horikawa Y, Iwasaki N, Hara M, Furuta H, Hinokio Y, Cockburn BN, Lindner T, Yamagata K, Ogata M, Tomonaga O, Kuroki H, Kasahara T, Iwamoto Y, Bell GI: Mutation in hepatocyte nuclear factor-1 β gene (TCF2) associated with MODY. *Nat Genet* 17:384–385, 1997
- Tager HS: Abnormal products of the human insulin gene. *Diabetes* 33:693–699, 1984
- Kadowaki T, Kadowaki H, Mori Y: A subtype of diabetes mellitus associated with a mutation of mitochondrial DNA. *N Engl J Med* 330:962–966, 1994
- Shih DQ, Stoffel M: Molecular etiologies of MODY and other early-onset forms of diabetes. *Curr Diab Rep* 2:125–134, 2002
- Aggarwal P, Gill-Randall R, Wheatley T, Buchalter MB, Metcalfe J, Alcolado JC: Identification of mtDNA mutation in a pedigree with gestational diabetes, deafness, Wolff-Parkinson-White syndrome and placenta accreta. *Hum Hered* 51:114–116, 2001
- Fajans SS, Bell GI, Polonsky KS: Molecular mechanisms and clinical pathophysiology of maturity onset diabetes of the young. *N Engl J Med* 345:971–980, 2001
- Byrne MM, Sturis J, Clement K, Vionnet N, Pueyo ME, Stoffel M, Takeda J, Passa P, Cohen D, Bell GI, Velho G, Froguel P, Polonsky KS: Insulin secretory abnormalities in subjects with hyperglycemia due to glucokinase mutations. *J Clin Invest* 93:1120–1130, 1994

17. Byrne MM, Sturis J, Menzel S, Yamagata K, Fajans SS, Dronsfield MJ, Bain SC, Hattersley AT, Velho G, Froguel P, Bell GI, Polonsky KS: Altered insulin secretory responses to glucose in diabetic and nondiabetic subjects with mutations in the diabetes susceptibility gene MODY3 on chromosome 12. *Diabetes* 45:1503–1510, 1996
18. Byrne MM, Sturis J, Fajans SS, Ortiz FJ, Stoltz A, Steffel M, Smith MJ, Bell GI, Halter JB, Polonsky KS: Altered insulin secretory responses to glucose in subjects with a mutation in the MODY1 gene on chromosome 20. *Diabetes* 44:699–704, 1997
19. Efendic S, Lins PE, Cerasi E: Potentiation and inhibition of insulin release in man following priming with glucose and with arginine: effect of somatostatin. *Acta Endocrinol (Copenh)* 90:259–271, 1979
20. Deeb SS, Fajas L, Nemoto M, Pihlajamäki J, Laakso M, Fujimoto W, Auwerx J: A Pro12Ala substitution in PPAR γ 2 associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. *Nat Genet* 20:284–287, 1998
21. Tontonoz P, Graves RA, Budavari AI, Erdjument-Bromage H, Jui M, Hu E, Tempst P, Spiegelman BM: Adipocyte-specific transcription factor ARF6 is a heterodimeric complex of two nuclear hormone receptors, PPAR γ and RXR α . *Nucleic Acid Res* 22:5628–5634, 1994
22. Berger J, Bailey P, Biswas C, Cullinan CA, Doebber TW, Hayes NS, Saperstein R, Smith RG, Leibowitz MD: Thiazolidinediones produce a conformational change in peroxisomal proliferator-activated receptor- γ : binding and activation correlate with antidiabetic actions in db/db mice. *Endocrinology* 137:4189–4195, 1996
23. Altshuler D, Hirschhorn JN, Klannemark M, Lindgren CM, Vohl M-C, Nemesh J, Lane CR, Schaffner SF, Bolk S, Brewer C, Tuomi T, Gaudet D, Hudson TJ, Daly M, Groop L, Lander ES: The common PPAR γ Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nat Genet* 26:76–80, 2000
24. McCarthy MI: Growing evidence for diabetes susceptibility genes from genome scan data. *Curr Diab Rep* 3:159–167, 2003
25. Hanis CL, Boerwinkle E, Chakraborty R, Ellsworth DL, Concannon P, Stirling B, Morrison VA, Wapelhorst B, Spielman RS, Gogolin-Ewens KJ, Shephard JM, Williams SR, Risch N, Hinds D, Iwasaki N, Ogata M, Omori Y, Petzold C, Rietzsch H, Schroder H-E, Schulze J, Cox NJ, Menzel S, Boriraj W, Chen X, Lim LR, Lindner T, Mereu LE, Wang Y-Q, Xiang K, Yamagata K, Yang Y, Bell GI: A genome-wide search for human non-insulin-dependent (type 2) diabetes genes reveals a major susceptibility locus on chromosome 2. *Nat Genet* 13:161–166, 1996
26. Horikawa Y, Oda N, Cox NJ, Li X, Orholm-Melander M, Hara M, Hinokio Y, Lindner TH, Mashima H, Schwarz PEH, del Bosque-Plata L, Horikawa Y, Oda Y, Yoshiuchi I, Colilla S, Polonsky KS, Wei S, Concannon P, Iwasaki N, Schulze J, Baier LJ, Bogardus C, Groop L, Boerwinkle E, Hanis CL, Bell GI: Genetic variation in the gene encoding calpain-10 is associated with type 2 diabetes mellitus. *Nat Genet* 26:163–175, 2000
27. Evans JC, Frayling TM, Cassell PG, Saker PJ, Hitman GA, Walker M, Levy JC, O'Rahilly S, Rao PVS, Bennett AJ, Jones EC, Menzel S, Prestwich P, Simecek N, Wishart M, Dhillon R, Fletcher C, Millward A, Demaine A, Wilkin T, Horikawa Y, Cox NJ, Bell GI, Ellard S, McCarthy MI, Hattersley AT: Studies of association between the gene for calpain-10 and type 2 diabetes mellitus in the United Kingdom. *Am J Hum Genet* 69:544–552, 2001
28. Weedon MN, Schwaz PE, Horikawa Y, Iwasaki N, Illig T, Holle R, Rathmann W, Selisko T, Schulze J, Owen KR, Evans J, del Bosque-Plata L, Hitman G, Walker M, Levy JC, Sampson M, Bell GI, McCarthy MI, Hattersley AT, Frayling TM: Meta-analysis and a large association study confirm a role for calpain-10 variation in type 2 diabetes susceptibility. *Am J Hum Genet* 73:1208–1212, 2003
29. Grant SF, Thorleifsson G, Reynisdottir I, Benediktsson R, Manolescu A, Sainz J, Helgason A, Stefansson H, Emilsson V, Helgadóttir A, Styrkarsdóttir U, Magnusson KP, Walters GB, Palsdóttir E, Jonsdóttir T, Gudmundsdóttir T, Gylfason A, Saemundsdóttir J, Wilensky RL, Reilly MP, Rader DJ, Bagger Y, Christiansen C, Gudnason V, Sigurdsson G, Thorsteinsdóttir U, Gulcher JR, Kong A, Stefansson K: Variant of transcription factor 7-like 2 (TCFL2) gene confers risk of type 2 diabetes. *Nat Genet* 38:320–323, 2006
30. Reynisdottir I, Thorleifsson G, Benediktsson R, Sigurdsson G, Emilsson V, Einarsdóttir AS, Hjorleifsdóttir EE, Orlygsdóttir GT, Bjornsdóttir GT, Saemundsdóttir J, Halldorsson S, Hrafnkelsdóttir S, Sigurjonsdóttir SB, Steinsdóttir S, Martin M, Kochan JP, Rhees BK, Grant SFA, Frigge ML, Kong A, Gudnason V, Stefansson k, Gulcher JR: Localization of a susceptibility gene for type 2 diabetes to chromosome 5q34–q35.2. *Am J Hum Genet* 73:323–335, 2003
31. McVie-Wyllie AJ, Lamson DR, Chen YT: Molecular cloning of a novel member of the GLUT family of transporters, SLC2A10 (GLUT10), localized on chromosome 20q13.1: a candidate gene for NIDDM susceptibility. *Genomics* 72:113–117, 2001
32. Anderson G, Rose CS, Hamid YH, Drivsholm T, Borch-Johnsen K, Hansen T, Pedersen O: Genetic variation of the GLUT10 glucose transporter (SLC2A10) and relationships to type 2 diabetes and intermediary traits. *Diabetes* 52:2445–2448, 2003
33. Mohlke KL, Skol AD, Scott LJ, Valle TT, Bergman RN, Tuomilehto J, Boehnke M, Collins FS, FUSION Study Group: Evaluation of SLC2A10 (GLUT10) as a candidate gene for type 2 diabetes and related traits in Finns. *Mol Genet Metab* 85:323–327, 2005
34. Bento JL, Palmer ND, Mychaleckyj JC, Lange LA, Langefeld CD, Rich SS, Freedman BI, Bowden DW: Association of protein tyrosine phosphatase 1B gene polymorphisms with type 2 diabetes. *Diabetes* 53:3007–3012, 2004
35. Palmer ND, Bento JL, Mychaleckyj JC, Langefeld CD, Campbell JK, Norris JM, Haffner SM, Bergman RN, Bowden DW: Association of protein tyrosine phosphatase 1B gene polymorphisms with measures of glucose homeostasis in Hispanic Americans: the Insulin Resistance Atherosclerosis Study (IRAS) Family Study. *Diabetes* 53:3013–3019, 2004
36. Florez JC, Agapakis CM, Burt NP, Sun M, Almgren P, Råstam L, Tuomi T, Gaudet D, Hudson TJ, Daly MJ, Ardlie KG, Hirschhorn JN, Groop L, Altshuler D: Association testing of the protein tyrosine phosphatase 1B gene (PTPN1) with type 2 diabetes in 7,883 people. *Diabetes* 54:1884–1891, 2005
37. Valle T, Tuomilehto J, Bergman RN, Ghosh S, Hauser ER, Eriksson J, Nylund SJ, Kohtamäki K, Toivanen L, Vidgren G, Tuomilehto-Wolf E, Ehnholm C, Blachak J, Langefeld CD, Watanabe RM, Magnuson V, Ally DS, Hagopian WA, Ross E, Buchanan TA, Collins F, Boehnke M: Mapping genes for NIDDM: design of the Finland-United States Investigation of NIDDM Genetics (FUSION) Study. *Diabetes Care* 21:949–958, 1998
38. Zouali H, Hani EH, Philippi A, Vionnet N, Beckmann JS, Demenais F, Froguel P: A susceptibility locus for early-onset non-insulin dependent (type 2) diabetes mellitus maps to chromosome 20q, proximal to the phosphoenolpyruvate carboxylase gene. *Hum Mol Genet* 6:1401–1408, 1997
39. Bowden DW, Sale M, Howard TD, Qadri A, Spray BJ, Rothschild CB, Akots G, Rich SS, Freedman BI: Linkage of genetic markers on human chromosomes 20 and 12 to NIDDM in Caucasian sib pairs with a history of diabetic nephropathy. *Diabetes* 46:882–886, 1997
40. Ji L, Malecki M, Warram JH, Yang Y, Rich SS, Krolewski AS: New susceptibility locus for NIDDM is localized to human chromosome 20q. *Diabetes* 46:876–881, 1997
41. Klupa T, Malecki MT, Pezolesi M, Ji L, Curtis S, Langefeld CD, Rich SS, Warram JH, Krolewski AS: Further evidence for a susceptibility locus for type 2 diabetes on

- chromosome 20q13.1-q13.2. *Diabetes* 49:2212–2216, 2000
42. Ghosh S, Watanabe RM, Hauser ER, Valle T, Magnuson VL, Erdos MR, Langefeld CD, Balow J Jr, Ally DS, Kohtamäki K, Chines P, Birznieks G, Kaleta HS, Musick A, Te C, Tannenbaum J, Eldridge W, Shapiro S, Martin C, Witt A, So A, Chang J, Shurtleff B, Porter R, Kudelko K: Type 2 diabetes: evidence for linkage on chromosome 20 in 716 Finnish affected sib pairs. *Proc Natl Acad Sci U S A* 96:2198–2203, 1999
 43. Silander K, Mohlke KL, Scott LJ, Peck EC, Hollstein P, Skol AD, Jackson AU, Deloukas P, Hunt S, Stavrides G, Chines PS, Erdos MR, Narisu N, Conneely KN, Li C, Fingerlin TE, Dhanjal SK, Valle TT, Bergman RN, Tuomilehto J, Watanabe RM, Boehnke M, Collins FS: Genetic variation near the hepatocyte nuclear factor-4 α gene predicts susceptibility to type 2 diabetes. *Diabetes* 53:1141–1149, 2004
 44. Love-Gregory LD, Wasson J, Ma J, Jin CH, Glaser B, Suarez BK, Permutt MA: A common polymorphism in the upstream promoter region of the hepatocyte nuclear factor-4 α gene on chromosome 20q is associated with type 2 diabetes and appears to contribute to the evidence for linkage in an Ashkenazi Jewish population. *Diabetes* 53:1134–1140, 2004
 45. Damcott CM, Hoppman N, Ott SH, Reinhardt LJ, Wang J, Pollin TI, O'Connell JR, Mitchell BD, Shuldiner AR: Polymorphisms in both promoters of hepatocyte nuclear factor 4- α are associated with type 2 diabetes in the Amish. *Diabetes* 53:3337–3341, 2004
 46. Hansen SK, Rose CS, Glümer C, Drivsholm T, Borch-Johnsen K, Jørgensen T, Pedersen O, Hansen T: Variation near the hepatocyte nuclear factor (HNF)-4 α gene associates with type 2 diabetes in the Danish population. *Diabetologia* 48:452–458, 2005
 47. Bagwell AM, Bento JL, Mychaleckyj JC, Freedman BI, Langefeld CD, Bowden DW: Genetic analysis of HNF4A polymorphisms in Caucasian-American type 2 diabetes. *Diabetes* 54:1185–1190, 2005
 48. National Diabetes Data Group: Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes* 28:1039–1057, 1979
 49. World Health Organization: *Report of the Expert Committee on Diabetes*. Geneva, World Health Org., 1985 (Tech. Rep. Ser. no. 727)
 50. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 20:1183–1197, 1997
 51. Metzger BE: Summary and recommendation of the Third International Workshop-Conference on Gestational Diabetes Mellitus. *Diabetes* 40:197–201, 1991
 52. Rich SS: Mapping genes in diabetes. *Diabetes* 39:1315–1319, 1990
 53. Dorner G, Plagemann A, Reinagel H: Familial diabetes aggregation in type I diabetes: gestational diabetes an apparent risk factor for increased diabetes susceptibility in the offspring. *Exp Clin Endocrinol* 89:84–90, 1987
 54. McLellan JA, Barrow BA, Levy JC, Hammersley MS, Hattersley AT, Gillmer MD, Turner RC: Prevalence of diabetes mellitus and impaired glucose tolerance in parents of women with gestational diabetes. *Diabetologia* 38:693–698, 1995
 55. Martin AO, Simpson JL, Ober C, Freinkel N: Frequency of diabetes mellitus in mothers of probands with gestational diabetes: possible maternal influence on the predisposition to gestational diabetes. *Am J Obstet Gynecol* 151:471–475, 1985
 56. Freinkel N, Metzger BE, Phelps RL, Simpson JL, Martin AO, Radvany R, Ober C, Dooley SL, Depp RO, Belton A: Gestational diabetes mellitus: a syndrome with phenotypic and genotypic heterogeneity. *Horm Metab Res* 18:427–439, 1986
 57. Ober C, Xiang KS, Thisted RA, Indovina KA, Wason CJ, Dooley S: Increased risk for gestational diabetes mellitus associated with insulin receptor and insulin-like growth factor II restriction fragment length polymorphisms. *Genet Epidemiol* 6:559–569, 1989
 58. Stoffel M, Bell KL, Blackburn CL, Powerll KL, Seo TS, Takeda J, Vionnet N, Xiang KS, Gidh-Jain M, Pilkis SJ, et al.: Identification of glucokinase mutations in subjects with gestational diabetes mellitus. *Diabetes* 42:937–940, 1993
 59. Chiu KC, Go RC, Aoki M, Riggs AC, Tanizawa Y, Acton RT, Bell DS, Goldenberg RL, Roseman JM, Permutt MA: Glucokinase gene in gestational diabetes mellitus: population association study and molecular scanning. *Diabetologia* 37:104–110, 1994
 60. Saker PJ, Hattersley AT, Barrow B, Hammersley MS, McLellan JA, Lo YM, Olds RJ, Gillmer MD, Holman RR, Turner RC: High prevalence of a missense mutation of the glucokinase gene in gestational diabetic patients due to a founder-effect in a local population. *Diabetologia* 39:1325–1328, 1996
 61. Allan CJ, Argyropoulos G, Bowker M, Zhu J, Lin PM, Stiver K, Golichowski A, Garvey WT: Gestational diabetes mellitus and gene mutations which affect insulin secretion. *Diabetes Res Clin Pract* 36:135–141, 1997
 62. Ellard S, Beards F, Allen LI, Shepherd M, Ballantyne E, Harvey R, Hattersley AT: A high prevalence of glucokinase mutations in gestational diabetic subjects selected by clinical criteria. *Diabetologia* 43:250–253, 2000
 63. Weng J, Ekelund M, Lehto M, Li H, Ekberg G, Frid A, Åberg A, Groop LC, Bertorp K: Screening for MODY mutations, GAD antibodies, and type 1 diabetes-associated HLA genotypes in women with gestational diabetes mellitus. *Diabetes Care* 25:68–71, 2002
 64. Ryan EA, Imes S, Liu D, McManus R, Finegood DT, Polonsky KS, Sturis J: Defects in insulin secretion and action in women with a history of gestational diabetes. *Diabetes* 44:506–512, 1995
 65. Sakamaki H, Yamasaki H, Matsumoto K, Izumino K, Kondo H, Sera Y, Yamaguchi Y, Eguchi K: No deterioration in insulin sensitivity, but impairment of both pancreatic beta-cell function and glucose sensitivity, in Japanese women with former gestational diabetes mellitus. *Diabet Med* 15:1039–1044, 1998
 66. Xiang AH, Peters RK, Trigo E, Kjos SL, Lee WP, Buchanan TA: Multiple metabolic defects during late pregnancy in women at high risk for type 2 diabetes. *Diabetes* 48:848–854, 1999
 67. Buchanan TA: Pancreatic B-cell defects in gestational diabetes: implications for the pathogenesis and prevention of type 2 diabetes. *J Clin Endocrinol Metab* 86:989–993, 2001
 68. Sakul H, Pratley R, Cardon L, Ravussin E, Mott D, Bogardus C: Familiarity of physical and metabolic characteristics that predict the development of non-insulin-dependent diabetes mellitus in Pima Indians. *Am J Hum Genet* 60:651–656, 1997
 69. Watanabe RM, Valle T, Hauser ER, Ghosh S, Eriksson J, Kohtamäki K, Ehnholm C, Tuomilehto J, Collins FS, Bergman RN, Boehnke M: Familiarity of quantitative metabolic traits in Finnish families with non-insulin-dependent diabetes mellitus. *Hum Hered* 49:159–168, 1999
 70. Elbein SC, Hasstedt SJ, Wegner K, Kahn SE: Heritability of pancreatic β -cell function among nondiabetic members of Caucasian familial type 2 diabetic kindreds. *J Clin Endocrinol Metab* 84:1398–1403, 1999
 71. Watanabe RM, Xiang A, Trigo E, Hernandez S, Berrios F, Langefeld C, Kjos S, Ouzunian J, Sacks D, Lawrence J, Buchanan T: Evidence of genetic predisposition for B-cell dysfunction in Mexican-American families of probands with gestational diabetes (Abstract). *Diabetes* 52 (Suppl. 1):A257, 2003
 72. Watanabe RM, Xiang AH, Allayee H, Hartiala J, Trigo E, Wang C, Berrios F, Hernandez J, Paredes G, Hernandez M, Cercado S, Patel L, Caro J, Kjos SA, Lawrence JM, Buchanan TA: Variation in the P2-promoter region of hepatocyte nuclear factor-4 α (HNF4A) is associated with β -cell function in Mexican American families of a proband with gestational diabetes (Abstract). *Diabetes* 54 (Suppl. 1):A28, 2005
 73. Muller Y, Infante AM, Hanson RL, Love-

- Gregory L, Knowler W, Bogardus C, Baier L: Variants in hepatocyte nuclear factor 4 α are modestly associated with type 2 diabetes in Pima Indians. *Diabetes* 54: 3035–3039, 2005
74. Bergman RN, Phillips LS, Cobelli C: Physiologic evaluation of factors controlling glucose tolerance in man: measurement of insulin sensitivity and β -cell glucose sensitivity from the response to intravenous glucose. *J Clin Invest* 68:1456–1467, 1981
75. Odom DT, Zizlsperger N, Gordon DB, Bell GW, Rinaldi NJ, Murray HL, Volkert TL, Schreiber J, Rolfe PA, Gifford DK, Fraenkel E, Bell GI, Young RA: Control of pancreas and liver gene expression by HNF transcription factors. *Science* 303: 1378–1381, 2004
76. Risch N, Merikangas K: The future of genetic studies of complex human diseases. *Science* 273:1616–1617, 1996
77. Oliphant A, Barker DL, Stuelpnagel JR, Chee MS: BeadArray technology: enabling an accurate, cost-effective approach to high-throughput genotyping. *Biotechniques* (Suppl.):56–58, 2002
78. International Human Genome Mapping Consortium: A physical map of the human genome. *Nature* 409:934–941, 2001
79. The Genome Sequencing Consortium: Initial sequencing and analysis of the human genome. *Nature* 409:860–921, 2001
80. The International HapMap Consortium: A haplotype map of the human genome. *Nature* 437:1299–1320, 2005