

# High Serum Uric Acid as a Novel Risk Factor for Type 2 Diabetes

ABBAS DEGHAN, MD, DSC<sup>1</sup>  
MANDY VAN HOEK, MD<sup>2</sup>  
ERIC J.G. SIJBRANDS, MD, PHD<sup>2</sup>

ALBERT HOFMAN, MD, PHD<sup>1</sup>  
JACQUELINE C.M. WITTEMAN, PHD<sup>1</sup>

**OBJECTIVE**— To investigate the association between serum uric acid level and risk of type 2 diabetes.

**RESEARCH DESIGN AND METHODS**— The population for analysis consisted of 4,536 subjects free from diabetes at baseline. During a mean of 10.1 years of follow-up, 462 subjects developed diabetes.

**RESULTS**— The age- and sex-adjusted hazard ratios (HRs) (95% CIs) for diabetes were 1.30 (0.96–1.76) for the second, 1.63 (1.21–2.19) for the third, and 2.83 (2.13–3.76) for the fourth quartile of serum uric acid, in comparison with the first quartile. After adjustment for BMI, waist circumference, systolic and diastolic blood pressure, and HDL cholesterol, the HRs decreased to 1.08 (0.78–1.49), 1.12 (0.81–1.53), and 1.68 (1.22–2.30), respectively.

**CONCLUSIONS**— The results of this population-based study suggest that serum uric acid is a strong and independent risk factor for diabetes.

*Diabetes Care* 31:361–362, 2008

Serum uric acid is positively associated with serum glucose in healthy subjects.<sup>(1,2)</sup> However, this association is not consistent between healthy and diabetic individuals<sup>(3–5)</sup>, as a low serum level of uric acid is reported in the hyperglycemic state<sup>(6)</sup>. Since most individuals experience a phase of impaired glucose tolerance before progression to diabetes, it is not clear whether raised serum uric acid predicts the risk of type 2 diabetes<sup>(4,5)</sup>. We investigated the association between serum uric acid and risk of diabetes in the Rotterdam Study, a large population-based, prospective cohort study among subjects aged 55 years and older.

## RESEARCH DESIGN AND METHODS

The Rotterdam Study has been described in detail elsewhere<sup>(7)</sup>. Written informed consent was obtained from all participants, and the Medical Ethics Committee of Erasmus Medical Center approved the study. At baseline,

serum uric acid was measured<sup>(8,9)</sup> and prevalent diabetes cases were excluded<sup>(10)</sup>. Incident cases of diabetes were diagnosed during follow-up based on the guidelines of the American Diabetes Association<sup>(11)</sup> and World Health Organization<sup>(12)</sup> using the information from general practitioners, pharmacies' databases, and fasting blood samples that were taken during follow-up examinations<sup>(10)</sup>. The association was assessed by Cox regression analysis. The population-attributable risk and 95% CIs were calculated with use of the Interactive Risk Assessment Program (IRAP)<sup>(13)</sup>.

**RESULTS**— Serum uric acid was ranged from 107 to 756  $\mu\text{mol/l}$  with a mean  $\pm$  SD of  $323.7 \pm 82.2 \mu\text{mol/l}$ . Age BMI, waist circumference, systolic and diastolic blood pressure, and HDL cholesterol were significantly correlated with serum uric acid. The correlation coefficient ranged from 0.03 for diastolic blood pressure to

0.35 for waist circumference. Except for HDL cholesterol, the correlations were positive.

During a mean follow-up of 10.1 years, 462 of 4,536 participants developed diabetes (incidence rate 10.1 per 1,000 person-years). The age- and sex-adjusted hazard ratios (HRs) (95% CIs) for diabetes were 1.30 (0.96–1.76) for the second, 1.63 (1.21–2.19) for the third, and 2.83 (2.13–3.76) for the fourth quartile of serum uric acid, in comparison with the first quartile. In a multivariate-adjusted model (Table 1), the HRs (95% CIs) decreased to 1.08 (0.78–1.49), 1.12 (0.81–1.53), and 1.68 (1.22–2.30), respectively. The population-attributable risk of high serum uric acid for diabetes was 0.24 (95% CI 0.17–0.30) for the fourth quartile, 0.09 (0.3–0.15) for the third quartile, and 0.04 (–0.01 to 0.10) for the second quartile.

**CONCLUSIONS**— We showed that the subjects with higher levels of serum uric acid are more at risk of developing type 2 diabetes. We also found that one-quarter of diabetes cases can be attributed to a high serum uric acid level.

Our finding is in agreement with previous studies. At least two studies in 1980s reported on the association of serum uric acid with the risk of diabetes<sup>(14,15)</sup>; however, the association was not adjusted for any potential confounder. Lately, it has been shown in middle-aged Japanese men<sup>(4)</sup> that serum uric acid level is significantly associated with the risk of diabetes. However, another study in middle-aged Japanese men in Osaka showed that the association was not significant after adjustment for BMI, alcohol consumption, smoking, physical activity, fasting blood glucose, and parental history of diabetes. The absence of an independent effect in this study could be explained by the fact that the study population only consisted of men. We observed in our study that, although not significantly, the association was weaker in men than in women (data not shown).

Recognition of high serum uric acid as a risk factor for diabetes has been a matter of debate for a few decades, since hyperuricemia has been presumed to be a consequence of insulin resistance rather than its precursor. However, recent findings sug-

From <sup>1</sup>Department of Epidemiology and Biostatistics, Erasmus Medical Center, Rotterdam, the Netherlands; and the <sup>2</sup>Department of Internal Medicine, Erasmus Medical Center, Rotterdam, the Netherlands.

Address correspondence and reprint requests to Dr. J.C.M. Witteman, PhD, Epidemiology & Biostatistics, Erasmus Medical Center, Rotterdam, P.O. Box 2040, 3000 CA Rotterdam, Netherlands. E-mail: j.witteman@erasmusmc.nl.

Received for publication 6 July 2007 and accepted in revised form 23 October 2007.

Published ahead of print at <http://care.diabetesjournals.org> on 31 October 2007. DOI: 10.2337/dc07-1276.

© 2008 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—HR (95% CI) for diabetes according to level of serum uric acid

Serum uric acid quartile	Participants (cases)	HR (95% CI)		
		Model 1	Model 2	Model 3
1 ( $\leq 267$ $\mu\text{mol/l}$ )	1,153 (77)	1.00 (Ref.)	1.00 (Ref.)	1.00 (Ref.)
2 (268–310 $\mu\text{mol/l}$ )	1,141 (94)	1.30 (0.96–1.76)	1.14 (0.83–1.57)	1.08 (0.78–1.49)
3 (311–370 $\mu\text{mol/l}$ )	1,175 (120)	1.63 (1.21–2.19)	1.23 (0.89–1.67)	1.12 (0.81–1.53)
4 ( $> 370$ $\mu\text{mol/l}$ )	1,067 (171)	2.83 (2.13–3.76)	1.92 (1.41–2.62)	1.68 (1.22–2.30)
P for trend		<0.001	<0.001	<0.001
1 SD increment	4,536 (462)	1.53 (1.39–1.67)	1.37 (1.23–1.52)	1.31 (1.18–1.46)

Model 1: adjusted for age and sex. Model 2: model 1 + BMI and waist circumference. Model 3: model 2 + systolic and diastolic blood pressure and HDL cholesterol.

gest that uric acid could be related to the development of diabetes. Serum uric acid has been shown to be associated with oxidative stress (16) and production of tumor necrosis factor- $\alpha$  (16), which are both related to the development of diabetes. In addition, a recent study in rats showed that fructose-induced hyperuricemia plays a pathogenic role in the metabolic syndrome (17). These findings support high serum uric acid as a precursor of type 2 diabetes.

Currently, gout and renal disorders are the only consequences considered for hyperuricemia. Recent studies have introduced serum uric acid as a potential risk factor for hypertension (18), stroke (8), and cardiovascular diseases (19). Our findings suggest that type 2 diabetes is another consequence of hyperuricemia. The importance of this finding is even clearer when considering that lowering serum uric acid in subjects in the highest quartile may decrease the incidence of diabetes by 24%, if the relationship is causal. Hence, the public health impact of high serum uric acid may be larger than currently thought. Even so, uric acid is neither a target for treatment in asymptomatic hyperuricemia nor a risk marker in clinical practice (20), but methods for assessment of serum uric acid are widely available and inexpensive. Moreover, xanthine oxidase inhibitors, which are currently used to decrease serum uric acid, are safe and inexpensive.

In conclusion, our findings, together with those from previous literature, indicate that lowering uric acid may be a novel treatment target for preventing diabetes and justify a prospective clinical trial on the possible benefits of the measurement and lowering of serum uric acid on multiple chronic disease end points.

**Acknowledgments**—The Rotterdam Study is supported by the Erasmus Medical Center and Erasmus University Rotterdam; the Netherlands

Organization for Scientific Research; the Netherlands Organization for Health Research and Development (ZonMw); the Research Institute for Diseases in the Elderly; the Ministry of Education, Culture and Science; the Ministry of Health, Welfare and Sports; the European Commission; and the Municipality of Rotterdam. A.D. is supported by a scholarship from Hormozgan University of Medical Science, Bandar Abbas, Iran.

#### References

1. Modan M, Halkin H, Karasik A, Lusky A: Elevated serum uric acid: a facet of hyperinsulinaemia. *Diabetologia* 30:713–718, 1987
2. Facchini F, Chen YD, Hollenbeck CB, Reaven GM: Relationship between resistance to insulin-mediated glucose uptake, urinary uric acid clearance, and plasma uric acid concentration. *JAMA* 266:3008–3011, 1991
3. Wun YT, Chan CSY, Lui CS: Hyperuricaemia in type 2 diabetes mellitus. *Diabetes Nutr Metab* 12:286–291, 1999
4. Nakanishi N, Okamoto M, Yoshida H, Matsuo Y, Suzuki K, Tataru K: Serum uric acid and risk for development of hypertension and impaired fasting glucose or type II diabetes in Japanese male office workers. *Eur J Epidemiol* 18:523–530, 2003
5. Taniguchi Y, Hayashi T, Tsumura K, Endo G, Fujii S, Okada K: Serum uric acid and the risk for hypertension and type 2 diabetes in Japanese men: the Osaka Health Survey. *J Hypertens* 19:1209–1215, 2001
6. Nan H, Dong Y, Gao W, Tuomilehto J, Qiao Q: Diabetes associated with a low serum uric acid level in a general Chinese population. *Diabetes Res Clin Pract* 2006
7. Hofman A, Grobbee DE, de Jong PT, van den Ouweland FA: Determinants of disease and disability in the elderly: the Rotterdam Elderly Study. *Eur J Epidemiol* 7:403–422, 1991
8. Bos MJ, Koudstaal PJ, Hofman A, Witteman JCM, Breteler MMB: Uric acid is a risk factor for myocardial infarction and stroke: the Rotterdam Study. *Stroke* 37:1503–1507, 2006
9. Trivedi R, Rebar L, Berta E, Stong L: New

- enzymatic method for serum uric acid at 500 nm. *Clin Chem* 24:1908–1911, 1978
10. Dehghan A, Kardys I, de Maat MP, Uitterlinden AG, Sijbrands EJ, Bootsma AH, Stijnen T, Hofman A, Schram MT, Witteman JC: Genetic variation, C-reactive protein levels, and incidence of diabetes. *Diabetes* 56:872–878, 2007
11. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 20:1183–1197, 1997
12. World Health Organization: *Diabetes Mellitus Fact Sheet* 138. Geneva, World Health Org., 2002
13. Gail DM: *Interactive Risk Assessment Program*. 2.2 ed., 2002
14. Brand FN, McGee DL, Kannel WB, Stokes J III, Castelli WP: Hyperuricemia as a risk factor of coronary heart disease: the Framingham study. *Am J Epidemiol* 121:11–18, 1985
15. Herman JB, Goldbourt U: Uric acid and diabetes: observations in a population study. *Lancet* 2:240–243, 1982
16. Butler R, Morris AD, Belch JJ, Hill A, Struthers AD: Allopurinol normalizes endothelial dysfunction in type 2 diabetics with mild hypertension. *Hypertension* 35:746–751, 2000
17. Nakagawa T, Hu H, Zharikov S, Tuttle KR, Short RA, Glushakova O, Ouyang X, Feig DI, Block ER, Herrera-Acosta J, Patel JM, Johnson RJ: A causal role for uric acid in fructose-induced metabolic syndrome. *Am J Physiol Renal Physiol* 290:F625–F631, 2006
18. Johnson RJ, Feig DI, Herrera-Acosta J, Kang DH: Resurrection of uric acid as a causal risk factor in essential hypertension. *Hypertension* 45:18–20, 2005
19. Baker JF, Krishnan E, Chen L, Schumacher HR: Serum uric acid and cardiovascular disease: recent developments, and where do they leave us? *Am J Med* 118:816–826, 2005
20. Duffy WB, Senekjian HO, Knight TF, Weinman EJ: Management of asymptomatic hyperuricemia. *JAMA* 246:2215–2216, 1981