

# Implementing a State-Based Cardiovascular Disease and Diabetes Prevention Program

KARL K. VANDERWOOD, MPH  
TARYN O. HALL, MPH  
TODD S. HARWELL, MPH  
MARCENE K. BUTCHER, RD, CDE

STEVEN D. HELGERSON, MD, MPH  
ON BEHALF OF THE MONTANA  
CARDIOVASCULAR DISEASE AND DIABETES  
PREVENTION PROGRAM WORKGROUP\*

**OBJECTIVE** — To evaluate weight loss and cardiometabolic risk reduction achieved through an adapted Diabetes Prevention Program intervention among adults at high risk for cardiovascular disease (CVD) and diabetes.

**RESEARCH DESIGN AND METHODS** — Eight health care facilities implemented a group-based lifestyle intervention beginning in 2008. Participants attended 16 weekly core sessions followed by 6 monthly after core sessions.

**RESULTS** — A total of 1,003 participants were enrolled, 816 (81%) completed the core and 578 (58%) completed the after core. Of participants completing the core and after core, 45 and 49% achieved the 7% weight loss goal, respectively. There were significant improvements in blood pressure, fasting glucose, and LDL cholesterol among participants completing the intervention.

**CONCLUSIONS** — Our findings indicate it is feasible for state-coordinated CVD and diabetes prevention programs to achieve significant weight loss and improve cardiometabolic risk.

*Diabetes Care* 33:2543–2545, 2010

The Diabetes Prevention Program (DPP) and other studies demonstrated that lifestyle intervention can prevent the development of type 2 diabetes and reduce cardiometabolic risk among participants, prompting many countries to begin implementing efforts to translate these studies into practice (1–5). In 2008, the Montana Department of Public Health and Human Services (DPHHS) implemented an adapted DPP, and preliminary results demonstrated feasibility and effectiveness (6). This report describes weight loss and cardiometabolic risk improvement among participants completing the intervention.

**RESEARCH DESIGN AND METHODS** — A description of the early phase of this intervention has been published previously, and the initial cohort of participants is included in this report (6). In brief, the DPHHS funded eight health care facilities with recognized diabetes self-management education (DSME) programs beginning in 2008. Sites used trained health professionals as lifestyle coaches to provide the 16-session core followed by 6 monthly after core sessions (7). DPHHS staff provided technical assistance, data collection and analyses, and evaluation.

Overweight (BMI  $\geq 25$  kg/m<sup>2</sup>) adults, with medical clearance from their refer-

ring provider and one or more of the following cardiovascular disease (CVD) and diabetes risk factors were eligible: a previous diagnosis of pre-diabetes; impaired glucose tolerance or impaired fasting glucose; high blood pressure ( $\geq 130/85$  mmHg or treatment) or dyslipidemia (triglycerides  $>150$  mg/dl, LDL cholesterol  $>130$  mg/dl or treatment, or HDL cholesterol  $<40$  mg/dl men and  $<50$  mg/dl women); or a history of gestational diabetes mellitus or gave birth to a baby  $>9$  pounds.

Height, weight, blood pressure, fasting blood glucose, and lipid values were collected at enrollment and at completion of the core and after core. Participants were weighed at the beginning of each session and submitted self-monitoring records. Participants were considered core completers if they did not drop out or miss three or more consecutive sessions and after core completers if they had completed follow-up laboratory measurements at 10 months.

Institutional review board approval was not required by the DPHHS because previous research established the safety and efficacy of the lifestyle intervention and only de-identified data were used for analyses.

Participant data were analyzed using SAS 9.1 (SAS Institute, Cary, NC). Baseline characteristics were compared among all enrolled participants and core and after core completers; *t* tests were used to compare continuous variables, and  $\chi^2$  tests were used to compare dichotomous variables. We calculated the proportion of completers who met the physical activity goal of  $>150$  min/week in the core and 5 and 7% weight loss in the core and after core. In the core and after core, the last observed weight of completers was used to calculate mean weight loss for those not attending the final session. Paired *t* tests were used to assess mean weight loss, and the mean systolic and diastolic blood pressure, HDL cholesterol, LDL cholesterol, and fasting blood glucose from baseline to the end of the core (4-month follow-up) and after core (10-month follow-up). Bonferroni correction was applied to the level of

From the Montana Department of Public Health and Human Services, Helena, Montana.

Corresponding author: Karl K. Vanderwood, kvanderwood@mt.gov.

Received 11 May 2010 and accepted 21 August 2010. Published ahead of print at <http://care.diabetesjournals.org> on 30 August 2010. DOI: 10.2337/dc10-0862.

\*A complete list of the members of the Montana Cardiovascular Disease and Diabetes Prevention Program Workgroup can be found in an online appendix at <http://care.diabetesjournals.org/cgi/content/full/dc10-0862/DC1>.

The contents of this report are solely the responsibility of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Weight loss and cardiometabolic risk factor outcomes among all participants completing the core and after core lifestyle intervention at 4 and 10 months, Montana, 2008–2010

	Baseline	Completed core	P value*	Completed after core	P value†
Weight (kg)					
Completed core, n=816	99.2 ± 20.7	92.4 ± 20.0	<0.001	—	—
Completed after core, n=578	97.4 ± 20.4	90.3 ± 19.5	<0.001	89.7 ± 19.3	<0.001
Systolic blood pressure (mmHg)					
Completed core, n=684	133.6 ± 15.7	126.5 ± 14.9	<0.001	—	—
Completed after core, n=453	132.7 ± 15.4	125.9 ± 14.3	<0.001	127.1 ± 14.6	<0.001
Diastolic blood pressure (mmHg)					
Completed core, n=683	82.0 ± 11.0	78.6 ± 9.6	<0.001	—	—
Completed after core, n=452	81.2 ± 10.9	77.7 ± 9.1	<0.001	77.7 ± 9.2	<0.001
HDL cholesterol (mg/dl)					
Completed core, n=692	48.8 ± 12.0	46.3 ± 10.8	<0.001	—	—
Completed after core, n=488	49.1 ± 11.6	46.1 ± 10.7	<0.001	51.0 ± 11.8	<0.001
LDL cholesterol (mg/dl)					
Completed core, n=663	125.2 ± 34.6	114.9 ± 32.6	<0.001	—	—
Completed after core, n=473	123.2 ± 33.2	112.0 ± 32.0	<0.001	118.7 ± 31.2	<0.001
Fasting blood glucose (mg/dl)					
Completed core, n=613	101.5 ± 14.7	97.4 ± 12.7	<0.001	—	—
Completed after core, n=418	101.6 ± 14.9	96.9 ± 11.7	<0.001	96.9 ± 15.2	<0.001

Data are means ± SD. \*Results of the paired-samples *t* test, comparisons of 4-month and baseline values. †Results of the paired-samples *t* test, comparisons of 10-month and baseline values.

significance ( $\alpha = 0.003$ ) to control for the number of paired *t* tests calculated.

**RESULTS**— Between February 2008 and January 2010, 1,003 participants were enrolled in the intervention; 816 (81%) completed the core and 578 (58%) completed the after core. Mean ± SD attendance was 14.9 ± 1.6 sessions during the core and 3.7 ± 2.1 sessions during after the core. The age of enrolled participants was 52.3 ± 11.6, and 80% (*n* = 805) were female. Core completers were significantly older than those who did not complete the core and after core completers were significantly older, had a lower BMI at baseline, and were more likely to have diagnosed dyslipidemia at baseline than those completing only the core (supplementary Table, available in an online appendix at <http://care.diabetesjournals.org/cgi/content/full/dc10-0862/DC1>).

At the conclusion of the core, 45% of completers achieved the 7% weight loss goal, 66% achieved 5% weight loss, and 66% met the physical activity goal. Among the after core completers, 49% met the 7% weight loss goal, 64% achieved 5% weight loss, and 70% achieved the physical activity goal at the end of core.

Core and after core completers achieved significant improvements in weight, systolic and diastolic blood pres-

sure, LDL cholesterol, and fasting blood glucose and a significant reduction in HDL cholesterol at the end of core (Table 1). Significant improvements in HDL cholesterol were seen for those completing the after core. Participants with and without impaired glucose values at baseline achieved significant improvements in weight, blood pressure, LDL cholesterol, and blood glucose values at completion of the core and after core (data not shown).

**CONCLUSIONS**— Core and after core completers achieved significant reductions in weight and improvements in cardiometabolic risk. However, HDL decreased significantly at the end of the core but was followed by a significant increase for those completing the after core. Other studies have found similar results, indicating reductions in HDL during initial weight loss, followed by increased HDL levels during weight maintenance (8,9).

Our lifestyle intervention has a number of strengths, which support translating this research into practice. We included overweight adults with risk factor(s) for CVD or diabetes, rather than only adults with pre-diabetes, an approach supported by recommendations from the American Diabetes Association and American Heart Association, acknowledging the importance of addressing an individual's global risk for CVD

and diabetes (10). We also relied on physician referrals rather than time-consuming screening events. Finally, offering the DPP in groups allowed for greater participant enrollment than a one-on-one intervention. There are several limitations to our study. First, there was a dropout rate of 19 and 42% at the end of the core and after core, respectively. Second, we used a pre- and post-evaluation with no comparison group. Third, we relied on self-reported physical activity and diet measures. Fourth, we were unable to obtain laboratory measures for all participants. Last, our analyses only included participants completing the intervention, which differed from the DPP, in which an intention-to-treat analysis was used.

Coordinated state and national approaches to implement diabetes prevention programs are needed. A recent assessment of Montana DSME programs indicated that these programs have the capacity to provide diabetes prevention services, the primary barrier being lack of reimbursement (11). Other promising models in the U.S. include regional training and implementation centers in Pittsburgh, Pennsylvania, and Indianapolis, Indiana (12,13). Because of the large number of individuals at high risk for diabetes in the U.S., many prevention sites will be needed, including DSME programs and other settings.

**Acknowledgments**— This project was funded through the Montana State Legislature and supported through a cooperative agreement with the Centers for Disease Control and Prevention, Division of Diabetes Translation (U32/CCU822743-05) in Atlanta, Georgia.

No potential conflicts of interest relevant to this article were reported.

K.K.V. and T.O.H. researched data, contributed to discussion, wrote the manuscript, and reviewed/edited the manuscript. T.S.H. researched data, contributed to discussion, and reviewed/edited the manuscript. M.K.B. and S.D.H. researched data and reviewed/edited the manuscript.

Parts of this study were presented in abstract form at the 70th Scientific Sessions of the American Diabetes Association, Orlando, Florida, 25–29 June 2010.

We thank and acknowledge Susan Day from the Montana DPHHS for her work and support on this project. We also thank Carol Percy and Cathy Manus from ACKCO American Indian Professional Services for providing the initial training and technical assistance for this project.

## References

1. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403
2. Ratner R, Goldberg R, Haffner S, Marcovina S, Orchard T, Fowler S, Temprosa M. Impact of intensive lifestyle and metformin therapy on cardiovascular disease risk factors in the diabetes prevention program. *Diabetes Care* 2005;28:888–894
3. Saaristo T, Peltonen M, Keinänen-Kiukkaanniemi S, Vanhala M, Saltevo J, Niskanen L, Oksa H, Korpi-Hyövälti E, Tuomilehto J. National type 2 diabetes prevention programme in Finland: FIN-D2D. *Int J Circumpolar Health* 2007;66:101–112
4. Schwarz PEH, Reddy P, Greaves CJ, J Dunbar JA, Schwarz J. *Diabetes Prevention in Practice*. Dresden, Germany, TUMAINI Institute for Prevention Management, 2010, p. 45–56
5. Schwarz PE, Lindström J, Kissimova-Scarbeck K, Szybinski Z, Barengo NC, Peltonen M, Tuomilehto J. The European perspective of type 2 diabetes prevention: diabetes in Europe—prevention using lifestyle, physical activity and nutritional intervention (DE-PLAN) project. *Exp Clin Endocrinol Diabetes* 2008;116:167–172
6. Amundson HA, Butcher MK, Gohdes D, Hall TO, Harwell TS, Helgerson SD, Vanderwood KK. Translating the Diabetes Prevention Program into practice in the general community: findings from the Montana Cardiovascular Disease and Diabetes Prevention Program. *Diabetes Educ* 2009;35:209–210, 213–214, 216–220
7. Healthy Native Community Partnership. Native lifestyle balance curriculum [article online], 2008. Available from <http://www.hncp.org/wst/hpdp/NLB>. Accessed 8 March 2008
8. Rossner S, Bjorvell H. Early and late effects of weight loss on lipoprotein metabolism in severe obesity. *Atherosclerosis* 1987;64:125–130
9. Fox AA, Thompson JL, Butterfield GE, Gylfadottir U, Moynihan S, Spiller G. Effects of diet and exercise on common cardiovascular disease risk factors in moderately obese older women. *Am J Clin Nutr* 1996;63:225–233
10. Eckel RH, Kahn R, Robertson RM, Rizza RA. Preventing cardiovascular disease and diabetes: a call to action from the American Diabetes Association and the American Heart Association. *Circulation* 2006;113:2943–2946
11. Butcher MK, Vanderwood KK, Hall TO, Gohdes D, Helgerson SD, Harwell TS. Capacity of diabetes education programs to provide both diabetes self-management education and to implement diabetes prevention services. *J Public Health Manag Pract*. In press
12. Ackermann RT, Marrero DG. Adapting the Diabetes Prevention Program lifestyle intervention for delivery in the community: the YMCA model. *Diabetes Educ* 2007;33:69, 74–75, 77–78
13. Kramer MK, Kriska AM, Venditti EM, Miller RG, Brooks MM, Burke LE, Siminerio LM, Solano FX, Orchard TJ. Translating the Diabetes Prevention Program: a comprehensive model for prevention training and program delivery. *Am J Prev Med* 2009;37:505–511