

High-Sensitivity CRP Discriminates HNF1A-MODY From Other Subtypes of Diabetes

TIM J. McDONALD, MSc^{1,2}
 BEVERLEY M. SHIELDS, PhD¹
 JANE LAWRY, BSc²
 KATHARINE R. OWEN, MD^{3,4}

ANNA L. GLOYN, DPHIL^{3,4}
 SIAN ELLARD, PhD^{1,5}
 ANDREW T. HATTERSLEY, DM¹

OBJECTIVE—Maturity-onset diabetes of the young (MODY) as a result of mutations in hepatocyte nuclear factor 1- α (*HNF1A*) is often misdiagnosed as type 1 diabetes or type 2 diabetes. Recent work has shown that high-sensitivity C-reactive protein (hs-CRP) levels are lower in HNF1A-MODY than type 1 diabetes, type 2 diabetes, or glucokinase (GCK)-MODY. We aim to replicate these findings in larger numbers and other MODY subtypes.

RESEARCH DESIGN AND METHODS—hs-CRP levels were assessed in 750 patients (220 *HNF1A*, 245 *GCK*, 54 HNF4- α [*HNF4A*], 21 HNF1- β [*HNF1B*], 53 type 1 diabetes, and 157 type 2 diabetes).

RESULTS—hs-CRP was lower in HNF1A-MODY (median [IQR] 0.3 [0.1–0.6] mg/L) than type 2 diabetes (1.40 [0.60–3.45] mg/L; $P < 0.001$) and type 1 diabetes (1.10 [0.50–1.85] mg/L; $P < 0.001$), HNF4A-MODY (1.45 [0.46–2.88] mg/L; $P < 0.001$), GCK-MODY (0.60 [0.30–1.80] mg/L; $P < 0.001$), and HNF1B-MODY (0.60 [0.10–2.8] mg/L; $P = 0.07$). hs-CRP discriminated HNF1A-MODY from type 2 diabetes with hs-CRP < 0.75 mg/L showing 79% sensitivity and 70% specificity (receiver operating characteristic area under the curve = 0.84).

CONCLUSIONS—hs-CRP levels are lower in HNF1A-MODY than other forms of diabetes and may be used as a biomarker to select patients for diagnostic *HNF1A* genetic testing.

Diabetes Care 34:1860–1862, 2011

Maturity-onset diabetes of the young (MODY) is a rare monogenic form of diabetes and is often misdiagnosed as type 1 diabetes or type 2 diabetes (1,2). A correct genetic diagnosis of MODY is important for predicting the clinical course of disease, risk to relatives, and optimizing treatment. Therefore, cheap and novel biomarkers that help identify these patients are desirable.

The *CRP* gene has hepatocyte nuclear factor 1- α (*HNF1A*) binding sites in its promoter, and common variants in and

around *HNF1A* are associated with circulating high-sensitivity C-reactive protein (hs-CRP) levels (3,4). Recently, Owen et al. (5) demonstrated that hs-CRP levels were reduced in 31 HNF1A-MODY patients compared with type 1 diabetes, type 2 diabetes, glucokinase (GCK) MODY, and normal control subjects, making it potentially a useful clinical test.

We aim to replicate this initial study in a larger cohort of patients and assess whether the reduction in hs-CRP is also seen in patients with mutations of other

genes encoding hepatic nuclear factors (hepatocyte nuclear factor 4- α [*HNF4A*] and hepatocyte nuclear factor 4- β [*HNF1B*]).

RESEARCH DESIGN AND

METHODS—Additional information about laboratory methods, study participants, and statistical analysis can be found in the Supplementary Data.

Study participants

hs-CRP level was assessed in plasma from 750 patients: 540 with a confirmed genetic diagnosis of MODY (220 *HNF1A*, 245 *GCK*, 54 *HNF4A*, and 21 *HNF1B*), 53 patients with type 1 diabetes, and 157 patients with type 2 diabetes.

RESULTS—Patient characteristics are presented in Supplementary Table 1. Subjects (29/750; 10.5%) had hs-CRP results > 10 mg/L ($n = 16$ GCK-MODY, $n = 10$ type 2 diabetes, and $n = 3$ HNF4A-MODY) and were considered to have an acute inflammatory response and excluded from further analysis. This is in line with the protocol used by Owen et al. and previous studies of CRP (5,6).

HNF1A-MODY versus type 2 diabetes and type 1 diabetes

Plasma hs-CRP was significantly lower in HNF1A-MODY than type 1 diabetes and type 2 diabetes (median [IQR] 0.3 [0.1–0.6] versus 1.1 [0.5–1.85] mg/L [$P < 0.001$] and 1.4 [0.6–3.45] mg/L; $P < 0.001$), respectively (Fig. 1).

For discriminating HNF1A-MODY from type 2 diabetes, receiver operating characteristic (ROC) curve showed good discrimination (area under the curve [AUC] 0.84) and identified a cutoff hs-CRP of 0.75 mg/L with 79% sensitivity and 71% specificity (Supplementary Fig. 1). This equates to a positive predictive value (PPV) of 2.7% and a negative predictive value (NPV) of 99.7%, assuming a MODY prevalence of 1% (7). hs-CRP was less discriminative between HNF1A-MODY and type 1 diabetes with a ROC AUC of 0.78 (79% sensitivity and 67% specificity at a cutoff of < 0.75 mg/L). This equates to a PPV of 2.4% and an NPV of 99.7%.

From the ¹Peninsula College of Medicine and Dentistry (University of Exeter), Peninsula NIHR Clinical Research Facility, Exeter, Devon, U.K.; the ²Royal Devon and Exeter NHS Foundation Trust, Department of Clinical Chemistry, Exeter, Devon, U.K.; the ³University of Oxford, Oxford Centre for Diabetes, Endocrinology and Metabolism, Oxford, U.K.; the ⁴Churchill Hospital, Oxford National Institute for Health Research Biomedical Research Centre, Oxford, U.K.; and the ⁵Royal Devon and Exeter NHS Foundation Trust, Department of Molecular Genetics, Exeter, U.K.

Corresponding author: Tim J. McDonald, tim.mcdonald@rdefnhs.uk.

Received 16 February 2011 and accepted 14 May 2011.

DOI: 10.2337/dc11-0323

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc11-0323/-DC1>.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.




Figure 1—Boxplot to show serum hs-CRP in type 1 diabetes ($n = 53$), type 2 diabetes ($n = 157$), HNF1A-MODY ($n = 220$), HNF4A-MODY ($n = 54$), HNF1B-MODY ($n = 21$), and GCK-MODY ($n = 245$). Box, median and IQR; whiskers, data range. \circ , outliers ($>1.5 \times$ IQR); \star , extreme values ($>3 \times$ IQR). The data exclude all subjects with hs-CRP levels >10 mg/L.

HNF1A-MODY versus other MODY subtypes

hs-CRP was lower in HNF1A-MODY (median [IQR] 0.3 [0.1–0.6]) than GCK-MODY (1.45 [0.46–2.88] mg/L; $P < 0.001$), HNF4A-MODY (0.6 [0.3–1.8] mg/L; $P < 0.001$), and HNF1B-MODY, although this did not reach statistical significance (0.6 [0.1–2.85] mg/L; $P = 0.07$).

To discriminate HNF1A-MODY from HNF4A-MODY, ROC curve analysis identified a cutoff hs-CRP <0.55 mg/L (AUC 0.76) achieving a sensitivity of 71% and a specificity of 70%. Discriminating HNF1A-MODY from all other MODY subtypes (HNF4A-, HNF1B-, GCK-MODY), a cutoff hs-CRP <0.55 mg/L (AUC 0.68) achieved 71% sensitivity and 55% specificity. This relationship persisted when discriminating HNF1A-MODY from all types of diabetes (71% sensitivity, 69% specificity).

Analysis of potential confounders can be found in the Supplementary Results.

CONCLUSIONS—We have confirmed in a large cohort of patients that plasma hs-CRP levels are lower in HNF1A-MODY than type 2 diabetes, type 1 diabetes, and GCK-MODY (5). We have shown for the first time that low hs-CRP levels are not seen in MODY

because of mutations in other related transcription factors (HNF4A, HNF1B).

The best discrimination achieved with hs-CRP is between HNF1A-MODY and type 2 diabetes (79% sensitivity, 70% specificity). However, because MODY is rare ($\sim 1\%$ of all diabetes [7]) compared with type 2 diabetes, this equates to a modest PPV of only 2.7%. Therefore, if hs-CRP was used as a marker for identifying HNF1A-MODY (hs-CRP <0.75 mg/L) in isolation the false-positive rate and number of unnecessary genetic testing would be high (37 to 1), suggesting that hs-CRP should be used in combination with other clinical characteristics. In contrast, a negative result (hs-CRP >0.75 mg/L), gives an NPV of 99.7% and therefore is good at ruling out HNF1A-MODY. Unlike type 1 diabetes where robust biomarkers, including pancreatic auto-antibodies and persistent C-peptide production (1,8), already exist, few biomarkers currently distinguish type 2 diabetes from MODY.

hs-CRP was only reduced in HNF1A-MODY patients and not in the other genetic subtypes of MODY. This result may have been predicted since the hs-CRP gene has HNF1A binding sites in its promoter (9). This may prove to be useful as a tool to help prioritize the order of

gene testing in patients with a high clinical suspicion of MODY. An hs-CRP level <0.55 mg/L should suggest HNF1A sequencing first, and levels >0.55 mg/L may warrant consideration of the other subtypes of MODY. This result may prove to be helpful as an adjunct to identify HNF4A-MODY patients, who are phenotypically similar to HNF1A-MODY.

An advantage of hs-CRP is that it is cheap and widely available. Current CRP assays, used to identify inflammation, measure down to levels of 0.2–0.3 mg/L and, therefore, a specific hs-CRP assay may not be necessary. It would be important to determine the comparability between routine CRP and hs-CRP assays before testing. A limitation of CRP is that it will be raised in inflammatory states, reducing its potential use as a discriminative tool. In addition, certain medications reduce CRP levels, such as statins (10), aspirin (11), and β -blockers (12), which are believed in some cases to reduce CRP by 20–30%. We did not have drug history available on individual patients, so we were unable to assess the impact of pharmacological agents on our results; however, Owen et al. (5) showed that removing subjects on statins and/or aspirin in their study did not alter their results.

In conclusion, we have confirmed that hs-CRP levels are lower in patients with HNF1A-MODY than other types of MODY, type 1 diabetes, and type 2 diabetes. hs-CRP is potentially a cheap, widely available biomarker that might aid in the cost-effective identification of patients with HNF1A-MODY.

Acknowledgments—This study was funded in Exeter by the European Community FP7 program CEED3 (HEALTH-F2-2008-223211). S.E., B.M.S., and A.T.H. are supported by the NIHR Peninsula Clinical Research Facility, University of Exeter. A.L.G. is supported by a Medical Research Council New Investigator Award (81696). K.R.O. is an NIHR-funded Clinician Scientist.

No potential conflicts of interest relevant to this article were reported.

T.J.M. wrote the manuscript and researched data. B.M.S. researched data and reviewed and edited the manuscript. J.L. researched data. K.R.O., A.L.G., and S.E. reviewed and edited the manuscript. A.T.H. researched data and reviewed and edited the manuscript.

The authors thank Annie Goodship (Royal Devon and Exeter NHS Foundation Trust, Department of Clinical Chemistry, Exeter, Devon, U.K.), Kevin Colclough (Royal Devon and Exeter NHS Foundation Trust, Department

of Molecular Genetics, Exeter, Devon, U.K.), and Elizabeth Wren (Royal Devon and Exeter NHS Foundation Trust, Department of Clinical Chemistry, Exeter, Devon, U.K.) for their technical assistance with the study.

References

1. Ellard S, Bellanné-Chantelot C, Hattersley AT; European Molecular Genetics Quality Network (EMQN) MODY group. Best practice guidelines for the molecular genetic diagnosis of maturity-onset diabetes of the young. *Diabetologia* 2008;51:546–553
2. Lambert AP, Ellard S, Allen LI, et al. Identifying hepatic nuclear factor 1- α mutations in children and young adults with a clinical diagnosis of type 1 diabetes. *Diabetes Care* 2003;26:333–337
3. Reiner AP, Barber MJ, Guan Y, et al. Polymorphisms of the HNF1A gene encoding hepatocyte nuclear factor-1 alpha are associated with C-reactive protein. *Am J Hum Genet* 2008;82:1193–1201
4. Ridker PM, Pare G, Parker A, et al. Loci related to metabolic-syndrome pathways including LEPR, HNF1A, IL6R, and GCKR associate with plasma C-reactive protein: the Women's Genome Health Study. *Am J Hum Genet* 2008;82:1185–1192
5. Owen KR, Thanabalasingham G, James TJ, et al. Assessment of high-sensitivity C-reactive protein levels as diagnostic discriminator of maturity-onset diabetes of the young due to HNF1A mutations. *Diabetes Care* 2010;33:1919–1924
6. Lee CC, Adler AI, Sandhu MS, et al. Association of C-reactive protein with type 2 diabetes: prospective analysis and meta-analysis. *Diabetologia* 2009;52:1040–1047
7. Shepherd M, Ellis I, Ahmad AM, et al. Predictive genetic testing in maturity-onset diabetes of the young (MODY). *Diabet Med* 2001;18:417–421
8. Besser REJ, Shepherd MH, McDonald TJ, Shields BM, Knight BA, Ellard S, Hattersley AT. Urinary C-peptide creatinine ratio is a practical outpatient tool for identifying hepatocyte nuclear factor 1- α /hepatocyte nuclear factor 4- α maturity-onset diabetes of the young from long-duration type 1 diabetes. *Diabetes Care* 2011;34:286–291
9. Toniatti C, Demartis A, Monaci P, Nicosia A, Ciliberto G. Synergistic trans-activation of the human C-reactive protein promoter by transcription factor HNF-1 binding at two distinct sites. *EMBO J* 1990;9:4467–4475
10. Mora S, Musunuru K, Blumenthal RS. The clinical utility of high-sensitivity C-reactive protein in cardiovascular disease and the potential implication of JUPITER on current practice guidelines. *Clin Chem* 2009;55:219–228
11. Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* 1997;336:973–979
12. Jenkins NP, Keevil BG, Hutchinson IV, Brooks NH. Beta-blockers are associated with lower C-reactive protein concentrations in patients with coronary artery disease. *Am J Med* 2002;112:269–274