

Elevated Depression Symptoms, Antidepressant Medicine Use and Risk of Developing Diabetes during the Diabetes Prevention Program

Richard R. Rubin, PhD^{1,2}
Yong Ma, MS³
David G. Marrero, PhD⁴
Mark Peyrot, PhD^{1,5}
Elizabeth L. Barrett-Connor, MD⁶
Steven E. Kahn, MD, ChB⁷
Steven M Haffner, MD, MPH⁸
David W. Price, MD⁹
William C. Knowler, MD, DrPH¹⁰

For The Diabetes Prevention Program Research Group*

From the¹ Department of Medicine, The Johns Hopkins University School of Medicine, Baltimore, Maryland; ² Department of Pediatrics, The Johns Hopkins University School of Medicine, Baltimore, Maryland; ³ Biostatistics Center, George Washington University, Rockville, Maryland; ⁴ Department of Medicine, Indiana University School of Medicine, Indianapolis, Indiana; ⁵ Department of Sociology, Loyola College of Maryland, Baltimore, Maryland; ⁶ Department of Family and Preventative Medicine, University of California at San Diego, La Jolla, California; ⁷ Department of Medicine, University of Washington, Seattle Washington; ⁸ Department of Medicine, Clinical Epidemiology, University of Texas Health Science Center, San Antonio, Texas; ⁹ Department of Family Medicine, University of Colorado Denver School of Medicine, Denver, Colorado; ¹⁰ The National Institute of Diabetes and Digestive and Kidney Diseases, Phoenix, Arizona.

* Members of the DPP Research Group appear in an on-line appendix

Running title: Depression symptoms, antidepressant medication, and diabetes risk

Corresponding Author:

Richard R. Rubin
Johns Hopkins University School of Medicine
946 E. Piney Hill Rd
Monkton, Maryland 21111
rrubin4@jhmi.edu

Received for publication 17 September 2007 and accepted in revised form 30 November 2007.

Additional information for this article can be found in an online appendix at <http://care.diabetesjournals.org>.

ABSTRACT

Objective: To assess the association between elevated depression symptoms or antidepressant medicine use on entry to the Diabetes Prevention Program (DPP) and during the study, and the risk of developing diabetes during the study.

Methods: DPP participants (N= 3187) in 3 treatment arms [intensive lifestyle (ILS), metformin (MET), placebo (PLB)] completed the Beck Depression Inventory (BDI) and reported their use of antidepressant medicines, at randomization and throughout the study (average duration in study = 3.2 yr).

Findings: When other factors associated with the risk of developing diabetes were controlled, elevated BDI scores at baseline or during the study were not associated with diabetes risk in any arm. Baseline antidepressant use was associated with diabetes risk in the PLB [Hazard Ratio (HR) = 2.25 (95% CI: 1.38-3.66) and ILS [HR= 3.48 (95% CI: 1.93-6.28)] arms. Continuous antidepressant use during the study (compared with no use) was also associated with diabetes risk in the same arms [PLB HR= 2.60 (95% CI: 1.37-4.94), ILS HR= 3.39 (95% CI: 1.61-7.13), as was intermittent antidepressant use during the study in the ILS arm [HR= 2.07 (95% CI: 1.18-3.62)]. Among MET arm participants, antidepressant use was not associated with developing diabetes.

Conclusions: A strong and statistically significant association between antidepressant use and diabetes risk in the PLB and ILS arms was not accounted for by measured confounders or mediators. If future research finds that antidepressant use independently predicts diabetes risk, efforts to minimize the negative effects of antidepressant agents on glycemic control should be pursued.

The concurrence of depression and diabetes is a serious problem. Among people with diabetes – whose risk of depression is 50% to 100% greater than the general population (1) – depression is associated with higher complication (2) and mortality rates (3), and higher health care costs (4). In 1674, Thomas Willis speculated that diabetes was *caused* by “long sorrow and other depressions” (5). Findings consistent with Willis’ hypothesis, as it applies to type 2 diabetes, were reported recently in a meta-analysis (6), but not all studies included in the meta-analysis reported positive findings. Some studies lacked sufficient statistical power or objective determination of baseline diabetes status. Even studies with positive findings suggested that they applied only to individuals with high levels of depression symptoms, or to certain demographic groups.

To our knowledge, no studies have considered the effects of elevated depression symptoms on diabetes risk in a large sample of individuals at high risk for developing diabetes because they have impaired glucose regulation or obesity. In addition, very few studies have identified an independent increment in the risk of developing diabetes associated with use of antidepressant medication (7), despite evidence suggesting that such medications may have an impact on glucose tolerance. Tricyclics appear to contribute to weight gain (8), and to increase blood glucose levels in patients with diabetes (9). Selective serotonin reuptake inhibitors (SSRI) and related agents appear to be associated with less weight gain, and perhaps to improve insulin sensitivity (10).

Data from the Diabetes Prevention Program (DPP) offer an opportunity to better understand the natural history of elevated depression symptoms, antidepressant medication, and type 2 diabetes in a large racially/ethnically diverse cohort of overweight individuals with elevated fasting

glucose and impaired glucose tolerance. The protocol included objective measures of glucose tolerance and potential mediators of the effect of depression symptoms or antidepressant medication on diabetes risk, including weight, weight change, physical activity, and insulin secretion and resistance.

We reported earlier that on entry to the DPP 10.3% of participants had symptoms consistent with at least mild depression, and 5.7% reported taking antidepressants, with 0.9% reporting both elevated symptoms and antidepressant use (11). Depression symptoms were higher among women and those having less education; antidepressant use was more common among women, those having more education, and those of white race/ethnicity. During the study the proportion of participants taking antidepressants increased (to 8.7% at year 3), while the proportion with elevated depression symptoms declined (to 8.4% at year 3), with no significant differences among DPP treatment arms.

The current study was designed to determine: 1) whether depression symptoms or antidepressant medicine use were associated with progression to type 2 diabetes in the DPP cohort, 2) if these associations varied by DPP treatment arm, and 3) whether these associations were mediated by potential diabetes risk factors.

RESEARCH DESIGN AND METHODS

The DPP involved persons at high risk for developing type 2 diabetes, and was conducted at 27 centers. DPP methods (12-13) and results (14) are described in detail elsewhere; the protocol is available at <http://www.bsc.gwu.edu/dpp>. At each DPP center an institutional review board approved the protocol; all participants gave written informed consent.

Participants. This analysis is based on the 3187 of 3234 enrolled DPP participants who

completed the Beck Depression Inventory (BDI) at baseline. Participants were ≥ 25 years of age, had a body-mass index of ≥ 24 kg/m² (≥ 22 kg/m² in Asian Americans), and a plasma glucose concentration of 95 to 125 mg/dl (5.3 to 6.9 mmol/l) in the fasting state (≤ 125 mg/dl in American Indians), and 140 to 199 mg/dl (7.8 to 11.0 mmol/l) two hours after a 75-g oral glucose load. Individuals were excluded from the study if they were taking antidepressant medication that might contribute to weight loss - bupropion or a selective serotonin reuptake inhibitor (SSRI) at more than the lowest usual dose (i.e., > 20 mg of fluoxetine or the equivalent) (15), if they had conditions that could seriously reduce their ability to participate in the DPP (including major psychiatric disorders), or if they could not successfully complete the 3-week run-in period during which participants took placebo medicines and recorded eating and activity. Recruitment was designed to randomize approximately half the participants from racial/ethnic minority groups.

Interventions. Eligible participants were randomly assigned to one of three interventions: standard lifestyle recommendations plus metformin at a dose of 850 mg twice daily (MET arm), standard lifestyle recommendations plus a placebo pill twice daily (PLB arm), or an intensive lifestyle modification program (ILS arm). Goals for ILS participants were to achieve and maintain a reduction of $\geq 7\%$ of initial body weight through a calorie-controlled, low-fat diet, and to engage in physical activity of moderate intensity, such as brisk walking, for ≥ 150 minutes per week (16).

Outcomes. As part of a comprehensive protocol, DPP participants completed the BDI (17) and a measure of leisure activity (18) prior to randomization and subsequently at each annual visit. Participants brought all prescription medicines, including antidepressants, to each quarterly clinic visit, and all were recorded.

Diabetes was diagnosed according to the 1997 American Diabetes Association criteria (19) based on an annual oral glucose-tolerance test or a semiannual fasting plasma glucose test. A confirmation test was performed, usually within 6 weeks (12). Fasting insulin was measured at annual visits with the OGTT. Detailed measurement methods for glucose and insulin have been previously published (20).

We report follow-up through July 2001, after which the primary results were announced and the interventions unmasked. This was four months longer than the results reported previously (14), resulting in a total mean follow-up of 3.2 years per participant.

We identified participants with BDI scores ≥ 11 as having elevated depression symptoms; others have chosen scores ranging from 10 to 16 (21-23), generally as a function of the importance placed on depression recognition. We chose a symptom threshold toward the low end of the severity range because we believed few severely depressed participants would pass the screening process for eligibility in the DPP. We identified participants who had elevated BDI scores at baseline and those who ever had elevated BDI scores during the DPP. We used this measure of depression symptoms (rather than the absolute BDI score) because we were interested in assessing the association between possible depression (i.e. elevated symptoms) and diabetes risk.

Analysis. At baseline, categorical variables are reported as percentages and continuous variables as mean \pm SD. Chi-square tests were used to compare baseline differences in categorical variables. To test for baseline differences among groups on continuous variables, the Wilcoxon rank sum test (24) was used for those with non-normal distributions; otherwise the student's t-test was used. During DPP, anti-depressant use was reported quarterly and depression symptoms annually. Mixed effects modeling

(25) was used to compare the difference in continuous variables such as weight by antidepressant use or depression symptoms and least square means with standard errors were reported. For categorical variables such as sex, repeated measures modeling with GEE (26) was used to compare differences by antidepressant use or elevated depression symptoms.

Cox proportional hazard models (27) were used to evaluate whether elevated depression symptoms or taking antidepressants were associated with developing diabetes. Having elevated depression symptoms during the DPP was defined as a time-dependent covariate: ever having BDI score ≥ 11 up to each time point evaluated. Antidepressant use during DPP was also defined as a time-dependent categorical variable up to each time point evaluated with three levels: never used, used intermittently (at least once but not always), and used continuously (at all visits). Time dependent covariate analyses (28) were used to model the above covariates and diabetes risk with adjustment for race/ethnicity and factors associated with an increased risk of developing diabetes (age, sex, education, fasting plasma glucose at baseline, weight at baseline and weight change during the study).

The life-table method (29) was used to estimate cumulative diabetes incidence rates at three years. These rates and the Greenwood estimate (30) of the standard error were used to calculate the number of persons needed to treat (NNT) with an antidepressant to associated with one case of diabetes and its 95% confidence interval.

All analyses were performed using the Statistical Analysis System (SAS Institute, Cary, NC).

RESULTS

At baseline (Table 1), 10.3% of DPP participants had BDI scores indicating at least mild depression (≥ 11), and 5.7% were taking

antidepressants. During the DPP (Table 1), intermittent antidepressant use was reported for 6.5% of total person-years, and continuous antidepressant use for 3.4% of total person-years.

At baseline the median BDI score for all three treatment arms was 3 with an interquartile range of 1-7; at follow-up the median BDI score for all three treatment arms was 2 with an interquartile range of 0-5 for the ILS arm and 0-6 for the MET and PLB arms. Elevated BDI scores at baseline and during the study were associated with: 1) female sex, 2) race/ethnicity other than non-Hispanic White, 3) less education, and 4) higher BMI, higher fasting insulin levels, and lower levels of physical activity. Elevated BDI scores were not related to age at baseline, but during the DPP older participants (≥ 60 years old) were less likely to have elevated BDI scores than younger participants.

Antidepressant use on entry to the study and use during the study (Table 1) was associated with female sex, non-Hispanic White race/ethnicity, and (on entry to the study only) more education. In contrast to elevated depression symptoms, antidepressant use was not associated with fasting insulin levels or physical activity level on entry to the DPP. Antidepressant use was associated with BMI on entry to and during the study, and with lower levels of physical activity during the study. Antidepressant use at baseline was not related to age, but during the DPP older participants (≥ 60 years old) were less likely to take these agents than younger participants.

At baseline, 7.4% of women, but only 2.0% (21 of 1029) of men, took antidepressants. The associations between antidepressant use and age, race/ethnicity, and education were similar for male and female participants, but when analyzed separately these associations were no longer significant among men, possibly due to lack of statistical power. Higher BMI at randomization was significantly associated with antidepressant

use at baseline and during the study for women only.

When other factors associated with an increased risk of developing diabetes (age, sex, fasting glucose, baseline weight, and weight change during the study) as well as race/ethnicity were controlled, neither elevated BDI scores at baseline nor during the study were significantly associated with diabetes risk in any treatment arm (see lines labeled HRa in Table 2). Leisure activity was not controlled because it was not associated with diabetes risk. (Absolute BDI scores at baseline were significantly associated with diabetes risk in the ILS arm only [$p=0.04$]).

Baseline antidepressant use, on the other hand, was strongly associated with diabetes risk for participants in the PLB [HR= 2.25 (95% CI: 1.38-3.66) and ILS [HR= 3.48 (95% CI: 1.93-6.28)] arms. Continuous antidepressant use during the study (compared with no use) was also significantly associated with diabetes risk in the same arms [PLB HR= 2.60 (95% CI: 1.37-4.94), ILS HR= 3.39 (95% CI: 1.61-7.13)], as was intermittent antidepressant use during the study in the ILS arm [HR= 2.07 (95% CI: 1.18-3.62)]. In the PLB arm the association between intermittent antidepressant uses during the study and diabetes risk approached significance [HR= 1.50 (95% CI: 0.97-2.33)]. In the MET arm, antidepressant use was not associated with diabetes risk. There was a significant difference between the ILS and MET arms in the association between antidepressant medication and diabetes risk.

We further analyzed baseline antidepressant use, comparing only participants who were taking SSRI, selective serotonin and norepinephrine reuptake inhibitors (SSNRI), or serotonin modulators (SM) with those who did not take any antidepressant, since these agents were considered less likely to increase the risk of diabetes. The results for this subgroup, which

included 78% of all those taking antidepressants at baseline, were similar to those for all antidepressant users: taking SSRI or related agents at baseline was significantly associated with the risk of developing diabetes during the DPP in the PLB and ILS arms, but not in the MET arm. For this reason, our analysis of the association between antidepressant use during the DPP and diabetes risk included all antidepressants.

Our findings suggest that antidepressant use might have increased diabetes risk in the PLB and ILS arms. Under this assumption, we calculated that the treating 5.4 PLB participants (95% CI: 3.1-24.8) or 5.2 ILS participants (95% CI: 3.2-15.5) with antidepressants would have caused one additional case of diabetes three years later (31). We did not perform this calculation for the MET arm, in which it would not be meaningful because there was no association between antidepressant use and diabetes risk.

CONCLUSIONS

Elevated depression scores were not associated with diabetes risk. Elevated BDI scores (≥ 11 , consistent with mild depression) at baseline and during the study were not associated with diabetes risk during the DPP, despite the fact that elevated depression symptoms *were* associated with several risk factors for diabetes (BMI, fasting glucose, and lower levels of physical activity). Absolute BDI scores were weakly associated with diabetes risk in the ILS arm only. These findings differ from the conclusions of a meta-analysis which suggested that elevated depressive symptoms are associated with the risk of developing type 2 diabetes (6). DPP participants were selected because their risk for developing diabetes was high, and rates of developing diabetes in the DPP were 5-fold to 10-fold higher than those in earlier studies included in the meta-analysis where subjects did not have impaired glucose regulation at baseline. In addition, the DPP cohort was

relatively free of depression; at baseline only 2.7% of participants had BDI scores indicating even moderate levels of depression (>16). Perhaps in a population at high risk of developing diabetes, generally mild depression symptoms were not potent enough to significantly affect overall diabetes rates. In some studies increased diabetes risk was seen only in those with more severe depression symptoms (6).

Antidepressant medication was associated with diabetes risk. The present study of individuals at very high risk for developing diabetes, with excellent ascertainment of diabetes onset and exposure to antidepressant medication, found an increased risk of diabetes associated with antidepressant medication. Two recent studies also assessed antidepressant-related diabetes risk. One study found no increased risk of diabetes among subjects taking antidepressant medication (7), but the study did not include a definitive determination of diabetes. The other study (32) compared the risk of developing diabetes for patients taking different antidepressants, and reported that the risk associated with taking SSRI in addition to TCA was greater than the risk associated with taking TCA alone. Since that study did not include those not taking antidepressants, the authors could not assess the diabetes-related risk associated with taking any antidepressant.

Among psychotropic agents, only second generation atypical antipsychotic agents (SGA) have been associated with increased diabetes risk (33). Weight gain – a known risk factor for diabetes – is a recognized side effect of tricyclic antidepressant (TCA) use, and TCA-associated hyperglycemia has been reported in patients who already have diabetes (9). SSRI or related agents, taken by 78% of DPP participants who took antidepressants, are generally considered to have less effect on weight, with some reports that in people with type 2 diabetes they can actually contribute to lowering weight (34) and improving insulin

sensitivity (10). In contrast, SSRI use in the DPP was associated with weight gain during the study; use of antidepressants was still associated with diabetes risk in the PLB and ILS arms when weight gain was controlled.

Explaining the association between antidepressant use and diabetes risk. The association between antidepressant use and diabetes risk remained significant when likely mediators (e.g., fasting glucose, weight, and weight gain) of this association were controlled. Thus we are unable to say how use of antidepressants increases diabetes risk, if it does at all. Antidepressant use could simply be a marker for the actual cause(s) of increased diabetes risk, perhaps more severe, chronic, or recurrent depression. Perhaps antidepressant users were more severely depressed in the past, or had a history of chronic or recurrent depression. If so, the lingering effects of past depression, or the current effects of that depression, even with its symptoms controlled by medication, could explain the association between antidepressant use and diabetes risk. This argument depends on the presence of past moderate or severe depression (or a history of chronic or recurrent depression), and on an effect of depression that operates after symptoms have resolved (since these participants did not have elevated depression symptoms), or on an unidentified continuing effect of depression not detected by our questionnaire.

Antidepressant use was not associated with diabetes risk in the MET arm. Antidepressant use was not associated with diabetes risk in the MET arm. Although there is no obvious explanation for this finding, it is similar to our previous report that metformin treatment not only reduced the incidence of diabetes but also obliterated the predictive effect of BMI (14).

Strengths and Limitations. Strengths of the current study include the large, racially and ethnically diverse population, the definitive

assessment of glucose tolerance and diabetes, and the fact that data were collected about antidepressant medication and depression symptoms. We were also able to pinpoint the diagnosis of diabetes, a considerable advance over studies that rely on clinical records which may not accurately capture when diabetes actually developed.

The absence of data on specific antidepressants is a limitation, but use of the most commonly used class of agents (SSRI) was associated with diabetes risk in the PLB and ILS study arms, just as use of any antidepressant was. Excluding participants from the DPP if they were taking bupropion, or SSRI in greater than low doses might limit the generalizability of our findings, but antidepressant medication and dosage were not limited during the study, where we still found an association between antidepressant use and diabetes risk.

We did not confirm that all patients using antidepressants were taking them for depression. Some SSRI and related agents are used to treat other disorders (35), but patients with these disorders probably represent a small proportion of those taking SSRI. TCA antidepressants are sometimes prescribed for relief of neuropathic symptoms in patients with diabetes, but DPP participants did not have diabetes at the outset of the study. The antidepressant bupropion is also indicated for smoking cessation, but only 6% of DPP participants reported taking this agent.

Our categorical classification of antidepressant use, based on patient self-report, precludes an examination of the association between incremental antidepressant dosage and diabetes risk. We also could not determine whether chronicity of depressive symptoms or the number of past episodes of depression were related to diabetes risk in this cohort, or modulated the noted association with antidepressant use.

Implications. Research implications. The finding of an association between

antidepressant use and diabetes risk requires replication. Ideally, future studies will examine the diabetes risk associated with different classes of antidepressant medication, and even specific agents. Such studies will require large populations, e.g., prescription databases, to obtain the necessary sample sizes. If the association between antidepressant use and diabetes risk is replicated, studies are needed to determine whether use of these agents is a causal risk factor for developing diabetes or a marker of depression severity. Studies should seek to clarify the metabolic effects of various antidepressants to determine how each agent affects diabetes risk. Comparing the diabetes risk associated with antidepressant use to the risk associated with psychological treatment could also be illuminating.

Clinical implications. If antidepressants prove to be an independent diabetes risk factor, clinicians will need to consider this when prescribing depression treatment in patients at high risk for diabetes. One possibility is to consider psychological treatment to avoid potential iatrogenic effects of antidepressants (although limited resources may often make this infeasible). In one study of depressed patients with diabetes who had high A1c levels, cognitive behavioral therapy counseling was associated with improved glycemic control (36); this suggests a potential benefit for patients at high risk for developing diabetes.

Public health implications. According to recent estimates there are over 40 million people in the U.S. with “prediabetes” – IGT or impaired fasting glucose (37) – and this number is rising rapidly. More than 10% of DPP participants were taking antidepressants when the study ended in 2001. Antidepressant use in the U.S. has continued to rise, so probably at least four million people in the U.S. who have prediabetes are taking antidepressants.

If future research confirms an etiologic role for antidepressants in diabetes, efforts to minimize the potentially negative effects of these agents on glycemic control should be pursued.

ACKNOWLEDGEMENTS

The Investigators gratefully acknowledge the commitment and dedication of the participants of the DPP. Support to the clinical centers and the Coordinating Center was provided by the National Institute of Diabetes and Digestive and Kidney Diseases through a Cooperative Agreement. The

Southwestern American Indian Centers were supported directly by the NIDDK, including its Intramural Research Program, and the Indian Health Service. The General Clinical Research Center Program, National Center for Research Resources supported many of the clinical centers. Funding was also provided by the Office of Research on Minority Health, the National Institute of Child Health and Human Development, the National Institute on Aging, the Centers for Disease Control and Prevention, the American Diabetes Association, Bristol-Myers Squibb and Parke-Davis.

REFERENCES

1. Egede LE, Zheng D: Independent factors associated with major depressive disorder in a national sample. *Diabetes Care* 26:104-111, 2003
2. de Groot M, Anderson R, Freedland KE, Clouse RE, Lustman PJ: Association of depression and diabetes complications: a meta-analysis. *Psychosom Med* 63:619-630, 2001
3. Katon WJ, Rutter C, Simon G, Lin EHB, Ludman E, Chiechanowski P, Kinder L, Young B, Von Korff M: The association of comorbid depression with mortality in patients with type 2 diabetes. *Diabetes Care* 28:2668-2672, 2005
4. Egede LE, Zheng D, Simpson K: Comorbid depression is associated with increased health care use and expenditures in individuals with diabetes. *Diabetes Care* 25:464-470, 2002
5. Willis T. *Diabetes: A Medical Odyssey*. New York, Tuckahoe, 1971
6. Knol MJ, Twisk JWR, Beekman ATF, Heine RJ, Snoek FJ, Pouwer F: Depression as a risk factor for the onset of type 2 diabetes. A meta-analysis. *Diabetologia* 49:837-845, 2006
7. Knol MJ, Geerlings MI, Egberts ACG, Gorter KJ, Grobbee DE, Heerdink ER: No increased incidence of diabetes in antidepressant users. *Int Clin Psychopharmacol* 22:382-386, 2007
8. Aronne LJ, Segal KR: Weight gain in the treatment of mood disorders. *J Clin Psychiatry* 64:22-29, 2003
9. Lustman PJ, Griffith LS, Clouse RE, Freedland KE, Eisen SA, Rubin EH, Carney RM, McGill JB: Effects of nortriptyline on depression and glycemic control in diabetes: Results of a double-blind, placebo-controlled trial. *Psychosom Med* 59: 241-250,1997
10. Maheux P, Ducros F, Bourque J, Garon J, Chiasson J-L : Fluoxetine improves insulin sensitivity in obese patients with non-insulin-dependent diabetes mellitus independent of weight loss. *Int J Obes Relat Metab Disord* 21:97-102,1997
11. The Diabetes Prevention Program Research Group: Depression symptoms and antidepressant medicine use in Diabetes Prevention Program participants. *Diabetes Care* 28:830-837,2005
12. The Diabetes Prevention Program Research Group: The Diabetes Prevention Program: design and methods for a clinical trial in the prevention of type 2 diabetes. *Diabetes Care* 22:623-634,1999
13. The Diabetes Prevention Program Research Group: The Diabetes Prevention Program: recruitment methods and results. *Controlled Clinical Trials* 23:157-171, 2002
14. The Diabetes Prevention Program Research Group: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *NEJM* 346:393-403, 2002
15. American Psychiatric Association: *American Psychiatric Association Practice Guideline for the Treatment of Patients with Major Depressive Disorder*. 2nd ed. Washington, DC, American Psychiatric Association, 2000, p. 424
16. The Diabetes Prevention Program (DPP) Research Group: The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care* 25:623-634, 2002
17. Beck AT, Steers RA: *Manual of the Beck Depression Inventory*. San Antonio, TX, Psychological Corporation, 1993
18. Kriska AM, Caspersen CJ: Introduction to a collection of physical activity questionnaires. *Med Sci Sports Exerc* 29:Suppl:S5-S9, 1997
19. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 20:1183-1197,1997

20. The Diabetes Prevention Program Research Group: The Diabetes Prevention Program: baseline characteristics of the randomized cohort. *Diabetes Care* 23:1619–1629, 2000
21. Leedom I, Meehan WT, Procci W, Zeidler: Symptoms of depression in patients with type II diabetes mellitus. *Psychosomatics* 32:280-286, 1991
22. Palinkas LA, Barrett-Connor E, Wingard DL: Type 2 diabetes and depressive symptoms in older adults: a population-based study. *Diabet Med* 8:532-539, 1991
23. Wing RR, Marcus MD, Blair EH, Epstein LH, Burton LR: Depressive symptomatology in obese adults with type II diabetes. *Diabetes Care* 13:170-172, 1990
24. Wilcoxon F: Individual comparisons by ranking methods. *Biometrics Bulletin* 1:80-83, 1945
25. Lindsey JK: *Models for repeated measurements*. Oxford, Clarendon Press, 1993
26. Liang KY, Zeger SL: Longitudinal data analysis using generalized linear models. *Biometrika* 73:13-22, 1986
27. Cox, DR: Partial likelihood. *Biometrika* 62, 269 -276, 1975
28. Allison, PD: *Survival Analysis Using the SAS System: A Practical Guide*. Cary, NC, SAS Institute Inc., 1995
29. Lachin JM: *Biostatistical methods: the assessment of relative risk*. New York, John Wiley, 2000
30. Greenwood M: *The natural duration of cancer*. *Reports on public health and medical subjects* (33:1-26), 1926
31. Hutton JL: Number needed to treat: properties and problems. *Journal of the Royal Statistical Society. Series A (Statistics in Society)* 163:403-19, 2000
32. Brown LC, Majumdar SR, Johnson JA: Type of antidepressant therapy and risk of type 2 diabetes in people with depression. *Diab Res Clin Pract* doi:1016/j.diabres.2007.07.009
33. Cohen D, Stolk RP, Grobbee DE, Gispen-De Wied CC: Hyperglycemia and diabetes in patients with schizophrenia or schizoaffective disorders. *Diabetes Care* 29:786-791, 2006
34. Sussman N, Ginsberg DL, Bikoff J: Effects of nefazodone on body weight: a pooled analysis of selective serotonin reuptake inhibitor- and imipramine-controlled trials. *J Clin Psychiatry* 62:256-260, 2001
35. Thompson PDR: *Physicians Desk Reference*. 18th ed. Montvale, NJ, 2004
36. Lustman PJ, Griffith LS, Freedland KE, Kissel SS, Clouse RE: Cognitive behavior therapy for depression in type 2 diabetes: A randomized controlled trial. *Ann Intern Med* 129:613-621, 1998
37. Centers for Disease Control and Prevention: *Number of Americans with diabetes continues to increase*. Fact sheet released October 26, 2005

TABLE 1. Depression Symptoms and Antidepressant Use at Baseline and during DPP

Characteristic	At baseline					During DPP					
	Total N	BDI ≥ 11 N(%)		Antidepressant medicine use N(%)		Total person-years	BDI ≥ 11 Person-years and % of total person-years		Intermittent use Person-years and % of total person-years		Continuous use Person-years and % of total person-years
Total Participants	3187	328 (10.3%)		181 (5.7%)		8637.1	1302.1 (15.1%)		622.0 (7.2%)		272.4 (3.2%)
Age (years)							*				
25-44	983	115 (11.7%)		47 (4.8%)		2628.2	460.3 (17.5%)		223.4 (8.5%)		53.7 (2.0%)
45-59	1567	160 (10.2%)		102 (6.5%)		4267.2	639.6 (15.0%)		318.6 (7.5%)		151.7 (3.6%)
≥ 60	637	53 (8.3%)		32 (5.0%)		1741.7	202.1 (11.6%)		80.0 (4.6%)		66.9 (3.8%)
Sex		***		***			***		***		
Female	2158	255 (11.8%)		160 (7.4%)		5854.1	1011.2 (17.3%)		520.0 (8.9%)		240.4 (4.1%)
Male	1029	73 (7.1%)		21 (2.0%)		2783.0	290.9 (10.5%)		102.0 (3.7%)		32.0 (1.1%)
Race/Ethnicity		***		***			***		***		
White	1746	132 (7.6%)		144 (8.2%)		4736.0	578.6 (12.2%)		436.2 (9.2%)		238.0 (5.0%)
African American	636	81 (12.7%)		11 (1.7%)		1690.0	299.0 (17.7%)		66.8 (4.0%)		9.2 (0.5%)
Hispanic	498	67 (13.5%)		18 (3.6%)		1343.8	244.3 (18.2%)		75.4 (5.6%)		21.0 (1.6%)
American Indian	165	33 (20.0%)		6 (3.6%)		460.4	125.4 (27.2%)		32.4 (7.0%)		1.3 (0.3%)
Asian	142	15 (10.6%)		2 (1.4%)		406.8	54.8 (13.5%)		11.1 (2.7%)		3.0 (0.7%)
Education in Years		***		*			***		*		
≤ 12	822	122 (14.8%)		33 (4.0%)		2249.2	453.2 (20.1%)		137.8 (6.1%)		54.5 (2.4%)
13-16	1534	152 (9.9%)		91 (5.9%)		4176.7	615.2 (14.7%)		302.7 (7.2%)		129.8 (3.1%)
≥ 17	831	54 (6.5%)		57 (6.9%)		2211.2	233.7 (10.6%)		181.5 (8.2%)		88.1 (4.0%)
		BDI ≥ 11		Antidepressant use			BDI ≥ 11		Antidepressant use		
		No	Yes	No	Yes		No	Yes	Never exposed	Intermittent use	Continuous use
Weight (kg)	Mean (SD)	94.1 (20.1)	95.6 (21.3)	94.0 (20.2)	97.8 (20.9) *	LS Mean (SE)	93.2 (0.4)	93.8 (0.4) *	93.3 (0.4)	93.1 (0.4)	94.7 (0.6) *
BMI (kg/m ²)	Mean (SD)	33.8 (6.6)	35.5 (7.0) ***	33.9 (6.7)	35.8 (7.1) **	LS Mean (SE)	33.6 (0.1)	33.9 (0.1) *	33.6 (0.1)	33.6 (0.1)	34.2 (0.2) **
Fasting Glucose (mg/dl)	Mean (SD)	106.5 (8.2)	106.6 (8.9)	106.6 (8.3)	105.3 (8.2) *	LS Mean (SE)	105.7 (0.1)	106.2 (0.3)	105.8 (0.1)	105.8 (0.4)	105.6 (0.6)
Fasting Insulin (uU/ml)	Mean (SD)	26.6 (15.2)	28.5 (14.9) *	26.8 (15.3)	26.5 (13.9)	LS Mean (SE)	25.0 (0.2)	27.4 (0.5) ***	25.3 (0.2)	26.7 (0.7)	25.1 (0.9)
Leisure activity (Met-hours)	Mean (SD)	16.6 (26.1)	14.5 (25.0) ***	16.5 (26.2)	14.4 (22.4)	LS Mean (SE)	18.6 (0.3)	15.8 (0.7) **	18.4 (0.3)	15.7 (1.0)	15.0 (1.4) *

Significance levels of differences in depression symptoms or antidepressant use are noted as follows: * $p < 0.05$, ** $p < 0.001$, *** $p < 0.0001$. Thus *** noted for sex in column 'BDI \geq 11' (at baseline) indicates that the proportion of those with elevated baseline BDI scores differed by sex with $p < 0.001$. During DPP, variables age, sex, race/ethnicity and education remain the same. Variables weight, BMI, fasting glucose, fasting insulin and leisure activity are measured repeatedly during the study and repeated model based means are reported.

TABLE 2. Prediction of Diabetes by Depression Symptoms and Antidepressant Drug Exposure

		At baseline				During follow-up				
		BDI<11	BDI≥11	No meds	Taking meds	BDI always <11	BDI ever ≥11	No meds	Intermittent meds	Continuous meds
PLB	Pyr	2451.6	300.4	2624.9	127.1	2316.7	435.3	2460.0	218.8	73.2
	Cases	270	30	277	23	245	55	259	28	13
	IR	11.0	10.0	10.6	18.1	10.6	12.6	10.5	12.8	17.8
	HRu	1	0.90 (0.61-1.32)	1	1.80 (1.15-2.82)	1	1.11 (0.82-1.50)	1	1.17 (0.78-1.76)	1.80 (1.00-3.25)
	HRa	1	0.84 (0.56-1.26)	1	2.25 (1.38-3.66)	1	0.99 (0.72-1.37)	1	1.50 (0.97-2.33)	2.60 (1.37-4.94)
MET	Pyr	2584.9	289.0	2675.7	198.2	2441.0	432.9	2549.2	224.4	100.3
	Cases	199	25	213	11	187	37	200	19	5
	IR	7.7	8.7	8.0	5.5	7.7	8.5	7.8	8.5	5.0
	HRu	1	1.14 (0.74-1.75)	1	0.69 (0.37-1.27)	1	1.00 (0.69-1.43)	1	0.96 (0.59-1.57)	0.68 (0.28-1.68)
	HRa	1	1.07 (0.68-1.68)	1	0.72 (0.38-1.36)	1	0.91 (0.62-1.33)	1	0.99 (0.59-1.65)	0.71 (0.28-1.78)
ILS	Pyr	2711.2	299.9	2863.5	147.6	2577.2	433.9	2733.4	178.7	99.0
	Cases	130	21	134	17	120	31	123	19	9
	IR	4.8	7.0	4.7	11.5	4.7	7.1	4.5	10.6	9.1
	HRu	1	1.49 (0.93-2.39)	1	2.68 (1.59-4.52)	1	1.35 (0.90-2.02)	1	1.91 (1.15-3.19)	2.51 (1.29-4.89)
	HRa	1	1.15 (0.69-1.93)	1	3.48 (1.93-6.28)	1	0.87 (0.56-1.39)	1	2.07 (1.18-3.62)	3.39 (1.61-7.13)

Abbreviations:

BDI = Beck depression inventory score

Meds = antidepressant drugs

Pyr = person-years of follow-up

Cases = incident cases of diabetes developing during follow-up

IR = incidence rate (cases/100 Pyr)

HRu = Hazard ratio (from Cox proportional hazard model) with 95% confidence interval, unadjusted for covariates

HRa = Hazard ratio (from Cox proportional hazard model) with 95% confidence interval, adjusted for age, sex, race/ethnicity, education, fasting plasma glucose, weight, and weight change. In addition, depression symptom and anti-depressant drug use are adjusted for each other.