

International tables of glycemic index and glycemic load values: 2008

Fiona S. Atkinson (RD), Kaye Foster-Powell (RD), Jennie C. Brand-Miller (PhD)

Institute of Obesity, Nutrition and Exercise
University of Sydney, NSW, 2006, Australia

Correspondence:

Professor J. Brand-Miller

Email: j.brandmiller@biochem.usyd.edu.au

Additional information for this article can be found in an online appendix at
<http://care.diabetesjournals.org>.

Submitted 8 July 2008 and accepted 13 September 2008.

This is an uncopyedited electronic version of an article accepted for publication in *Diabetes Care*. The American Diabetes Association, publisher of *Diabetes Care*, is not responsible for any errors or omissions in this version of the manuscript or any version derived from it by third parties. The definitive publisher-authenticated version will be available in a future issue of *Diabetes Care* in print and online at <http://care.diabetesjournals.org>.

Objective: To systematically tabulate published and unpublished sources of reliable GI values.

Research Design and Methods: A literature search identified 205 articles published between 1981 and 2007. Unpublished data were also included where the data quality could be verified. The data were separated into 2 lists, the first representing more precise data derived from testing healthy subjects, and the second primarily from individuals with impaired glucose metabolism.

Results: The tables list the GI of over 2,480 individual food items. Dairy products, legumes and fruits were found to be low GI. Breads, breakfast cereals and rice, including wholegrain, were available in both high and low GI versions. The correlation coefficient for 20 staple foods tested in both healthy and diabetic subjects was $r = 0.94$ ($p < 0.001$).

Conclusions: These tables improve the quality and quantity of GI data available for research and clinical practice.

The relevance of dietary glycemic index (GI) and glycemic load (GL) is debated. While the World Health Organisation (1), the American Diabetes Association (2), Diabetes UK (3) and the Canadian Diabetes Association (4) give qualified support for the concept, many health professionals still consider it complex and too variable for clinical practice (5). The availability of reliable tables of GI is critical for continuing research and resolution of the controversy. New data have become available since previous tables were published in 2002 (6). Our aim was to systematically tabulate published and unpublished sources of reliable GI values, with derivation of the GL.

RESEARCH DESIGN AND METHODS

We conducted a literature search of MEDLINE from January 1981 through December 2007 using the terms 'glyc(a)emic index' and 'glyc(a)emic load'. We restricted the search to human studies published in English using standardised methodology. We performed a manual search of relevant citations and contacted experts in the field. Unpublished values from our laboratory and elsewhere were included. Values listed in previous tables (6, 7) were not automatically entered but reviewed first. Final data were divided into two lists. Values derived from groups of 8 or more healthy subjects were included in the first list. Data derived from testing individuals with diabetes or impaired glucose metabolism, too few subjects ($n \leq 5$) or showing wide variability ($SEM > 15$), were included in the second list. Some foods were tested in only 6 or 7 normal subjects but otherwise appeared reliable and were included in the first list. Two columns of GI values were created because both glucose and white bread continue to be used reference foods. The conversion factor 100/70 or 70/100 was used to convert from one scale to the other.

In instances whether other reference foods (eg rice) were used, this was accepted provided the conversion factor to the glucose scale had been established. To avoid confusion, the glucose scale is recommended for final reporting. GL values were calculated as the product of the amount of available carbohydrate in a specified serving size and the GI value (using glucose as the reference food), divided by 100. Carbohydrate content was obtained from the reference paper or food composition tables (8). The relationship between GI values determined in normal subjects versus diabetic subjects was tested by linear regression. Common foods ($n = 20$), including white bread, cornflakes, rice, oranges, corn, apple juice, sucrose and milk were used for this analysis.

RESULTS

Tables A1 and A2 list 2487 separate entries, citing 205 separate studies (Table A1 and Table A2 available online at <http://care.diabetesjournals.org>). Table A1, representing reliable data derived from subjects with normal glucose tolerance, contains 1879 individual entries (75% of the total). Table A2 contains 608 entries, of which 491 values were determined in individuals with diabetes or impaired glucose metabolism (20% of the total). The correlation coefficient for 20 foods tested in both normal and diabetic subjects was $r = 0.94$ ($p < 0.001$, line of best fit: $y = 0.9x + 9.7$ where x is the value in normal subjects). Table A2 also lists 60 values derived from groups of 5 or fewer subjects, and 57 values displaying wide variability ($SEM > 15$). A summary table comprising values for 62 common foods appears below. More reliable values are available for many foods, including carrots (GI = 39) and bananas (GI = 51).

CONCLUSIONS

The 2008 edition of tables of GI and GL has doubled the amount of data available for research and other applications. Most varieties of legumes, pasta, fruits and dairy products are still classified as low GI foods (55 or less on the glucose reference scale). Breads, breakfast cereals, rice and snack products, including wholegrain versions, are available in both high (70 or greater) and low GI forms. Most varieties of potato and rice are high GI, but lower GI cultivars were identified. Many confectionary items, such as chocolate, have a low GI but their high saturated fat content reduces their nutritional value. The GI should not be used in isolation, the energy density and macronutrient profile of foods should also be considered (1). The high correlation coefficient ($r = 0.94$) between values derived from testing the same foods in normal and diabetic subjects indicates that GI values in Table A1 are relevant to dietary interventions in people with diabetes.

Although data quality has been improved, many foods have been tested only once in 10 or fewer subjects, and caution is needed. Repeated testing of certain products indicates that white and wholemeal bread have remained remarkably consistent over the past 25 y, but other products appear to be increasing in GI. This secular change may arise because of efforts on the part of the food industry to make food preparation more convenient and faster cooking. Some foods, such as porridge oats, show variable results, which may reflect true differences in refining and processing that affect the degree of starch gelatinization (9). Users should note that manufacturers sometimes give the same product different names in different countries, and in some cases, the same name for different items. Kellogg's Special K™ and All-Bran™, for example, are different formulations in North America, Europe and Australia.

Assignment of GI values to foods requires knowledge of local foods. Ideally,

branded product information is available because manufacturers prepare and process foods, particularly cereal products, in different ways. This variability is not unique to the GI, but true of many nutrients, including saturated fat and fiber. In the absence of specific product GI information, these tables provide the basis for extrapolation. In the case of low carbohydrate products, a GI value of 40 for vegetables, 70 for flour products and 30 for dairy foods could be assigned.

In summary, the 2008 edition of the international tables of GI improves the quality and quantity of reliable data available for research and clinical practice. The data in Table A1 should be preferred for research and coding of food databases. The values listed in Table A2 may be helpful in the absence of other data.

ACKNOWLEDGEMENTS

JBM and KFP conceived the study. FSA and KFP searched the literature, evaluated the quality of the data and composed the tables. All authors contributed to the interpretation of the findings, and the writing of the final paper.

JBM and KFP are co-authors of *The New Glucose Revolution* book series (Hodder and Stoughton, London; Marlowe and Co, New York; Hodder Headline, Sydney and elsewhere). JBM is the Director of a not-for-profit GI-based food endorsement program in Australia. FSA is employed to manage the University of Sydney GI testing service.

REFERENCES

1. Mann J, Cummings J, Englyst H, Key T, Liu S, Riccardi G, Summerbell C, Uauy R, van Dam R, Venn B, Vorster H, Wiseman M. *FAO/WHO Scientific Update on carbohydrates in human nutrition: conclusions*. Eur J Clin Nutr, 2007. 61: S132-7.
2. Sheard N, Clark N, Brand-Miller J, Franz M, Pi-Sunyer FX, Mayer-Davis E, Kulkarni K, Geil P. *Dietary carbohydrate (amount and type) in the prevention and management of diabetes*. Diabetes Care, 2004. 27: 2266-71.
3. Nutrition Subcommittee of the Diabetes Care Advisory Committee of Diabetes UK. *The implementation of nutritional advice for people with diabetes*. Diabet Med, 2003. 20: 786-807.
4. Canadian Diabetes Association. *Guidelines for the nutritional management of diabetes mellitus in the new millennium. A position statement by the Canadian Diabetes Association*. Can J Diabetes Care, 2000. 23: 56-69.
5. Franz M. *The glycemic index: not the most effective nutrition therapy intervention*. Diabetes Care, 2003. 26: 2466-8.
6. Foster-Powell K, Holt SH, Brand-Miller JC. *International table of glycemic index and glycemic load values: 2002*. Am J Clin Nutr, 2002. 76(1): 5-56.
7. Foster-Powell K, Miller J. *International tables of glycemic index*. Am J Clin Nutr, 1995. 62(4): 871S-90S.
8. US Department of Agriculture, Agricultural Research Service, 2007. *USDA National Nutrient Database for Standard Reference*. Release 20. Nutrient Data laboratory Home Page, <http://www.ars.gov/ba/bhnrc/ndl>. Accessed 20 May 2008.
9. Bjorck I, Granfeldt Y, Liljeberg H, Tovar J, Asp N-G. *Food properties affecting the digestion and absorption of carbohydrates*. Am J Clin Nutr, 1994. 59: 699S-705S.

Summary Table: The average glycemic index (GI, mean \pm SEM) of 62 common foods derived from multiple studies by different laboratories.

High carbohydrate foods		Breakfast cereals		Fruit & fruit products	
White wheat bread ¹	75 \pm 2	Cornflakes	81 \pm 6	Apple, raw ²	36 \pm 2
Wholewheat/wholemeal bread	74 \pm 2	Wheat flake biscuits	69 \pm 2	Orange, raw ²	43 \pm 3
Specialty grain bread	53 \pm 2	Porridge, rolled oats	55 \pm 2	Banana, raw ²	51 \pm 3
Unleavened wheat bread	70 \pm 5	Instant oat porridge	79 \pm 3	Pineapple, raw	59 \pm 8
Wheat roti	62 \pm 3	Rice porridge/congee	78 \pm 9	Mango, raw ²	51 \pm 5
Chapati	52 \pm 4	Millet porridge	67 \pm 5	Watermelon, raw	76 \pm 4
Corn tortilla	46 \pm 4	Muesli	57 \pm 2	Dates, raw	42 \pm 4
White rice, boiled ¹	73 \pm 4			Peaches, canned ²	43 \pm 5
Brown rice, boiled	68 \pm 4	Vegetables		Strawberry jam/jelly	49 \pm 3
Barley	28 \pm 2	Potato, boiled	78 \pm 4	Apple juice	41 \pm 2
Sweet corn	52 \pm 5	Potato, instant mash	87 \pm 3	Orange juice	50 \pm 2
Spaghetti, white	49 \pm 2	Potato, French Fries	63 \pm 5		
Spaghetti, wholemeal	48 \pm 5	Carrots, boiled	39 \pm 4	Sugars	
Rice noodles ²	53 \pm 7	Sweet potato, boiled	63 \pm 6	Fructose	15 \pm 4
Udon noodles	55 \pm 7	Pumpkin, boiled	64 \pm 7	Sucrose	65 \pm 4
Couscous ²	65 \pm 4	Plantain/Green banana	55 \pm 6	Glucose	103 \pm 3
		Taro, boiled	53 \pm 2	Honey	61 \pm 3
Dairy products & alternatives		Vegetable soup	48 \pm 5		
Milk, full fat	39 \pm 3			Snack products	
Milk, skim	37 \pm 4	Legumes		Chocolate	40 \pm 3
Ice cream	51 \pm 3	Chickpeas	28 \pm 9	Popcorn	65 \pm 5
Yoghurt, fruit	41 \pm 2	Kidney beans	24 \pm 4	Potato crisps	56 \pm 3
Soy milk	34 \pm 4	Lentils	32 \pm 5	Soft drink/Soda	59 \pm 3
Rice milk	86 \pm 7	Soya beans	16 \pm 1	Rice crackers/crisps	87 \pm 2

Footnotes

1. Low GI varieties were also identified
2. Average of all available data