

Diabetic foot prevention: a neglected opportunity in high risk patients

Lawrence A. Lavery, DPM¹; Nathan A. Hunt, DPM¹; Javier LaFontaine, DPM^{1,2}; Cory L. Baxter, DPM¹; Agbor Ndip, MD³; Andrew J.M. Boulton, MD³

1. Department of Surgery, Scott and White Hospital and Texas A & M Health Science Center College of Medicine, Temple, Texas
2. Podiatry Section, Central Texas Veterans Healthcare System, Temple, Texas
3. Department of Diabetes and Medicine, Manchester Royal Infirmary, Manchester, UK

Address Correspondence to:

Nathan A. Hunt, DPM
Email: nhunt@swmail.sw.org

Submitted 17 February 2010 and accepted 19 April 2010.

This is an uncopyedited electronic version of an article accepted for publication in Diabetes Care. The American Diabetes Association, publisher of Diabetes Care, is not responsible for any errors or omissions in this version of the manuscript or any version derived from it by third parties. The definitive publisher-authenticated version will be available in a future issue of Diabetes Care in print and online at <http://care.diabetesjournals.org>.

Objective: To evaluate the frequency of foot prevention strategies among high risk patients with diabetes.

Research design and methods: Electronic medical records were used to identify 150 patients on dialysis and 150 patients with previous foot ulceration or amputation with 30 months follow-up to determine the frequency patients received education, podiatry care, and therapeutic shoes and insoles as prevention services.

Results: Few patients had formal education (1.3%), therapeutic shoes/insoles (7%), or preventative podiatric care (30%). The cumulative ulcer incidence was the same in both groups (210 per thousand person-years). In contrast, the cumulative amputation incidence was higher in the dialysis group compared to the ulcer group (58.7 vs. 13.1 per thousand person-years, $p < 0.001$). Patients on dialysis were younger and more likely to be of non-Hispanic white descent ($p = 0.006$) than patients with a previous history of ulcer or amputation.

Conclusions: Prevention services are infrequently provided to high risk patients.

The prevalence of foot complications is 250% higher among dialysis-treated patients when compared to patients without chronic kidney disease (1,2,3). Similarly, patients with a past ulcer history have a 34 times greater risk of developing another ulcer (4,5). Programs to prevent foot ulcers and amputations generally involve therapeutic shoes and insoles, regular foot care, and patient education (6,7,8). This study evaluated the frequency of prevention services among high risk patients.

RESEARCH DESIGN AND METHODS:

We used claims data for diabetes (ICD-9 250.X), ulceration (ICD-9 707.10, 707.14 and 707.15) and dialysis (CPT 90935-90937) from the Scott and White Health Plan to identify 150 consecutive patients in each group with at least 30 months follow-up from the time of diagnosis. We enrolled subjects from 2000 to 2006. We verified these diagnoses by reviewing comprehensive electronic medical records (EMR): including all patient care notes, imaging, labs, and prescriptions. Scott and White is an integrated, multi-specialty physician group with approximately 550 physicians and 14 clinics, 3 dialysis centers and a 535 bed hospital.

For the ulcer group, our evaluation began after the initial ulcer healed. For the dialysis group, our evaluation began with the initiation of dialysis. Subjects with HIV/AIDS, trauma from motor vehicle accidents, bilateral amputations, and patients with less than 30 months follow-up were excluded.

Three prevention therapies were evaluated: pedorthic care (professionally fitted therapeutic shoes and insoles), diabetes education, and podiatry services. Pedorthic services were identified from notes in the EMR and durable medical equipment codes (Codes A5501, A5503-A5508, A5512, A5513). Diabetes education was defined as a

session with a certified diabetes educator (CPT code S9445, S9460, S9465). Our diabetes education program addressed “the diabetic foot” in the third of four education sessions. Podiatry care was assessed by review of the EMR to identify the number of visits, and determine whether the visit was for prevention, ulcer treatment, or other pathology. Foot assessment by any health care provider was also identified.

Peripheral vascular disease was defined as at least two non-palpable foot pulses or abnormal ankle-brachial indices (<0.9). Neuropathy was defined as at least one site insensate to a 10 gram Semmes-Weinstein monofilament, abnormal vibration perception (>25 volts), or abnormal light touch sensation. Pearson’s Chi-square and Fisher’s exact tests were used to compare categorical data between study groups. Student’s T-test was used to compare continuous data.

RESULTS

We studied 300 patients (Dialysis n=150, Ulcer group n=150), 92.3% had type 2 diabetes (Table). Compared to the ulcer group, dialysis patients were 10 years younger on average and less likely to be of Hispanic (p=0.006) or African descent (p<0.001).

The incidence of ulceration and amputation was high in both study groups.–Incidence of ulceration was 210 per thousand person-years in both groups. However, amputation incidence was significantly higher in the dialysis group (58.7 vs. 13.1 per thousand person-years, p<0.001).

Few patients received prevention services (Table). Two patients (1.3%) in the dialysis group had formal diabetes education, neither attended the diabetic foot care session. No one in the ulcer group received formal education. A small proportion of patients received therapeutic shoes. During the first 12-month evaluation period 21 patients (7%) received shoes and insoles. Only 4 patients

(1.3%) received a second pair of therapeutic shoes and insoles during the second 12-month study period and no one received a third pair of shoes in the final six months. There was no difference in the proportion of patients that received therapeutic shoes between the dialysis and ulcer groups (7.3% vs. 6.7%, $p=1.0$).

During the 30 month evaluation period, 195 patients (65%) received care by a podiatrist. However, the majority of patients (70%) were seen after they developed a foot ulcer. Only 90 patients (30%) were seen for preventative care prior to ulceration. Significantly fewer patients in the ulcer group were seen by a podiatrist for preventative care (18%) compared to the dialysis group (42%, $p<0.001$). Additionally, neuropathy (35%) and vascular assessments (62.4%) were infrequently performed.

CONCLUSIONS

This study focused on two high risk groups for developing diabetic foot ulcers and amputations (1,9). As expected, the cumulative amputation incidence was high in both groups (ulcer group 13.1 and dialysis group 58.7 per thousand person-years). The amputation incidence in the general population with diabetes ranges from 4.4-9.5 per thousand person-years (10).

Prevention services were infrequently provided to patients in both risk groups. In our study, only 7% of patients received therapeutic shoes, 1.3% received professional education, and 30% received preventative care by a podiatrist. Other reports suggest a poor referral pattern for therapeutic shoes as well. In a study by Sugarman only 2.9% of subjects with diabetes that met the criteria for "high risk" received therapeutic footwear (11). Although the high rate of amputation may be due to our study patients' inherent risk for foot complications, it is possible that poorly utilized prevention services played a role. We expect that appropriate prevention

services could have significantly reduced the high rate of amputation.

We believe the results of this study can be generalized to high risk patients in other health care settings. Perhaps, prevention services would be provided less frequently in community practices that are not integrated and that do not have electronic medical records, because it is more difficult to communicate and coordinate care.

Specialized diabetic foot programs have been reported to reduce the incidence of amputations by 50% (6,7,8). Uccioli demonstrated ~50% reduction in foot ulceration when therapeutic shoes were prescribed for patients with an ulcer history compared to patients that selected their own shoes (12), and others have demonstrated that patients receiving regular foot care have fewer recurrent ulcers (13).

Prevention services for the diabetic foot are simple to establish and can be made easily accessible through organized multi-disciplinary care. This data provide further evidence that preventative foot care is not regularly provided, even among patients with the highest risk for lower limb complications. It also highlights an opportunity to improve prevention services for the diabetic foot with simple protocols for evaluation and referral.

Author Contributions: L.A.L. researched data, wrote the manuscript, and reviewed/edited the manuscript. N.A.H. contributed to the discussion, and reviewed/edited the manuscript. J.L. reviewed/edited the manuscript. C.L.B. researched data. A.N. reviewed/edited the manuscript. A.J.M.B. reviewed/edited the manuscript.

ACKNOWLEDGEMENTS

This study was supported by a research grant provided by the American Podiatric Medical Association.

REFERENCES

1. Ndip A, Lavery LA, Lafontaine J, Rutter MK, Vardhan A, Vileikyte L, Boulton AJ. High levels of foot ulceration and amputation risk in a multiracial cohort of diabetic patients on dialysis therapy. *Diabetes Care* 33:1-2, 2010
2. Dossa CD, Shepard AD, Amos AM, Kupin WL, Reddy DJ, Elliott JP, Wilczwski JM, Ernst CB. Results of lower extremity amputations in patients with end-stage renal disease. *J Vasc Surg* 20:14-9, 1994
3. Eggers PW, Gohdes D, Pugh J. Nontraumatic lower extremity amputations in the Medicare end-stage renal disease population. *Kidney Int* 56:1524-33, 1999
4. Peters EJ, Lavery LA. Effectiveness of the diabetic foot risk classification system of the International Working Group on the Diabetic Foot. *Diabetes Care* 24:1442-7, 2001
5. Lavery LA, Peters EJ, Williams JR, Murdoch DP, Hudson A, Lavery DC. Reevaluating the way we classify the diabetic foot: restructuring the diabetic foot risk classification system of the International Working Group on the Diabetic Foot. *Diabetes Care* 31:154-6, 2008
6. Foster AV, Snowden S, Grenfell A, Watkins PJ, Edmonds ME. Reduction of gangrene and amputations in diabetic renal transplant patients: the role of a special foot clinic. *Diabet Med* 12:632-5, 1995
7. Rith-Najarian S, Branchaud C, Beaulieu O, Gohdes D, Simonson G, Mazze R. Reducing lower-extremity amputations due to diabetes. Application of the staged diabetes management approach in a primary care setting. *J Fam Pract* 47:127-32, 1998
8. Lavery LA, Wunderlich RP, Tredwell JL. Disease management for the diabetic foot: effectiveness of a diabetic foot prevention program to reduce amputations and hospitalizations. *Diabetes Res Clin Pract* 70:31-7, 2005
9. Game FL, Chipchase SY, Hubbard R, Burden RP, Jeffcoate WJ. Temporal association between the incidence of foot ulceration and the start of dialysis in diabetes mellitus. *Nephrol Dial Transplant* 21:3207-10, 2006
10. Lavery LA, Ashry HR, van Houtum W, Pugh JA, Harkless LB, Basu S. Variation in the incidence and proportion of diabetes-related amputations in minorities. *Diabetes Care* 19:48-52, 1996
11. Sugarman JR, Reiber GE, Baumgardner G, Prella CM, Lowery J. Use of the therapeutic footwear benefit among diabetic medicare beneficiaries in three states, 1995. *Diabetes Care* 21:777-81, 1998
12. Uccioli L, Faglia E, Monticone G, Favales F, Durola L, Aldeghi A, Quarantiello A, Calia P, Menzinger G. Manufactured shoes in the prevention of diabetic foot ulcers. *Diabetes Care* 18:1376-8, 1995
13. Plank J, Haas W, Rakovac I, Gorzer E, Sommer R, Siebenhofer A, Pieber TR. Evaluation of the impact of chiropodist care in the secondary prevention of foot ulcerations in diabetic subjects. *Diabetes Care* 26:1691-5, 2003

Table: Patient demographics and results

	Dialysis (n=150)	Ulcer History (n=150)	Total (n=300)
Male (%)	40.7% (61)	41.3% (62)	41.0% (123)
Race*			
White	77.3% (116)	42.7% (64)	60.0% (180)
Black	13.3% (20)	36.0% (54)	24.7% (74)
Hispanic	7.3% (11)	18.6% (28)	13.0% (39)
Other	2.0% (3)	2.7% (4)	2.3% (7)
Age (Mean \pm SE)*	64.9 \pm 0.98	74.25 \pm 1.04	70.22 \pm 0.58
% Type 2 diabetes	92.6% (139)	92.0% (138)	92.3% (277)
% Amputation (n)*	14.7% (22)	3.3% (5)	9% (27)
% Ulceration (n)	52.7% (79)	52.7% (79)	52.6 % (158)
Neuropathy testing			
Semmes Weinstein Monofilament	26.7% (40)	27.3% (41)	27% (81)
Vibration Perception Threshold	2.7% (4)	2.0% (3)	2.3% (7)
Other assessment	3.3% (5)	8.0% (12)	5.7% (17)
No neuropathy testing	67.3% (101)	62.7% (94)	65% (195)
Vascular Assessment			
Pedal pulse evaluated	47.3% (71)	56.0% (84)	51.7% (155)
ABI evaluated	12.7% (19)	8.7% (13)	10.7% (32)
No assessment	40.0% (60)	35.3% (53)	37.6% (113)
Diabetes Education			
Session 1 (n)	0.6% (1)	0	0.3% (1)
Session 2 (n)	0.6% (1)	0	0.3% (1)
Sessions 3-4	0	0	0
Podiatry			
Podiatry- anytime*	49.3% (74)	80.6% (121)	65% (195)
Podiatry before an ulcer*	42 % (63)	18% (27)	30% (90)
Total podiatry visits	296	362	658
Therapeutic Shoes and Insoles			
Received shoes or insoles	7.3% (11)	6.6% (10)	7.0% (21)
Received 2nd shoes or insoles	0.6% (1)	2.0% (3)	1.3% (4)
Received 3rd shoes or insoles	0	0	0

* P< 0.001