



Randomized Trial of Telephone Outreach to Improve Medication Adherence and Metabolic Control in Adults With Diabetes

DOI: 10.2337/dc14-0596

Patrick J. O'Connor,¹ Julie A. Schmittiel,²
 Ram D. Pathak,³ Ronald I. Harris,⁴
 Katherine M. Newton,⁵ Kris A. Ohnsorg,¹
 Michele Heisler,⁶ Andrew T. Sterrett,⁷
 Stanley Xu,⁷ Wendy T. Dyer,²
 Marsha A. Raebel,⁷ Abraham Thomas,⁸
 Emily B. Schroeder,⁷ Jay R. Desai,¹ and
 John F. Steiner⁷

OBJECTIVE

Medication nonadherence is a major obstacle to better control of glucose, blood pressure (BP), and LDL cholesterol in adults with diabetes. Inexpensive effective strategies to increase medication adherence are needed.

RESEARCH DESIGN AND METHODS

In a pragmatic randomized trial, we randomly assigned 2,378 adults with diabetes mellitus who had recently been prescribed a new class of medication for treating elevated levels of glycated hemoglobin (A1C) $\geq 8\%$ (64 mmol/mol), BP $\geq 140/90$ mmHg, or LDL cholesterol ≥ 100 mg/dL, to receive 1) one scripted telephone call from a diabetes educator or clinical pharmacist to identify and address nonadherence to the new medication or 2) usual care. Hierarchical linear and logistic regression models were used to assess the impact on 1) the first medication fill within 60 days of the prescription; 2) two or more medication fills within 180 days of the prescription; and 3) clinically significant improvement in levels of A1C, BP, or LDL cholesterol.

RESULTS

Of the 2,378 subjects, 89.3% in the intervention group and 87.4% in the usual-care group had sufficient data to analyze study outcomes. In intent-to-treat analyses, intervention was not associated with significant improvement in primary adherence, medication persistence, or intermediate outcomes of care. Results were similar across subgroups of patients defined by age, sex, race/ethnicity, and study site, and when limiting the analysis to those who completed the intended intervention.

CONCLUSIONS

This low-intensity intervention did not significantly improve medication adherence or control of glucose, BP, or LDL cholesterol. Wide use of this strategy does not appear to be warranted; alternative approaches to identify and improve medication adherence and persistence are needed.

Medication nonadherence remains a principal contributor to poor metabolic control in adults with diabetes. Previous studies (1–5) suggest that ~15–20% of new prescriptions for medications treating elevated levels of glucose, blood pressure (BP), or lipids are never filled and that only ~60% of those who fill a newly prescribed medication in these classes have persisted in taking the medication 12 months later.

¹HealthPartners Institute for Education and Research, Minneapolis, MN

²Kaiser Permanente Division of Research, Northern California, Oakland, CA

³Marshfield Clinic, Marshfield, WI

⁴Geisinger Center for Health Research, Danville, PA

⁵Group Health Research Institute, Seattle, WA

⁶University of Michigan, Center for Clinical Management Research, Ann Arbor, MI

⁷Institute for Health Research, Kaiser Permanente, Denver, CO

⁸Henry Ford Health System, Detroit, MI

Corresponding author: Patrick J. O'Connor, patrick.j.oconnor@healthpartners.com.

Received 7 March 2014 and accepted 15 September 2014.

Clinical trial reg. no. NCT02192255, clinicaltrials.gov.

© 2014 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

However, medication adherence and persistence vary depending on age, sex, number of prescribed medications, and seriousness of the medical condition.

Published reports indicate that medication adherence may be effectively improved through case management by nurses or clinical pharmacists, financial incentives, or other strategies (6,7). However, many of the effective case-management strategies require ongoing contact and are quite expensive (8,9). Despite these encouraging reports of interventions that improve medication adherence and persistence in adults with diabetes, the need for more cost-effective, sustainable intervention strategies is widely recognized. This must be balanced against the need for interventions that are potent enough to achieve their aims. Recent reports (10–12) suggest that low-intensity, well-timed telephone contact may improve primary adherence to statins or other medications.

This multisite pragmatic clinical trial was designed to assess the effectiveness of a single scripted telephone call to diabetes patients who 1) were currently above the recommended clinical goals for glucose, BP, or lipids, and 2) had recently been prescribed a new medication for that specific clinical domain. The goals of the intervention were to improve the primary adherence and persistence to the newly prescribed medication, and to improve control of glucose, BP, and lipids.

RESEARCH DESIGN AND METHODS

Hypothesis, Study Design, and Study Sites

This randomized trial tested the hypothesis that a telephone contact with a patient who had recently been prescribed a new medication for uncontrolled glycated hemoglobin (A1C), BP, or LDL cholesterol would improve 1) primary medication adherence; 2) medication persistence; 3) medication possession ratio (MPR); and 4) A1C, BP, or LDL cholesterol control. This study was part of the larger Agency for Healthcare Research and Quality–funded SUPREME-DM study. The clinical trial reported here was coordinated and led by HealthPartners Institute for Education and Research, while data collection was coordinated through Kaiser Permanente Northwest and analysis was conducted at Kaiser Permanente Colorado. The clinical intervention sites for this study included

Kaiser Permanente Northern California (KPNC), Group Health Cooperative (GHC), Marshfield Clinic, and Geisinger Clinic. These are large multispecialty medical groups that have sophisticated health informatics systems and use electronic medical records (EMRs). Nearly all patients have health insurance with modest copays for visits and prescription medications. Diabetes care is provided by primary care providers, with referrals to diabetes educators and endocrinologists as needed.

Study Subjects

At each clinical intervention site, study subjects were selected if they met the following criteria: 1) they were age 18–75 years of age; 2) they met study criteria for diabetes before study enrollment (13,14); 3) they had received clinical care at a designated clinic or medical center involved in this study for at least 15 months before study enrollment; and 4) were prescribed a new class of medication (not filled in the past 180 days) for A1C, BP, or LDL cholesterol that were uncontrolled at the time of medication prescription (A1C $\geq 8\%$ [64 mmol/mol], systolic BP [SBP] ≥ 140 mmHg, or LDL cholesterol ≥ 100 mg/dL).

Randomization Procedure

We randomized each cluster to receive usual care or intervention. We defined clusters of patients based on their linkage to primary care physicians at KPNC and Marshfield Clinic; linkage to primary care clinics at GHC; or linkage to certified diabetes educators at Geisinger Clinic. Every 2 weeks during the 12-month intervention period, programmers at each study site identified newly eligible study subjects, assigned an index date on which each was identified, and determined whether they were in a usual-care or intervention cluster. The “prescription date” was defined as the date on which a new class of medication was prescribed for uncontrolled A1C, BP, or LDL cholesterol.

Description of Intervention

Those subjects in the intervention arm received a single protocol-structured telephone call from an interventionist who was a nurse health manager (one site); diabetes educator or diabetes educator trainee (one site); or pharmacist (two sites). All interventionists followed the same structured telephone interview protocol to ascertain whether the

subject had started to take the newly prescribed medication; this protocol was modified from one that was effectively used in a prior study (15,16). If the patient was taking the new medication as prescribed, positive reinforcement was provided. If the new medication prescription had not been filled, or was filled but the patient was not taking the medication as directed, the interventionist probed for reasons for nonadherence and worked with the patient to identify and resolve barriers to adherence. The calls were designed to last a median time of <5 min, and up to three call attempts were made to reach each intervention patient. Most intervention telephone calls occurred within 2–6 weeks after the prescription date. Subjects who had filled and those who had not filled their prescription before the index date were identified.

Data Sources

Most study data were obtained directly from extracts of EMR data. These data included demographics, vital signs, laboratory values, encounter dates, and pharmacy prescriptions. Other data, including pharmacy fill data and study enrollment data, were obtained from medical group administrative databases.

Variable Definitions

“Primary medication adherence” was defined as having at least one prescription fill of the index medication within 60 days of the prescription date. “Medication persistence” was defined as having two or more fills of the index medication within 180 days of the prescription date. The MPR denominator was the number of days between the first prescription fill for the new medication and the last day of supplied drug from the last prescription fill for the new medication; the numerator was defined as the number of days of medication supplied in all fills in the denominator time period.

Plan of Analysis

The hypotheses that the intervention would improve 1) primary medication adherence; 2) medication persistence; 3) MPR; and 4) A1C, BP, and LDL cholesterol control were tested using hierarchical logistic regression models and nested linear regression models to assess the proportion of patients who

met medication and care-improvement goals and mean changes in clinical values. If a patient had more than one out-of-control intermediate outcome at the time of the index date, we included all treated intermediate outcomes in the main analyses. If a subject was initially assigned to the intervention or control group and then had a second eligible episode of care during the study period, the patient remained in the initial assignment group. If a subject had insufficient data to assess the change in intermediate outcomes, their adherence and persistence were still assessed if they had sufficient pharmacy coverage.

The principal dependent variable was a composite of improvement from prerandomization to postrandomization values of the clinical domain of interest. To be classified as "improved," each patient had to meet or exceed the following prespecified thresholds for improvement calculated as the difference between the postrandomization value and prerandomization value: A1C greater than or equal to -0.2% (-2 mmol/mol), SBP greater than or equal to -5 mmHg, or LDL cholesterol greater than or equal to -5 mg/dL. These thresholds were assigned a priori to assure that smaller improvements were not classified as clinically significant. These specified changes have a measurable impact on the occurrence of major clinical end points based on prior intervention studies (17–19). We also examined the mean changes in A1C, SBP, and LDL cholesterol levels among the subset of eligible subjects who had been prescribed a new medication for A1C, BP, or LDL cholesterol.

After completion of the primary analysis, a second prespecified analysis was conducted to identify whether the intervention effect varied in subgroups based on age, sex, race, and study site. After completing the intent-to-treat analysis that included all randomized subjects, we conducted a per-protocol analysis in which changes in study outcomes were compared for all usual-care patients versus only intervention-group patients who had been successfully contacted by the interventionist. For this analysis, successful contact was defined as a telephone conversation between the interventionist and the target patient. A last post hoc analysis was conducted to assess the impact of the

intervention in the subgroup of $\sim 20\%$ of subjects in both study arms who had not filled their new prescription between the prescription date and the index date.

Protection of Human Research

Subjects

The institutional review boards at KPNC, GHC, Marshfield Clinic, and Geisinger Clinic reviewed and approved all aspects of the randomized trial reported here, including the intervention, the randomization procedure, data collection, data-transfer protocols, and analysis. This study is registered at clinicaltrials.gov (NCT02192255).

RESULTS

The study randomization resulted in 1,220 eligible patients assigned to the intervention and 1,158 assigned to the control arm, with no significant differences in baseline clinical and demographic characteristics between groups (Table 1). Table 2 shows that, among all study subjects, 1,102 subjects were prescribed a new class of medications to control elevated glucose levels; of these, 969 subjects (88.0%) had both baseline and follow-up A1C values, and 892 (80.9%) had at least 180 days of follow-up and pharmacy coverage. Overall, 76% of subjects in both the intervention and control groups had already filled their new prescription before the index date. There were no significant differences between the intervention and control arms in primary medication adherence within 60 days of the prescription date (intervention arm 85.9%, control arm 87.6%, $P = 0.54$), medication persistence at 180 days (intervention arm 66.9%, control arm 62.2%, $P = 0.14$), MPR (intervention arm 0.80, control arm 0.79, $P = 0.90$), change in A1C level from baseline (intervention arm -1.16% [-12 mmol/mol], control arm -1.33% [-13 mmol/mol], $P = 0.15$), or the proportion of subjects with $\geq 0.2\%$ (2 mmol/mol) improvement in A1C level (intervention arm 73.7%, control arm 75.2%, $P = 0.64$). Among the 195 subjects whose new prescription was not filled before the index date, the change in A1C level from baseline was -1.08% (-11 mmol/mol) in the intervention arm vs. -0.90% (-9 mmol/mol) in the control arm ($P = 0.48$).

Table 2 shows that, among all study subjects, 791 were prescribed a new class of medications to control hypertension; of these, 731 subjects (92.4%) had both baseline and follow-up BP values, and 570 subjects (72.1%) had at least 180 days of follow-up and pharmacy coverage, among whom 76.0% (control arm) and 79.1% (intervention arm) had filled their new prescription before the index date ($P = 0.39$). There were no significant differences between the intervention and control study arms in primary medication adherence within 60 days of the prescription date (intervention arm 85.8%, control arm 83.0%, $P = 0.35$), medication persistence at 180 days (intervention arm 64.0%, control arm 60.3%, $P = 0.30$), MPR (intervention arm 0.90, control arm 0.92, $P = 0.13$), SBP change from baseline (intervention arm -18.1 mmHg, control arm -16.4 mmHg, $P = 0.26$), or the proportion of subjects with a ≥ 4 mmHg drop in SBP (intervention arm 78.5%, control arm 75.8%, $P = 0.38$). Among the 129 subjects whose new prescription was not filled before the index date, the change in SBP from baseline was -19.6 mmHg in the intervention arm vs. -15.6 mmHg in the control arm ($P = 0.25$).

Table 2 also shows that, among all study subjects, 663 subjects were prescribed a new class of medications to control lipid levels, which was almost always a new statin prescription. Of these, 539 subjects (81.3%) had both baseline and follow-up LDL cholesterol values, and 549 subjects (82.8%) had at least 180 days of follow-up and pharmacy coverage, among whom 70.0% (control arm) and 60.9% (intervention arm) had filled their new prescription before the index date ($P = 0.02$)—a difference observed before the intervention. There were no significant differences between the intervention and control arms of the study in primary medication adherence within 60 days of the prescription date (intervention arm 79.6%, control arm 81.9%, $P = 0.47$), medication persistence at 180 days (intervention arm 49.1%, control arm 49.5%, $P = 0.73$), MPR (intervention arm 0.851, control arm 0.846, $P = 0.84$), mean LDL cholesterol change from baseline (intervention arm -30.4 mg/dL, control arm -33.0 mg/dL, $P = 0.44$), or the proportion of subjects with a ≥ 5 mg/dL drop in LDL cholesterol

Table 1.—Baseline patient characteristics shown for 1) all patients and 2) three subsets of patients within both the intervention and the control group who met specific criteria for baseline elevated A1C, LDL cholesterol, or SBP

	Intervention				Control				P value (all patients)
	All patients (n = 1,220)	A1C ≥8% (64 mmol/mol) (n = 569)	LDL cholesterol ≥100 mg/dL (n = 348)	SBP >140/90 mmHg (n = 388)	All patients (n = 1,158)	A1C ≥8% 64 mmol/mol (n = 533)	LDL cholesterol ≥100 mg/dL (n = 315)	SBP ≥140/90 mmHg (n = 403)	
Site, n (%)									
Site A	161 (13.2)	52 (9.1)	31 (8.9)	91 (23.5)	136 (11.7)	46 (8.6)	25 (7.9)	73 (18.1)	0.576
Site B	834 (68.4)	421 (74.0)	269 (77.3)	203 (52.3)	806 (69.6)	400 (75.0)	246 (78.1)	228 (56.6)	
Site C	152 (12.5)	57 (10.0)	38 (10.9)	67 (17.3)	137 (11.8)	49 (9.2)	31 (9.8)	67 (16.6)	
Site D	73 (6.0)	39 (6.9)	10 (2.9)	27 (7.0)	79 (6.8)	38 (7.1)	13 (4.1)	35 (8.7)	
Female sex, n (%)	623 (51.1)	274 (48.2)	189 (54.3)	203 (52.3)	615 (53.1)	263 (49.3)	184 (58.4)	216 (53.6)	0.319
Race, n (%)									
Asian	171 (14.0)	88 (15.5)	62 (17.8)	35 (9.0)	183 (15.8)	89 (16.7)	56 (17.8)	48 (11.9)	0.422
Black	150 (12.3)	64 (11.2)	50 (14.4)	47 (12.1)	140 (12.1)	66 (12.4)	44 (14.0)	51 (12.7)	
Other/unknown	85 (7.0)	46 (8.1)	25 (7.2)	25 (6.4)	66 (5.7)	33 (6.2)	18 (5.7)	21 (5.2)	
White	814 (66.7)	371 (65.2)	211 (60.6)	281 (72.4)	769 (66.4)	345 (64.7)	197 (62.5)	283 (70.2)	
Hispanic									
No	358 (29.3)	134 (23.6)	77 (22.1)	171 (44.1)	329 (28.4)	125 (23.5)	61 (19.4)	165 (40.9)	0.825
Unknown	717 (58.8)	356 (62.6)	231 (66.4)	183 (47.2)	695 (60.0)	333 (62.5)	213 (67.6)	208 (51.6)	
Yes	145 (11.9)	79 (13.9)	40 (11.5)	34 (8.8)	134 (11.6)	75 (14.1)	41 (13.0)	30 (7.4)	
Age at new prescription (years), n (%)									
18–39	59 (4.8)	36 (6.3)	19 (5.5)	6 (1.5)	59 (5.1)	34 (6.4)	18 (5.7)	15 (3.7)	0.553
40–64	653 (53.5)	341 (59.9)	199 (57.2)	171 (44.1)	594 (51.3)	316 (59.3)	179 (56.8)	159 (39.5)	
≥65	508 (41.6)	192 (33.7)	130 (37.4)	211 (54.4)	505 (43.6)	183 (34.3)	118 (37.5)	229 (56.8)	
Age at new prescription (years)*									
Mean ± SD	61.67 ± 13.03	59.10 ± 12.43	59.70 ± 12.63	66.47 ± 12.49	62.04 ± 13.37	59.24 ± 12.79	59.93 ± 13.41	66.23 ± 13.22	0.390
Median (minimum–maximum)	62 (19–94)	60 (19–94)	60.00 (22–91)	66 (33–94)	63 (19–96)	60 (19–96)	60 (24–92)	67 (28–92)	
Drug coverage at new prescription, n (%)*	1,020 (83.6)	485 (85.2)	302 (86.8)	298 (76.8)	975 (84.2)	459 (86.1)	271 (86.0)	322 (79.9)	0.695
Baseline value									
Mean ± SD		9.76 ± 1.66% (83 mmol/mol)	135.1 ± 31.09	155.2 ± 13.51		9.83 ± 1.65% (84 mmol/mol)	138.0 ± 30.56	154.6 ± 13.78	
Median (minimum–maximum)		9.3% (8.0–17.9%) (78 mmol/mol)	126.5 mg/dL (100.0–281.0 mg/dL)	152.0 mmHg (140.0–214.0 mmHg)		9.4% (8.0–15.7%) (79 mmol/mol)	130.0 mg/dL (100.0–274.0 mg/dL)	150.0 mmHg (140.0–220.0 mmHg)	

Note that a given patient could have elevations in more than one clinical domain. *Time of new prescription for first uncontrolled condition when pooled over all conditions.

Table 2—Follow-up time and medication adherence among patients with baseline uncontrolled A1C, BP, or LDL cholesterol, by intervention vs. control condition

	Patients with uncontrolled A1C			Patients with uncontrolled SBP (mm Hg)			Patients with uncontrolled LDL cholesterol (mg/dL)		
	Control	Intervention	P value	Control	Intervention	P value	Control	Intervention	P value
Follow-up ≥ 60 days									
Yes	533 (99.4)	569 (98.9)	0.373	403 (98.0)	388 (98.7)	0.445	315 (98.4)	348 (98.9)	0.583
Follow-up ≥ 180 days									
Yes	533 (94.0)	569 (93.0)	0.593	403 (90.6)	388 (89.4)	0.646	$n = 315$ 294 (93.3)	$n = 348$ 327 (94.0)	0.668
Drug coverage and follow-up ≥ 60 days									
Yes	533 (85.9)	569 (84.5)	0.885	403 (78.7)	388 (76.3)	0.445	315 (85.7)	348 (85.9)	0.572
Drug coverage and follow-up ≥ 180 days									
Yes	533 (81.8)	569 (80.1)	0.792	403 (73.2)	388 (70.9)	0.590	315 (82.5)	348 (83.0)	0.551
Primary adherence by the index date									
Yes	458 (76.0)	481 (75.9)	0.932	317 (76.0)	296 (79.1)	0.389	270 (70.0)	299 (60.9)	0.023
Primary adherence within 60 days of new prescription									
Yes	458 (87.6)	481 (85.9)	0.540	317 (83.0)	296 (85.8)	0.354	270 (81.9)	299 (79.6)	0.474
Nonpersistence (< 2 fills) at 180 days									
Yes	436 (37.8)	456 (33.1)	0.141	295 (39.7)	275 (36.0)	0.301	260 (51.5)	289 (51.9)	0.725
MPR									
Mean \pm SD	0.793 \pm 0.24 0.912	0.802 \pm 0.22 0.881	0.903	0.922 \pm 0.129 1.000	0.900 \pm 0.159 0.995	0.126	0.846 \pm 0.178 0.909	0.851 \pm 0.184 0.930	0.839
Median (minimum–maximum)	(0.151–1.000)	(0.215–1.000)		(0.278–1.000)	(0.188–1.000)		(0.292–1.000)	(0.175–1.000)	
Change from baseline									
Mean \pm SD	–1.33 \pm 1.87% (–13 mmol/mol)	–1.16 \pm 1.80% (–12 mmol/mol)	0.149	–16.4 \pm 18.75 (–103 to 42)	–18.1 \pm 20.56 (–82 to 57)	0.255	–33.0 \pm 38.36 (–162 to 74)	–30.4 \pm 39.02 (–175 to 110)	0.438
Median (minimum to maximum)	(–8.8 to 6.0%)	(–11.0 to 4.7%)		(–16.0 to 42)	(–17.0 to 57)		(–30.0 to 74)	(–28.5 to 110)	
Decrease from baseline*									
Yes	463 (75.2)	506 (73.7)	0.639	368 (75.8)	363 (78.5)	0.380	251 (75.3)	288 (75.7)	0.901
A1C change from baseline among late fillers†									
Mean \pm SD	–0.90 \pm 1.85% (–9 mmol/mol)	–1.08 \pm 1.78% (–11 mmol/mol)	0.476	–15.6 \pm 16.74 (–54 to 30)	–19.6 \pm 21.60 (–68 to 25)	0.248	–22.3 \pm 37.78 (–112 to 74)	–29.2 \pm 35.53 (–108 to 110)	0.227
Median (minimum to maximum)	(–0.95% to 4.0%)	(–0.90% to 3.6%)		(–14.5 to 30)	(–17.5 to 25)		(–17.0 to 74)	(–29.0 to 110)	
	(–9 mmol/mol)	(–9 mmol/mol)							

Data are n (%), unless otherwise specified. *For A1C decrease from baseline of $\geq 0.2\%$. †Late filler is patient with drug coverage who filled a new prescription after the index date.

Table 3—Post hoc power analysis to detect observed differences in clinical parameters as significant within the subset of patients who had not filled their new prescription prior to the index date

Outcome	A1C (%)	SBP (mmHg)	LDL cholesterol (mg/dL)
Difference to detect (observed difference)	0.18 (2 mmol/mol)	4.0	6.9
Pooled SD	1.81	19.2	36.4
Power to detect the observed difference			
Total sample size (INT + CTL)			
100	0.078	0.178	0.155
200	0.108	0.311	0.266
500	0.199	0.642	0.562
1,000	0.349	0.908	0.850

CTL, control; INT, intervention.

(intervention arm 75.7%, control arm 75.3%, $P = 0.90$). Among the 164 subjects whose new prescription was not filled before the index date, the change in LDL cholesterol level from baseline was -29.2 mg/dL in the intervention arm vs. -22.3 mg/dL in the control arm ($P = 0.23$). A post hoc power analysis indicates that substantially larger sample sizes would be required to confirm or refute the observed positive trends in A1C, BP, and LDL cholesterol levels among the subgroup of those subjects who had not filled the new prescription by the index date (Table 3).

In the subgroup of subjects who had not filled their prescription prior to the index date, the differences in LDL cholesterol (6.9 mg/dL) and SBP (4.0 mmHg) are potentially clinically significant, and favored the intervention group. However, the post hoc power analysis shown in Table 3 indicates that substantially larger sample sizes would have been required to adequately assess the statistical significance of these observed differences, and to confirm or refute the observed positive trends in A1C, BP, and LDL cholesterol levels among the subgroup of those subjects who had not filled the new prescription by the index date (Table 3).

Table 4 gives the results of the intent-to-treat and per-protocol analyses of the intervention effect on a composite measure of improvement in A1C, BP, and LDL cholesterol control. Sufficient data to assess this measure were available for 1,090 of 1,220 intervention subjects (89.3%) and for 1,012 of 1,158 control subjects (87.4%). Clinically significant improvement in the clinical domain of interest occurred in 76.3% of intervention group subjects and 75.9% of control group subjects, a statistically and clinically nonsignificant difference.

Additional analyses were performed by the number of clinical domains; among the combined 2,378 study subjects, 2,933 clinical domains were not at goal and had a new medication prescribed. There were sufficient data to adjudicate clinical improvement for 1,157 of 1,501 clinical domains (77.1%) in the intervention group and 1,082 of 1,432 clinical domains (75.6%) in the control group. Clinical improvement occurred in 75.7% of the 1,157 intervention group clinical domains and in 75.4% of the 1,082 control group clinical domains ($P = 0.87$). No significant differences in improvement were noted in subgroups defined by patient sex, age group, race, or ethnicity. There were no significant differences in response to the intervention across the four study sites, or based on whether the case manager was a pharmacist, diabetes educator, or nurse health manager. A secondary analysis that compared differences in improvement between intervention subjects who did and did not have successful contact with the case manager showed no significant main effect ($P = 0.74$) or significant differences in subgroups defined by sex, age group, race, ethnicity, or type of case manager. Results were similar in those with one versus more than one clinical domain targeted by the intervention.

CONCLUSIONS

In this population-based, pragmatic randomized controlled trial, we studied the impact of a very brief structured telephone intervention on clinical and adherence measures among patients with diabetes who had been prescribed a new class of medication for the treatment of uncontrolled A1C, BP, or LDL cholesterol. The telephone intervention was delivered by interventionists who

were pharmacists, diabetes educators, or nurse health managers trained in the use of the study protocol and intervention. The intervention failed to significantly improve primary adherence, medication persistence, MPR, or the main composite measure of improved glucose, BP, or LDL cholesterol control.

There are several possible explanations for the failure of this intervention to have its desired effect. Although the intervention was grounded in a coherent conceptual model and delivered by clinicians carefully trained and monitored for intervention fidelity, it was brief. The intervention included a single telephone contact with a median duration of <5 min delivered a median time of 2–3 weeks after the patient was prescribed a new class of medication for uncontrolled glucose, BP, or lipids. Successful case-manager interventions in prior randomized trials (7,8) of adults with diabetes have typically included much more patient contact, sometimes delivered by a clinic-based nurse working with the patient's primary care provider. However, some prior reports (11,12) indicate that brief telephone contact or other brief interventions may be sufficient to improve primary adherence rates.

It is notable that, in our study, an estimated 76%, 78%, and 66% of those subjects prescribed new medications for glucose, BP, and lipids, respectively, had already filled them before the interventionist called. The high medication fill rate before the intervention call in effect reduced the power for the study to detect intervention effects. We observed differences of 0.18% (2 mmol/mol) in A1C, 4 mmHg in SBP, and 6.9 mg/dL in LDL cholesterol in favor of the intervention group when limiting the analysis to the subset of patients who had not

Table 4—Proportion of patients with improvement in composite measure of improvement in A1C, LDL, or SBP among patients with that clinical condition uncontrolled at baseline

	CTL (n = 1,158 patients; 1,432 clinical domains)	INT (n = 1,220 patients; 1,501 clinical domains)	INT subgroups		P value	
			SC (n = 563 patients; 690 clinical domains)	No SC (n = 657 patients; 811 clinical domains)	INT vs. CTL	SC vs. no SC + CTL
Patient-specific analysis						
Baseline and follow-up measurement	1,158	1,220	563	657		
Yes	1,012 (87.4)	1,090 (89.3)	510 (90.6)	580 (88.3)		
Improvement	1,012	1,090	510	580		
Yes	768 (75.9)	832 (76.3)	391 (76.7)	441 (76.0)	0.813	0.739
Condition-specific analysis						
Baseline and follow-up measurement	1,432	1,501	690	811		
Yes	1,082 (75.6)	1,157 (77.1)	543 (78.7)	614 (75.7)		
Improvement	1,082	1,157	543	614		
Yes	816 (75.4)	876 (75.7)	414 (76.2)	462 (75.2)	0.865	0.676
Improvement by interventionist						
Diabetes educator	102/132 (77.3)	116/148 (78.4)	82/108 (75.9)	34/40 (85.0)	0.536	0.248
Nurse health manager	47/70 (67.1)	49/68 (72.1)	12/18 (66.7)	37/50 (74.0)		
Pharmacist	667/880 (75.8)	711/941 (75.6)	320/417 (76.7)	391/524 (74.6)		
Improvement by sex						
Female	428/564 (75.9)	441/596 (74.0)	220/292 (75.3)	221/304 (72.7)	0.227	0.962
Male	388/518 (74.9)	435/561 (77.5)	194/251 (77.3)	241/310 (77.7)		
Improvement by race						
Asian	126/168 (75.0)	112/161 (69.6)	49/66 (74.2)	63/95 (66.3)	0.597	0.413
Black	98/136 (72.1)	114/151 (75.5)	50/71 (70.4)	64/80 (80.0)		
Other/unknown	54/65 (83.1)	69/79 (87.3)	25/27 (92.6)	44/52 (84.6)		
White	538/713 (75.5)	581/766 (75.8)	290/379 (76.5)	291/387 (75.2)		
Improvement by Hispanic ethnicity						
No	221/307 (72.0)	264/341 (77.4)	119/155 (76.8)	145/186 (78.0)	0.124	0.184
Unknown	500/653 (76.6)	508/677 (75.0)	248/323 (76.8)	260/354 (73.4)		
Yes	95/122 (77.9)	104/139 (74.8)	47/65 (72.3)	57/74 (77.0)		
By age group						
18–39	38/53 (71.7)	34/43 (79.1)	16/20 (80.0)	18/23 (78.3)	0.501	0.202
40–64	420/560 (75.0)	462/621 (74.4)	209/269 (77.7)	253/352 (71.9)		
≥65	358/469 (76.3)	380/493 (77.1)	189/254 (74.4)	191/239 (79.9)		

Data are n (%) or n/T (%), unless otherwise indicated. CNT, control; INT, intervention; SC, successful contact; T, total number of uncontrolled conditions belonging to the indicated subgroup for which baseline and follow-up measurements are available.

already filled their prescription before the intervention call. However, in this patient subgroup, our power to detect these clinically significant improvements in SBP and LDL cholesterol as statistically significant was <20%; the sample size needed for >80% power to detect these differences as significant would have been ~1,000 patients (total for intervention and control) for each of these clinical domains. Future efforts to integrate prescription data with medication fill data would enable more precise identification of early primary nonadherence, and might improve the clinical impact and efficiency of our intervention strategy.

The potential advantages of a more targeted approach are suggested by a recent report that evaluated the impact of an automated telephone call to patients with uncontrolled dyslipidemia who did not fill their new statin prescription within 2 weeks. Compared with a randomized control group, intervention-group patients filled their initial statin prescription more often (42.3% vs. 26.0%, $P < 0.001$) and had significantly better medication persistence at 1 year ($P < 0.001$), although the impact on subsequent LDL cholesterol levels was not reported (10). However, at present, this type of precisely targeted intervention can be delivered in a timely fashion only

in integrated care systems that have close to real-time consolidation of prescription data from EMRs and prescription fill data from pharmacy databases (20).

Notably, only a small fraction of subjects who had levels of A1C $\geq 8\%$ (64 mmol/mol), BP $\geq 140/90$ mmHg, or LDL cholesterol > 100 mg/dL had a new class of medications prescribed at a given clinical encounter. Similar low rates have been previously reported (21–23) in both primary care and subspecialty outpatient practices, and suggest that more attention needs to be directed to the timely initiation and intensification of medications in patients

who are not at key evidence-based clinical goals.

Although primary adherence to newly prescribed medications was in the 75–80% range, the relatively low medication persistence rate we observed at 180 days of follow-up, which was confirmed in data from other studies, underscores the importance of efforts to improve longer-term persistence with key medications (e.g., metformin and statins) that have conferred substantial clinical benefits in multiple randomized trials (19,24–27). It is possible that synchronizing interventions to the date of a second or third medication refill may be a strategy to consider (28,29).

A number of factors limit the interpretation of our data. First, the study was conducted at four medical groups with sophisticated health informatics systems, and the generalization of results to other care settings may be limited. Second, the study was underpowered to detect observed clinically meaningful changes in BP and LDL cholesterol control in the subset of patients who had not filled their new prescription before the intervention call (Table 3). Our inability to rapidly link prescription data with claims data at all sites prevented us from implementing an intervention that targeted only those who had not yet filled their new prescription. Future studies that explore a more targeted approach are needed. Finally, our intervention involved resource-intensive personal calls to patients; more experimentation with less expensive automated communication approaches is needed.

In summary, this low-intensity intervention did not significantly improve medication adherence or intermediate outcomes of diabetes care. Wide use of this strategy may not be warranted, and alternative approaches to identify and improve medication adherence are needed.

Funding. This study was funded by grant HS-019859 from the Agency for Healthcare Research and Quality.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. P.J.O.'C. and K.A.O. wrote, reviewed, and edited the manuscript. J.A.S., R.D.P., R.I.H., K.M.N., and M.H. contributed to the discussion, and revised and edited the manuscript. A.T.S., S.X., and W.T.D. researched the data, and reviewed and edited the manuscript. M.A.R., A.T., E.B.S., J.R.D., and J.F.S.

reviewed and edited the manuscript. P.J.O.C. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- O'Connor PJ. Improving medication adherence: challenges for physicians, payers, and policy makers. *Arch Intern Med* 2006;166:1802–1804
- Karter AJ, Parker MM, Moffet HH, Ahmed AT, Schmittiel JA, Selby JV. New prescription medication gaps: a comprehensive measure of adherence to new prescriptions. *Health Serv Res* 2009;44:1640–1661
- Raebel MA, Ellis JL, Carroll NM, et al. Characteristics of patients with primary non-adherence to medications for hypertension, diabetes, and lipid disorders. *J Gen Intern Med* 2012;27:57–64
- Steiner JF. Rethinking adherence. *Ann Intern Med* 2012;157:580–585
- Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med* 2005;353:487–497
- Kripalani S, Yao X, Haynes RB. Interventions to enhance medication adherence in chronic medical conditions: a systematic review. *Arch Intern Med* 2007;167:540–550
- McDonald KM, Chang C, Schultz E. *Through the Quality Kaleidoscope: Reflections on the Science and Practice of Improving Health Care Quality. Closing the Quality Gap: Revisiting the State of the Science*. Rockville, MD, Agency for Healthcare Research and Quality, 2013
- Aubert RE, Herman WH, Waters J, et al. Nurse case management to improve glycemic control in diabetic patients in a health maintenance organization. A randomized, controlled trial. *Ann Intern Med* 1998;129:605–612
- Tricco AC, Ivers NM, Grimshaw JM, et al. Effectiveness of quality improvement strategies on the management of diabetes: a systematic review and meta-analysis. *Lancet* 2012;379:2252–2261
- Deroose SF, Green K, Marrett E, et al. Automated outreach to increase primary adherence to cholesterol-lowering medications. *JAMA Intern Med* 2013;173:38–43
- Vollmer WM, Feldstein A, Smith DH, et al. Use of health information technology to improve medication adherence. *Am J Manag Care* 2011;17:SP79–SP87
- Viswanathan M, Golin CE, Jones CD, et al. Interventions to improve adherence to self-administered medications for chronic diseases in the United States: a systematic review. *Ann Intern Med* 2012;157:785–795
- Desai JR, Wu P, Nichols GA, Lieu TA, O'Connor PJ. Diabetes and asthma case identification, validation, and representativeness when using electronic health data to construct registries for comparative effectiveness and epidemiologic research. *Med Care* 2012;50(Suppl.):S30–S35
- Nichols GA, Desai J, Elston Lafata J, et al.; SUPREME-DM Study Group. Construction of a multisite DataLink using electronic health records for the identification, surveillance, prevention, and management of diabetes mellitus: the SUPREME-DM project. *Prev Chronic Dis* 2012;9:E110
- Heisler M, Hofer TP, Schmittiel JA, et al. Improving blood pressure control through a clinical pharmacist outreach program in patients with diabetes mellitus in 2 high-performing health systems: the adherence and intensification of medications cluster randomized, controlled pragmatic trial. *Circulation* 2012;125:2863–2872
- Heisler M, Hofer TP, Klamers ML, et al. Study protocol: the Adherence and Intensification of Medications (AIM) study—a cluster randomized controlled effectiveness study. *Trials* 2010;11:95
- Law MR, Morris JK, Wald NJ. Use of blood pressure lowering drugs in the prevention of cardiovascular disease: meta-analysis of 147 randomised trials in the context of expectations from prospective epidemiological studies. *BMJ* 2009;338:b1665
- Baigent C, Blackwell L, Emberson J, et al.; Cholesterol Treatment Trialists' (CTT) Collaboration. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* 2010;376:1670–1681
- Holman RR, Paul SK, Bethel MA, Matthews DR, Neil HA. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008;359:1577–1589
- Fischer MA. Chipping away: improving primary medication adherence. *JAMA Intern Med* 2013;173:44–45
- Schmittiel JA, Uratsu CS, Karter AJ, et al. Why don't diabetes patients achieve recommended risk factor targets? Poor adherence versus lack of treatment intensification. *J Gen Intern Med* 2008;23:588–594
- Bramley TJ, Gerbino PP, Nightengale BS, Frech-Tamas F. Relationship of blood pressure control to adherence with antihypertensive monotherapy in 13 managed care organizations. *J Manag Care Pharm* 2006;12:239–245
- O'Connor PJ, Sperl-Hillen JM, Rush WA, et al. Impact of electronic health record clinical decision support on diabetes care: a randomized trial. *Ann Fam Med* 2011;9:12–21
- Colhoun HM, Betteridge DJ, Durrington PN, et al.; CARDS investigators. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet* 2004;364:685–696
- Stuart BC, Simoni-Wastila L, Zhao L, Lloyd JT, Doshi JA. Increased persistency in medication use by U.S. Medicare beneficiaries with diabetes is associated with lower hospitalization rates and cost savings. *Diabetes Care* 2009;32:647–649
- Rasmussen JN, Chong A, Alter DA. Relationship between adherence to evidence-based pharmacotherapy and long-term mortality after acute myocardial infarction. *JAMA* 2007;297:177–186
- Aubert RE, Yao J, Xia F, Garavaglia SB. Is there a relationship between early statin compliance and a reduction in healthcare utilization? *Am J Manag Care* 2010;16:459–466
- Franklin JM, Shrank WH, Pakes J, et al. Group-based trajectory models: a new approach to classifying and predicting long-term medication adherence. *Med Care* 2013;51:789–796
- Bryson CL, Au DH, Young B, McDonnell MB, Fihn SD. A refill adherence algorithm for multiple short intervals to estimate refill compliance (ReComp). *Med Care* 2007;45:497–504