



Diabetic Ketoacidosis and Related Events in the Canagliflozin Type 2 Diabetes Clinical Program

DOI: 10.2337/dc15-1251

Ngozi Erondu, Mehul Desai, Kirk Ways, and Gary Meininger

OBJECTIVE

This study assessed the incidence of serious adverse events of diabetic ketoacidosis (DKA) among patients with type 2 diabetes treated with canagliflozin.

RESEARCH DESIGN AND METHODS

All serious adverse events of DKA and related events (ketoacidosis, metabolic acidosis, and acidosis) from 17,596 patients from randomized studies of canagliflozin through 11 May 2015 were analyzed.

RESULTS

Serious adverse events of DKA and related events were reported in 12 patients (0.07%), including 4 (0.07%), 6 (0.11%), and 2 (0.03%) treated with canagliflozin 100 and 300 mg and comparator, respectively; corresponding incidence rates were 0.522, 0.763, and 0.238 per 1,000 patient-years, respectively. Most patients with DKA and related events had a blood glucose >300 mg/dL (16.7 mmol/L) at presentation of DKA, were on insulin, and had DKA-precipitating factors, including some with type 1 diabetes/latent autoimmune diabetes of adulthood.

CONCLUSIONS

DKA and related events occurred at a low frequency in the canagliflozin type 2 diabetes program, with an incidence consistent with limited existing observational data in the general population with type 2 diabetes.

On 15 May 2015, the U.S. Food and Drug Administration (FDA) issued a Drug Safety Communication based upon a search of the FDA Adverse Event Reporting System database that indicated that medicines for type 2 diabetes in the sodium glucose cotransporter 2 (SGLT2) inhibitor class (which includes canagliflozin, empagliflozin, and dapagliflozin) may lead to ketoacidosis. The FDA also noted that patients may present atypically, with only slightly increased levels of blood glucose (1). In addition, several case reports and series have described diabetic ketoacidosis (DKA) in patients with type 1 diabetes or type 2 diabetes treated with SGLT2 inhibitors (2–4).

RESEARCH DESIGN AND METHODS

An analysis of all serious adverse events of DKA and related terms of ketoacidosis, metabolic acidosis, and acidosis was performed using a database that contained data from 17,596 patients, with nearly 24,000 patient-years of exposure, compiled from completed and ongoing randomized, controlled clinical studies of canagliflozin. The overall mean exposure in this analysis was 1.4 years. Table 1 includes details regarding the studies included in this analysis, which was conducted by Janssen Research & Development, LLC (the sponsor of canagliflozin). A history of type 1

Janssen Research & Development, LLC, Raritan, NJ

Corresponding author: Gary Meininger, gmeininger@its.jnj.com.

Received 11 June 2015 and accepted 26 June 2015.

© 2015 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

See accompanying article, p. XXX.

Table 1—Randomized, controlled studies of canagliflozin included in the analysis of DKA and related events

Study/status	Study design/population	Key inclusion criteria			Treatment groups	Reference
		Age, years	HbA _{1c} , % (mmol/mol)	eGFR, mL/min/1.73 m ²		
DIA3002 completed	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 3-arm, parallel-group study (with a 26-week, core double-blind period plus a 26-week, extension double-blind period) Men and women with type 2 diabetes on metformin and sulfonyleurea therapy 	18 to 80	7.0 to 10.5 (53 to 91)	≥55	Canagliflozin 100 mg Canagliflozin 300 mg Placebo (1:1:1)	(9)
DIA3004 completed	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 3-arm, parallel-group study (with a 26-week, core double-blind period plus a 26-week, extension double-blind period) Men and women with type 2 diabetes who have moderate renal impairment on currently available standard of care AHA therapies 	≥25	7.0 to 10.5 (53 to 91)	≥30 to <50	Canagliflozin 100 mg Canagliflozin 300 mg Placebo (1:1:1)	(10,11)
DIA3005 completed	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 3-arm, parallel-group study (with a 26-week, core double-blind period plus a 26-week, active-controlled, extension double-blind period)* Men and women with type 2 diabetes (monotherapy) 	18 to 80	7.0 to 10.0 (53 to 86)	≥50	Canagliflozin 100 mg Canagliflozin 300 mg Placebo (1:1:1)	(12,13)
DIA3006 completed	<ul style="list-style-type: none"> Randomized, double-blind, parallel-group study (with a 26-week placebo- and active-controlled, core double-blind period and a 26-week active-controlled, extension double-blind period) Men and women with type 2 diabetes on metformin therapy 	18 to 80	7.0 to 10.5 (53 to 91)	≥55	Canagliflozin 100 mg Canagliflozin 300 mg Sitagliptin Placebo (2:2:2:1)	(14)
DIA3008 (CANVAS) ongoing	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, parallel-group cardiovascular assessment study Men and women with type 2 diabetes on currently available standard of care AHA therapies 	≥30	7.0 to 10.5 (53 to 91)	≥30	Canagliflozin 100 mg Canagliflozin 300 mg Placebo (1:1:1)	(15,16)
DIA3009 completed	<ul style="list-style-type: none"> Randomized, double-blind, active-controlled, parallel-group study (with a 52-week, core double-blind period plus a 52-week, extension double-blind period) Men and women with type 2 diabetes on metformin therapy 	18 to 80	7.0 to 10.5 (53 to 91)	≥55	Canagliflozin 100 mg Canagliflozin 300 mg Glimepiride (1:1:1)	(17,18)
DIA3010 completed	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, parallel-group study (with a 26-week, core double-blind period plus a 78-week, extension double-blind period) Men and women with type 2 diabetes on currently available standard of care AHA therapies 	55 to 80	7.0 to 10.0 (53 to 86)	≥50	Canagliflozin 100 mg Canagliflozin 300 mg Placebo (1:1:1)	(19,20)
DIA3012 completed	<ul style="list-style-type: none"> Randomized, double-blind, parallel-group, 3-arm study (with a 26-week, placebo-controlled, core double-blind period plus a 26-week, active-controlled, extension double-blind period) Men and women with type 2 diabetes on metformin and pioglitazone therapy 	18 to 80	7.0 to 10.5 (53 to 91)	≥55	Canagliflozin 100 mg Canagliflozin 300 mg Placebo (1:1:1)	(21)

Continued on p. 3

Table 1—Continued

Study/status	Study design/population	Key inclusion criteria			Treatment groups	Reference
		Age, years	HbA _{1c} , % (mmol/mol)	eGFR, mL/min/1.73 m ²		
DIA3015 completed	<ul style="list-style-type: none"> Randomized, double-blind, 52-week, active-controlled study Men and women with type 2 diabetes on metformin and sulfonylurea therapy 	≥18	7.0 to 10.5 (53 to 91)	≥55	Canagliflozin 300 mg Sitagliptin (1:1)	(22)
DNE3001 (CREDESCENCE) ongoing	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 2-arm, parallel-group, event-driven, multicenter study Men and women with type 2 diabetes and diabetic nephropathy 	≥30	6.5 to 12.0 (48 to 108)	30 to 90	Canagliflozin 100 mg Placebo (1:1)	—
DIA4002 ongoing†	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 3-arm, parallel-group, multicenter study Men and women with type 2 diabetes and hypertension 	18 to <75	7.0 to <10.0 (53 to <86)	≥60	Canagliflozin 100 mg Canagliflozin 300 mg Placebo (1:1:1)	—
DIA4003 (CANVAS-R) ongoing	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 2-arm, parallel-group, multicenter study Men and women with type 2 diabetes receiving standard of care, but with inadequate glycemic control and at elevated risk of cardiovascular events 	≥30	7.0 to 10.5 (53 to 91)	≥30	Canagliflozin 100 mg (with titration to canagliflozin 300 mg) Placebo (1:1)	—
DIA4004 ongoing	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 2-arm, parallel-group, multicenter study Men and women with type 2 diabetes on metformin and sitagliptin therapy 	18 to 75	7.5 to 10.5 (58 to 91)	≥60	Canagliflozin 100 mg (with titration to canagliflozin 300 mg) Placebo (1:1)	—
DIA2003 completed	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled, 3-arm, parallel-group, 18-week, multicenter study Men and women with type 2 diabetes with inadequate glycemic control on metformin therapy 	18 to 80	7.5 to 10.5 (58 to 91)	≥55	Canagliflozin 50 mg BID Canagliflozin 150 mg BID Placebo (1:1:1)	(23)
DIA3011 completed	<ul style="list-style-type: none"> Randomized, double-blind, active-controlled, parallel-group, 26-week multicenter study of initial combination therapy with canagliflozin and metformin Men and women with drug-naïve type 2 diabetes 	18 to <75	7.5 to 12.0 (58 to 108)	≥60	Metformin XR Canagliflozin 100 mg Canagliflozin 300 mg Canagliflozin 100 mg/ metformin XR Canagliflozin 300 mg/ metformin XR (1:1:1:1:1)	(24)

AHA, antihyperglycemic agent; BID, twice daily; CREDESCENCE, Canagliflozin and Renal Events in Diabetes with Established Nephropathy Clinical Evaluation; eGFR, estimated glomerular filtration rate; XR, extended release. *DIA3005 also had a 26-week high glycemic substudy that was not included in the current analysis. No adverse events of DKA, ketoacidosis, metabolic acidosis, or acidosis were reported in this substudy. †Clinical conduct is completed; final clinical study report is in progress.

diabetes or DKA was an exclusion criterion in all studies. Ascertainment of potential events for inclusion in this analysis was done using investigator-reported adverse events. Four adverse event terms (i.e., diabetic ketoacidosis, ketoacidosis, metabolic acidosis, and acidosis) from the *Medical Dictionary for Regulatory Activities* (MedDRA) were searched. Cases meeting standard criteria for a regulatory definition of a serious adverse event (e.g., resulting in hospitalization or a medically important

event) were included in this analysis. All unblinded cases in this analysis came from completed studies or unblinded data sets previously used to support canagliflozin global marketing dossiers or required for responses to health authorities. Through 11 May 2015, there were 12 patients with 13 unblinded serious adverse events of DKA, ketoacidosis, metabolic acidosis, and acidosis, and 3 additional serious adverse events that remain blinded and were not included in the current analysis. These 3 additional events

come from the ongoing CANagliflozin cardiovascular Assessment Study (CANVAS), which is blinded and is being monitored by an independent data monitoring committee. Data from the 12 unblinded patients with serious adverse events are discussed below.

RESULTS

The incidence of serious adverse events of DKA and related events in the canagliflozin randomized clinical trial database was 0.07% (12 of 17,596). The incidence

of serious adverse events of DKA and related events by treatment group was 0.07% (4 of 5,337), 0.11% (6 of 5,350), and 0.03% (2 of 6,909) with canagliflozin 100 and 300 mg and comparator, respectively; corresponding incidence rates were 0.522, 0.763, and 0.238 per 1,000 patient-years, respectively. After being diagnosed with a DKA-related event, 6 patients on canagliflozin (3 on canagliflozin 100 mg, 3 on canagliflozin 300 mg, and none on comparator) were reported to have autoimmune diabetes (latent autoimmune diabetes of adulthood [LADA] or type 1 diabetes) or tested positive for GAD65 antibodies. Excluding these 6 patients, the incidences of serious adverse events of DKA and related events by treatment group in patients with type 2 diabetes were 0.02% (1 of 5,334), 0.06% (3 of 5,347), and 0.03% (2 of 6,909) with canagliflozin 100 and 300 mg and comparator, respectively, with corresponding incidence rates of 0.130, 0.381, and 0.238 per 1,000 patient-years, respectively. The race and ethnicity of these patients were as follows: 1 Hispanic or Latino, American Indian, or Alaska native patient; 1 Hispanic or Latino white patient; and 10 non-Hispanic or Latino white patients. Compared with other patients in the canagliflozin program, these 12

patients were predominantly male, white, older, and had a longer duration of diabetes, lower BMI, higher HbA_{1c}, and lower estimated glomerular filtration rate at baseline (Table 2). Specific details of the 12 patients with serious adverse events of DKA and related events are reported in Table 3. Eight of the 12 patients in this analysis were enrolled in the CANVAS trial, which included patients with significant comorbid conditions; of these 8 patients, all 7 in the canagliflozin treatment groups were on insulin. The 10 patients with blood glucose values reported at presentation had levels that were >300 mg/dL (16.7 mmol/L) and ranged from 347 to 571 mg/dL (19.3 to 31.7 mmol/L). One other patient on canagliflozin 300 mg had several blood glucose levels ranging from 148 to 320 mg/dL (8.2 to 17.8 mmol/L), but dates and times of these measurements were not provided. Of the 10 patients on canagliflozin with a DKA-related event, 8 were receiving insulin therapy (note that ~31% of patients [n = 5,407] in the canagliflozin type 2 diabetes program were on background insulin therapy), with 4 having questionable compliance with insulin therapy at the time of the event. One of these 4 patients also had a second event postoperatively after a cholecystectomy. The other 4 patients on insulin

therapy with a DKA-related event had concomitant diagnoses of pancreatic cancer, myocardial infarction, gastroenteritis, and viral infection. Among the two canagliflozin patients not on insulin therapy with an event, one had type 1 diabetes and one had a subcutaneous abscess and chronic pancreatitis.

CONCLUSIONS

In summary, DKA and related events occurred at a low frequency in patients participating in the randomized, controlled, canagliflozin type 2 diabetes clinical trial program. Although there are limited epidemiological data on the incidence of DKA in patients with type 2 diabetes, the overall incidence rates of these events in the current analysis are consistent with the broad range reported in existing observational data. Specifically, a study in Northern Sweden reported an estimated DKA incidence rate of 0.5 per 1,000 patient-years (5), and an analysis of four large U.S. commercial claims databases (i.e., the Truven MarketScan Commercial Claims and Encounters, MarketScan Medicare Supplemental Beneficiaries, the MarketScan Multistate Medicaid Database, and the Optum ClinFormatics database) found a DKA incidence rate in the range of 0.32 to 2.0 per 1,000 patient-years (data on file). However, given the potential for incomplete reporting or underreporting of DKA, the incidence of DKA in patients with type 2 diabetes, including patients treated with canagliflozin and other SGLT2 inhibitors, may be underestimated.

Although there were some differences in baseline characteristics between all patients and the subset of patients who developed DKA and related events, there was no clear baseline clinical phenotype that allowed the identification of specific individual patients at risk for developing DKA. Nevertheless, most patients had a known precipitating factor for DKA at the time of these events. Some reports note that patients who presented with DKA had atypically low blood glucose values; however, of the 10 patients treated with canagliflozin who presented with DKA and related events and had available blood glucose values at presentation, 9 patients had blood glucose values >250 mg/dL (13.9 mmol/L). We postulate that patients diagnosed as having type 2 diabetes or misdiagnosed as

Table 2—Background demographic and disease characteristics of patients with and without serious adverse events of DKA and related events

	Patients with DKA (n = 12)	Patients without DKA (n = 17,584)
Sex, n (%)		
Male	9 (75.0)	7,182 (40.8)
Female	3 (25.0)	10,401 (59.2)
Age, years	69.5 (47, 78)	61.0 (20, 96)
Race, n (%)		
White	11 (91.7)	13,480 (76.7)
Black/African American	0	703 (4.0)
Asian	0	2,148 (12.2)
Other†	1 (8.3)	1,253 (7.1)
Ethnicity, n (%)		
Hispanic or Latino	2 (16.7)	3,118 (17.7)
Not Hispanic or Latino	10 (83.3)	14,385 (81.8)
Other‡	0	81 (0.5)
HbA _{1c} , %	8.9 (7, 11)	8.0 (5, 14)
HbA _{1c} , mmol/mol	74 (53, 97)	66 (31, 130)
BMI, kg/m ²	27.1 (23, 34)	31.3 (15, 73)
eGFR, mL/min/1.73 m ²	69.0 (33, 127)	79.0 (10, 227)
Duration of diabetes, years	13.5 (1, 29)	9.0 (0, 55)

Data are median (range) unless otherwise indicated. eGFR, estimated glomerular filtration rate. †Includes American Indian or Alaskan Native, Native Hawaiian or other Pacific Islander, multiple, other, unknown, and not reported. ‡Includes unknown and not reported.

Table 3—Summary of patients with treatment-emergent serious adverse events of DKA and related events in the canagliflozin development program for type 2 diabetes

Patient	1	2	3	4	5	6	7	8	9	10	11	12
Age, years	73	66	73	76	50	74	73	78	47	66	57	62
BMI, kg/m ²	25.7	27.1	28.8	22.7	22.7	25.4	34.2	27.0	29.6	30.5	24.9	29
Sex	M	M	M	M	M	M	F	F	F	M	M	M
Evidence of autoimmune diabetes (Type 1 diabetes, LADA, GAD65 antibody positive)	Yes	No	No	Yes	No	Yes	Yes	Yes	Yes	No	No	No
Diabetes duration (at randomization), years	21	22	20	14	10	30	11	20	1	12	13	1
Baseline HbA _{1c} , % (mmo/mol)	9.1 (76)	7.8 (62)	8.7 (72)	8.4 (68)	8.0 (64)	7.9 (63)	10.5 (91)	9.6 (81)	9.3 (78)	7.2 (65)	9.9 (85)	10.5 (91)
Baseline C-peptide value, nmo/L (ng/mL)	0.17 (0.51) Low*	1.19 (3.57)	N/A	<0.02 (<0.07) Low*	<0.02 (<0.07) Low*	0.03 (0.10) Low*	N/A	N/A	0.14 (0.43) Low*	N/A	0.34 (1.02)	N/A
Treatment group	CANA 300 mg	Placebo	CANA 100 mg	CANA 100 mg	CANA 300 mg	CANA 300 mg	CANA 300 mg	CANA 100 mg	CANA 100 mg	CANA 300 mg	SITA 100 mg	CANA 300 mg
Adverse event	Acidosis DKA (non-TEAE)	Metabolic acidosis	DKA	DKA	Metabolic acidosis	DKA	Ketoacidosis	DKA	DKA	DKA	DKA	Ketoacidosis
Onset day relative to first dose	Acidosis: 618 DKA: 1,226 (stopped treatment day 1,194)	Admitted day 731 (stopped treatment day 693)	454	21	54	288	744	536	212	720 (stopped treatment day 719)	256	18
Background AHA(s)	INS	MET, GLIP	INS	INS	INS	INS	INS, MET	INS, MET	MET, GLIM	INS (started 2 days prior to DKA onset), EXEN, GLIC, MET	INS	None
Blood glucose, mg/dL (mmo/L)†	Acidosis: 369 (20.5) DKA: 533 (29.6)	N/A	400 (22.2)	347 (19.3)	>500 (>27.8)	>500 (>27.8)	148–320 (8.2–17.8)†	481 (26.7)	400 (22.2)	470 (26.1)	481 (26.7)§	571 (31.7)
pH	Acidosis: 7.24 DKA: N/A	N/A	7.14	N/A	6.82	N/A	N/A	7.23	7.022	N/A	7.22§	N/A
Bicarbonate, mEq/L	Acidosis: 15 DKA: 15	N/A	15	N/A	3.4	N/A	13.6§	11.7	1.8	N/A	11.4§	N/A
Anion gap, mmo/L	Acidosis: 6 DKA: 17	N/A	25	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Ketones (blood or urine)	Acidosis: +blood DKA: +blood, +urine	N/A	+blood	N/A	+blood	N/A	N/A	+blood	N/A	N/A	N/A	+Urine

Continued on p. 6

Table 3—Continued

Patient	1	2	3	4	5	6	7	8	9	10	11	12
Confounder factors	<ul style="list-style-type: none"> Acidosis: History of LADA Acute cholecystitis requiring laparoscopic cholecystectomy Acidosis developed postoperatively 	<ul style="list-style-type: none"> History of alcohol abuse Admitted with left lower lobe infiltrate, sepsis, respiratory failure, metastatic colorectal cancer 	<ul style="list-style-type: none"> Hyperglycemia the day before hospitalization for DKA thought to be due to “bad insulin” Changed reservoir, tubing, and site of the insulin pump Self-administered 116–117 units of insulin because blood glucose levels remained elevated Blood glucose still remained elevated and he went to the ER, where he presented with dehydration, hypotension, tachycardia, and elevated CK-MB Elevated troponin levels were noted the next day, and the event was adjudicated as an MI 	<ul style="list-style-type: none"> Nausea, vomiting, and diarrhea the day before hospitalization for DKA Patient did not take usual insulin dose on day of hospitalization Nonfatal STEMI occurred day after DKA Subsequently tested positive for GAD65 and insulin antibodies[§] 	<ul style="list-style-type: none"> Vomiting 2 days before hospitalization, which apparently led to the interruption of insulin 	<ul style="list-style-type: none"> On insulin since diagnosis (~30 years) Reduced usual insulin dose due to reduced blood glucose after study start Unintentional weight loss of ~13.6 kg over ~6 months Medication and dietary noncompliance Infectious gastroenteritis with continuous vomiting 3 days prior to DKA Elevated transaminases noted during hospitalization (nonviral hepatitis) Subsequently tested positive for GAD65 and insulin antibodies[§] 	<ul style="list-style-type: none"> Did not take insulin injections for 4 days prior to hospitalization due to technical problems with insulin pen Subsequently diagnosed with LADA 	<ul style="list-style-type: none"> Associated with RSV infection and faulty insulin injection technique (assessed during the hospitalization) Subsequently tested positive for GAD65 and insulin antibodies[§] 	<ul style="list-style-type: none"> 45.4 kg weight loss within <2 years Subsequent diagnosis with type 1 diabetes (positive for GAD65) 	<ul style="list-style-type: none"> UTI from days 656 to 678, 692 to 718, and day 736 Pancreatic cancer with liver metastasis diagnosed on day 786 	<ul style="list-style-type: none"> Acute gastroenteritis started on day 255 Developed septic shock and acute renal failure in addition to DKA Patient died of acute MI (cause of death from autopsy report) on day 258 	<ul style="list-style-type: none"> Heart failure class II and on indapamide Absessed boil of the anterior abdominal wall which required dissection and antibiotics Abdominal ultrasound showed chronic pancreatitis

Abbreviations: AHA, antihyperglycemic agent; CANA, canagliflozin; CK-MB, creatinine kinase-myoglobin; ER, emergency room; EXEN, exenatide; F, female; GLUC, glimepiride; GULM, glimepiride; GIP, glipeptide; GIIP, glipeptide; M, male; MET, metformin; MI, myocardial infarction; N/A, not available; RSV, respiratory syncytial virus; SFA, sitagliptin; STEMI, ST-segment elevation myocardial infarction; TEAE, treatment-emergent adverse event (defined as an adverse event that occurred during the treatment period or within 30 days since the last dose of the study medication); UTI, urinary tract infection. *Per normal laboratory range of 0.27 to 1.28 mmol/L (0.81 to 3.85 ng/mL). †Blood glucose value at presentation of the adverse event. ‡Range of all values reported; specific days and times not reported. §Specific date not reported. ||GAD65 antibody titers ≥17× the upper limit of normal (20 DK units/mL); IA-2 antibody titers were negative.

type 2 diabetes (e.g., LADA, type 1 diabetes), and who have a low β-cell reserve coupled with a potential SGLT2 inhibitor-associated increase in glucagon (6–8), are unable to produce sufficient insulin to suppress hepatic ketogenesis and peripheral lipolysis, which in the setting of an acute illness (and associated increase in insulin resistance), can develop DKA. Further prospective research is needed to better understand the incidence and underlying mechanism(s) of DKA associated with SGLT2 inhibitors.

Funding. This article was supported by Janssen Research & Development, LLC. Technical editorial assistance was provided by Alaina DeToma, PhD, of MedErgy HealthGroup, and was funded by Janssen Global Services, LLC. Canagliflozin has been developed by Janssen Research & Development, LLC, in collaboration with Mitsubishi Tanabe Pharma Corporation.

Duality of Interest. N.E., M.D., K.W., and G.M. are full-time employees of Janssen Research & Development, LLC. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. N.E., M.D., K.W. and G.M. contributed to the design, acquisition, analysis and interpretation of the data, and development of the manuscript. All authors approved the final version of the manuscript submitted. G.M. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. U.S. Food and Drug Administration. FDA drug safety communication: FDA warns that SGLT2 inhibitors for diabetes may result in a serious condition of too much acid in the blood. [article online], 2015. Available from <http://www.fda.gov/Drugs/DrugSafety/ucm446845.htm>. Accessed 19 May 2015
2. Hine J, Paterson H, Abrol E, Russell-Jones D, Herring R. SGLT inhibition and euglycaemic diabetic ketoacidosis. *Lancet Diabetes Endocrinol* 2015;3:503–504
3. Hayami T, Kato Y, Kamiya H, et al. Case of ketoacidosis by a sodium-glucose cotransporter 2 inhibitor in a diabetic patient with a low-carbohydrate diet. *J Diabetes Investig* 20 February 2015 [E-pub ahead of print]. DOI: 10.1111/jdi.12330
4. Burr K, Nguyen A-T, Rasouli N. A case report of ketoacidosis associated with canagliflozin (Invokana). *In Diabetes Case Reports II*. *Endocr Rev* 2015;36. Abstract SAT-595
5. Wang ZH, Kihl-Selstam E, Eriksson JW. Ketoacidosis occurs in both type 1 and type 2 diabetes—a population-based study from Northern Sweden. *Diabet Med* 2008;25:867–870
6. Ferrannini E, Muscelli E, Frascerra S, et al. Metabolic response to sodium-glucose cotransporter 2 inhibition in type 2 diabetic patients. *J Clin Invest* 2014;124:499–508

7. Merovci A, Solis-Herrera C, Daniele G, et al. Dapagliflozin improves muscle insulin sensitivity but enhances endogenous glucose production. *J Clin Invest* 2014;124:509–514
8. Bonner C, Kerr-Conte J, Gmyr V, et al. Inhibition of the glucose transporter SGLT2 with dapagliflozin in pancreatic alpha cells triggers glucagon secretion. *Nat Med* 2015;21:512–517
9. Wilding JP, Charpentier G, Hollander P, et al. Efficacy and safety of canagliflozin in patients with type 2 diabetes mellitus inadequately controlled with metformin and sulphonylurea: a randomised trial. *Int J Clin Pract* 2013;67:1267–1282
10. Yale JF, Bakris G, Cariou B, et al. Efficacy and safety of canagliflozin in subjects with type 2 diabetes and chronic kidney disease. *Diabetes Obes Metab* 2013;15:463–473
11. Yale JF, Bakris G, Cariou B, et al.; DIA3004 Study Group. Efficacy and safety of canagliflozin over 52 weeks in patients with type 2 diabetes mellitus and chronic kidney disease. *Diabetes Obes Metab* 2014;16:1016–1027
12. Stenlöf K, Cefalu WT, Kim KA, et al. Efficacy and safety of canagliflozin monotherapy in subjects with type 2 diabetes mellitus inadequately controlled with diet and exercise. *Diabetes Obes Metab* 2013;15:372–382
13. Stenlöf K, Cefalu WT, Kim KA, et al. Long-term efficacy and safety of canagliflozin monotherapy in patients with type 2 diabetes inadequately controlled with diet and exercise: findings from the 52-week CANTATA-M study. *Curr Med Res Opin* 2014;30:163–175
14. Lavallo-González FJ, Januszewicz A, Davidson J, et al. Efficacy and safety of canagliflozin compared with placebo and sitagliptin in patients with type 2 diabetes on background metformin monotherapy: a randomised trial. *Diabetologia* 2013;56:2582–2592
15. Neal B, Perkovic V, de Zeeuw D, et al. Rationale, design, and baseline characteristics of the CANagliflozin cardioVascular Assessment Study (CANVAS)—a randomized placebo-controlled trial. *Am Heart J* 2013;166:217–223.e11
16. Neal B, Perkovic V, de Zeeuw D, et al.; CANVAS Trial Collaborative Group. Efficacy and safety of canagliflozin, an inhibitor of sodium-glucose cotransporter 2, when used in conjunction with insulin therapy in patients with type 2 diabetes. *Diabetes Care* 2015;38:403–411
17. Cefalu WT, Leiter LA, Yoon KH, et al. Efficacy and safety of canagliflozin versus glimepiride in patients with type 2 diabetes inadequately controlled with metformin (CANTATA-SU): 52 week results from a randomised, double-blind, phase 3 non-inferiority trial. *Lancet* 2013;382:941–950
18. Leiter LA, Yoon KH, Arias P, et al. Canagliflozin provides durable glycemic improvements and body weight reduction over 104 weeks versus glimepiride in patients with type 2 diabetes on metformin: a randomized, double-blind, phase 3 study. *Diabetes Care* 2015;38:355–364
19. Bode B, Stenlöf K, Sullivan D, Fung A, Usiskin K. Efficacy and safety of canagliflozin treatment in older subjects with type 2 diabetes mellitus: a randomized trial. *Hosp Pract (1995)* 2013;41:72–84
20. Bode B, Stenlöf K, Harris S, et al. Long-term efficacy and safety of canagliflozin over 104 weeks in patients aged 55–80 years with type 2 diabetes. *Diabetes Obes Metab* 2015;17:294–303
21. Forst T, Guthrie R, Goldenberg R, et al. Efficacy and safety of canagliflozin over 52 weeks in patients with type 2 diabetes on background metformin and pioglitazone. *Diabetes Obes Metab* 2014;16:467–477
22. Scherthaner G, Gross JL, Rosenstock J, et al. Canagliflozin compared with sitagliptin for patients with type 2 diabetes who do not have adequate glycemic control with metformin plus sulphonylurea: a 52-week randomized trial. *Diabetes Care* 2013;36:2508–2515
23. Qiu R, Capuano G, Meininger G. Efficacy and safety of twice-daily treatment with canagliflozin, a sodium glucose co-transporter 2 inhibitor, added on to metformin monotherapy in patients with type 2 diabetes mellitus. *J Clin Transl Endocrinol* 2014;1:54–60
24. Rosenstock J, Chuck L, González-Ortiz M, et al. Initial combination therapy with canagliflozin (CANA) plus metformin extended-release (MET XR) in drug-naïve type 2 diabetes mellitus (T2DM) (Abstract). *Diabetes* 2015;64:LB34